

# STUDY OF CORRELATION OF ECG AND ECHOCARDIOGRAPHY LOCALIZATION IN ACUTE MYOCARDIAL INFARCTION

Pothula Rama Rao<sup>1</sup>, Prathap Bingi<sup>2</sup>

<sup>1</sup>Associate Professor of General Medicine, GMC, Kadapa

<sup>2</sup>Assistant Professor of General Medicine, GMC, Kadapa

Received: 07-11-2021 / Revised: 09-12-2021 / Accepted: 20-12-2021

Corresponding author: Dr. Prathap Bingi

Conflict of interest: Nil

## Abstract:

**Background:** Acute myocardial infarction (AMI) is a major cause of morbidity and mortality, and accurate localization of the infarct is essential for early diagnosis, risk stratification, and management. Electrocardiography (ECG) and 2D echocardiography (Echo) are widely available, non-invasive tools used to identify the site and extent of myocardial damage. This study evaluates the correlation between ECG and Echo in localizing infarction in AMI patients.

**Aim:** To assess the correlation between ECG findings and echocardiographic localization of infarction in patients with acute myocardial infarction.

**Methodology:** A total of 40 AMI patients admitted to the CCU at GGH Kadapa were included based on WHO diagnostic criteria. Detailed clinical evaluation, ECG at admission, and early 2D Echocardiography were performed. Patients with prior MI, conduction defects, cardiomyopathies, or valvular and pericardial diseases were excluded. Infarct localization on ECG was compared with regional wall motion abnormalities on Echo.

**Results:** ECG showed anterior wall MI in 40%, inferior wall in 37.5%, anteroseptal in 12.5%, and anteroinferior in 10%. Echo demonstrated more detailed involvement, including extensive anterior, anteroseptal-apical, and combined inferior-RV infarctions. A strong correlation was observed for anterior wall MI, while inferior MI showed broader extension on Echo.

**Conclusion:** ECG and Echo correlate well in localizing the infarct site in AMI. Echocardiography provides greater clarity on the extent of regional wall motion abnormalities, making both modalities complementary in diagnosis and management.

**Keywords:** Myocardial infarction, ECG, Echocardiography, Infarct localization, Wall motion abnormality.

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

## Introduction

Acute myocardial infarction is one of the common diagnosis in hospitalized patients in both developed and developing countries, which is a serious complication of atherosclerotic coronary heart disease. In most patients it results from atherosclerotic occlusion of the related

vessel resulting in infarct. Myocardial infarction may occur at any age but the frequency increases progressively with age. Ischemic heart disease is thus a great killer of mankind accounting for 15% of all mortality in India [1,2]. Acute myocardial infarction is characterized by

retrosternal chest pain, radiating to ulnar aspect of left arm. In some patients particularly in elderly, acute myocardial infarction is not manifested clinically by chest pain but rather by symptoms of acute left ventricular failure [3]. It is observed that various risk factors such as age, male sex, smoking, obesity, hyperlipidemia, diabetes mellitus, hypertension, family history of IHD, type A personality, play a role in the occurrence of myocardial infarction [4]. Various methods such as QRS scoring index by ECG & left ventricular EF and wall motion abnormality by 2D Echo help in diagnosis and prognostification of myocardial infarction. These investigations are non-invasive and can be done at less advanced centres. Hence, this study is undertaken to correlate the site of infarction by ECG and 2D Echo.

**Materials & Methods:** A total of 40 patients of myocardial infarction admitted in CCU, GGH Kadapa were selected for study. Data is collected by taking a detailed history from the patients (as per the performa) particularly keeping the following points in view.

Time of onset of typical chest pain, nature of pain and associated symptoms like excessive sweating, vomiting, breathlessness, giddiness, fatigue and abdominal pain. A history of smoking, alcohol consumption, hypertension, diabetes mellitus, obesity according to BMI, personality type and family history of IHD.

A thorough clinical examination was carried out in each case with special reference to pulse, BP, CVS and

respiratory examination for the presence of any cardiac enlargement, S3 gallop, rub, murmur and basal crepitations in the lungs. Investigations like fasting lipid profile & enzymes like CPK- MB and SGOT were done. ECG is taken at the time of admission for the ECG diagnosis of myocardial infarction, the criteria consisting of ST segment elevation of  $\geq 2\text{mm}$ , 0.08 second from J point in  $\geq 2$  related electric fields, with typical evolutionary changes or presence of new pathological Q waves.

As soon as feasible, a 2D-Echo was performed by means of commercially available mechanical sector scanner. With the patient in left lateral decubitus position, multiple parasternal long axis views, short axis and apical views were taken to study regional wall motion abnormalities

#### **Inclusion Criteria:**

Patients above 20 years of age and WHO criteria for the diagnosis of acute MI are included A history of ischemic type of chest pain, Evolutionary changes on serially obtained ECG tracings and A rise and fall in serum cardiac markers.

**Exclusion Criteria:** Patients presenting with: Previous history of MI, Sub endocardial infarction, true posterior wall infarction, LVH, hemi-block, bundle branch blocks, intraventricular conduction defects and complete heart blocks. Valvular heart disease, Cardiomyopathy, Pericardial diseases viii. Congenital heart disease, Previous cardiac surgeries were excluded from the study.

#### **Results:**

**Table 1: Site of infarction in ECG**

Site of infarction	Total no. of cases	%
Anterior wall	16	40
Inferior wall	15	37.5
Anteroinferior wall	4	10
Antero septal wall	5	12.5
Total	40	100

**Table 2: Site of infarction on 2D echo in 16 patients of anterior wall MI**

Site of infarction on 2D echo	No. of cases	%
Extensive anterior wall	6	37.5
Anteroseptal and apical	5	31.25
Anterior and apical	3	18.75
Anterior and septal	2	12.5
Total	16	100

**Table 3: Site of infarction on 2D echo in 15 patients of inferior wall MI**

Site of infarction on 2D echo	No. of cases	%
Inferior wall	8	53.3
Inferior wall and RVMI	4	26.61
Inferior wall and anteroseptal	3	20
Total	15	100

### Discussion:

In our study, as shown in results in table No.1, 16 patients out of 40 patients had anterior wall myocardial infarction on ECG. Echocardiography in these patients further elaborated that 6 patients had extensive anterior wall infarction, 5 patients had antero-septal and apical wall myocardial infarction, 3 had anterior & apical, 2 had anterior & septal infarction & none of the patients showed no regional wall motion abnormality, thus elaborating the extensive anterior infarction seen on electrocardiography in great details.

15 patients, out of 40 patients had inferior wall myocardial infarction and inferior wall with right ventricle infarction on ECG. When echo was done in these patients, 8 patients had inferior wall myocardial infarction, 4 patients had inferior wall and right ventricle infarction, 3 patients had inferior wall and anterior-septal myocardial infarction and none of the patients showed no regional wall motion abnormality, again giving a more lucid interpretation.

5 patients out of 40 patients had antero-septal infarction on ECG. On echocardiographic examination in these patients, 3 patients had antero-septal myocardial infarction, 1 patients had antero-septal apical infarction, 1 patients

had antero-septal and interventricular septum infarction, thereby lending credence to the fact that echocardiography delineates ischemic changes more extensively. 4 patients out of 40 patients had antero-inferior wall myocardial infarction on ECG. When echo was done in these patients, 3 patients had global hypokinesia and one patient had anterior, inferior and lateral wall hypokinesia.

According to Penco M et al. (1996) ECG and echo cardiography showed good correlation in evaluating the infarct site, but echo showed larger extension [5]. Regarding the infarct site, a good correlation was found in anterior acute myocardial infarction but not in inferior acute myocardial infarction.

Our study is in agreement with the above findings. Shah et al. (1980) in his study agreed that ECG and echocardiography have good correlation to localize the site of infarction, but in echocardiography segmental wall motion abnormalities are frequently more extensive than on ECG and may occur in areas, apparently remote from the putatively infarcted zone [6].

In our study, this is also seen to be true. According to Scharti et al (1984) 2D echocardiography should be regarded as a supporting method to the ECG but not as an essential one in the diagnosis of acute

myocardial infarction [7]. According to Kuch (1993) ECG and 2D Echo are compatible methods but not replacable ones [8]. Mahajan Devinder Singh concluded that localization of the site of myocardial infarction on ECG correlated broadly with that seen on Echocardiography and was able to elaborate regional wall motion abnormalities in detail i.e., Echo could detect abnormalities in those areas which could not be shown by ECG [9]. Our study is in agreement with the findings of Penco M (1996) Shah (1980), Scharti et al. (1984), Kuch (1993), Izumi et al. (1995) & Mahajan Devinder Singh (2002), and electrocardiography and echocardiography have a good correlation in localizing the site of infarction but Echo was able to elaborate the site of infarction in much greater detail.

### Conclusion:

Localization of MI with ECG very much correlated with echocardiography findings in our study.

Echo was able to elaborate regional wall motion abnormalities in detail than ECG. ECG and Echo serve as two major tools for diagnosis and management of acute MI patients.

### References:

1. Park K. Park's Text Book of Preventive and Social Medicine. 17th Edn., M/s BanarsidasBhanot publishers; P-272 2002.
2. Siddharth N. Shah. API Text book of Medicine. 7th Edn. Association of Physicians of India; P-441; 2003.
3. DJ Weatherall, JGG Ledingham & DA Warrell. Oxford Text Book of Medicine. Vol.II, 3rd Edn, P – 2331; 1996.
4. Valentin Fuster et al., Hurst's. The Heart, 11th Edn. Vol. 1, P – 5, Mc Graw Hill publications, 2004.
5. Penco M, Degwanti A, Rosant G, Romani S: "Comparison of ECG and echocardiographic findings in acute myocardial infarct" *Cardiologica* 1996; 41(8): p.1751-58.
6. Shah PK., Pichler M., Berman DS., Maddohi Peter J., Singh BN. "Non – invasive identification of high risk subset of patients with acute inferior wall myocardial infarction". *Am.J. Cardiol.* 1980; 46: p.915.
7. Scharti M, Reitsch W, Muller U.J. *Kardiol* 1984; 73(1): p.56-65.
8. Kuch J., Brakstor W., Sczanieka O., Ruchalski M., Wysokinski A., Kryzm inska E.: *Kardiol Pol.* 1993; 38(1): p.21-5.
9. Mahajan Devinder Singh. "ECG and echocardiography to correlate the location of acute myocardial infarction.". *JAPI* 2002; 50: p.1494.