

Different Changes in Electrocardiogram in Cases of Acute Cerebrovascular Accidents

Pothula Rama Rao¹, Prathap Bingi²

¹Associate Professor of General Medicine, GMC, Kadapa

²Assistant Professor of General Medicine, GMC, Kadapa

Received: 28-12-2021 / Revised: 15-01-2022 / Accepted: 08-02-2022

Corresponding author: Dr. Prathap Bingi

Conflict of interest: Nil

Abstract:

Background: Acute cerebrovascular accidents (CVA) frequently present with electrocardiographic (ECG) abnormalities, which may mimic myocardial ischemia and complicate clinical interpretation. These ECG changes reflect neurogenic cardiac effects and are associated with increased morbidity and mortality.

Aim: To analyze the different ECG changes occurring in acute cerebrovascular accidents and correlate them with the type of stroke.

Methodology: This study included 40 CVA patients admitted within 72 hours to GGH Kadapa. Patients with head injury, known cardiac disease, renal/hepatic disease, or delayed presentation were excluded. All patients underwent detailed clinical evaluation, 12-lead ECG at admission and on days 3, 7, and 30, along with CT brain, cardiac biomarkers, 2D Echo, and routine investigations. ECG findings such as T-wave changes, ST-segment abnormalities, QTc prolongation, U-wave changes, and pathological Q-waves were documented and compared across ischemic stroke, intracerebral hemorrhage (ICH), and subarachnoid hemorrhage (SAH).

Results: Most patients were >60 years (70%) and male (67.5%). Ischemic stroke constituted 70%, ICH 22.5%, and SAH 7.5%. The most common ECG abnormalities were T-wave changes (42.5%), QTc prolongation (35%), ST-segment depression (12.5%), ST elevation (10%), pathological Q-waves (15%), and U-wave changes (5%). T-wave changes predominated in ischemic and hemorrhagic strokes, whereas QTc prolongation was most frequent in SAH.

Conclusion: Abnormal T-waves and QTc prolongation are the most frequent ECG findings in acute CVA. These neurogenic ECG changes often resemble myocardial ischemia, and recognizing those aids in accurate diagnosis and appropriate management.

Keywords: Cerebrovascular accident, ECG changes, QTc prolongation, T-wave abnormalities, Stroke.

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

Introduction

Stroke is defined as a "neurological deficit of cerebrovascular cause that persists beyond 24 hours or is interrupted by death within 24 hours"[1]. CVA can be classified into two major categories- ischemic and hemorrhagic strokes.[2] About 87% are ischemic

strokes and the rest being hemorrhagic in origin. Studies have demonstrated the fact that primary neurologic abnormalities may produce ECG changes without any myocardial lesion. ECG changes affecting T waves, U wave, ST segment, QT interval and

arrhythmias have been reported. These changes may resemble those of myocardial ischemia and myocardial infarction, leading to misinterpretation and delay in operative management of sub arachnoid hemorrhage [3]. There are evidences suggesting, that patients who had ECG changes following CVA had poor prognosis compared to who don't have changes in ECG. Approximately 2-6% of all CVA patients die from cardiac causes in first 3 months after ischemic stroke [4].

Materials & Methods: The present study was conducted in the department of medicine in GGH Kadapa.

Inclusion Criteria: cases of cerebrovascular accident and admitted within 72 hours and patients having ECG changes.

Exclusion Criteria:

Stroke cases which came after 72 hours, the individuals with head injury and known cardiac, hepatic and renal diseases. A total of 40 cases of cerebrovascular accident were enrolled

in the study by using the above mentioned inclusion and exclusion criteria. The diagnosis of CVA was confirmed by using following criteria [5]. Temporal profile of the clinical syndrome, evidence of focal brain damage/disease, clinical setting.

A 12 Lead Electrocardiogram was taken for all the cases within 24 hours of admission and subsequently repeated on 3rd day, 7th day and 30th day for follow up. CT scan brain was taken and all patients were subjected to investigations like Trop-i or Trop-t, 2d echocardiogram, serum electrolytes, urine analysis, blood sugar, blood urea, lipid profile and relevant serological tests. Detailed analysis and interpretation of ECG changes like sinus arrhythmia, sinus bradycardia, sinus tachycardia, abnormal Q wave, U wave, T wave, right and left ventricular hypertrophy, S-T segment elevation/depression, T wave inversion and prolonged QT interval was done.

Results:

Table 1: Distribution according to age, sex and type of CVA

	Variable	No of patients	%
Age	<40 years	2	5
	40-59 years	10	25
	>60 years	28	70
Sex	Male	27	67.5
	Female	13	32.5

Table 2: Distribution according to ECG changes

ECG changes	No of patients	%
Abnormal T wave	17	42.5
QTc prolongation	14	35
ST segment elevation	4	10
ST segment depression	5	12.5
Abnormal U wave	2	5
Pathological Q waves	6	15

Table 3: Distribution of ECG changes according to types of CVA

ECG changes	Ischemic stroke		Intracerebral hemorrhage		SAH	
	No of patients	%	No of patients	%	No of patients	%
Abnormal T waves	10	35.7	3	33.3	0	0
QTc prolongation	8	28.5	2	22.2	2	66
ST segment elevation	3	10.7	1	11.1	1	33
ST segment depression	2	7.1	2	22.2	0	0
Abnormal U wave	2	7.1	0	0	0	0
Pathological Q waves	3	10.7	1	11.1	0	0

Discussion:

In the present study we tried to study the electrocardiography findings in acute cerebrovascular diseases. For this purpose we selected 40 cases of acute cerebrovascular diseases. It was observed that majority of the cases were more than 60 years of age. Male predominance was also observed with male: female ratio of 2.07. It was seen that 70% patients were of ischemic stroke followed by intra-cerebral hemorrhage (22.5%) and sub arachnoid hemorrhage (7.5%) was observed.

The most common ECG change observed in the present study was abnormal T wave (42.5%), prolong QTc (35%). Similar findings were also reported by Goldstein DS et al, Sommargren CE et al and Villa A et al.[6,7] The higher incidence of abnormal T wave in ischemic stroke patients with cardiovascular diseases implies the influences of other factors on the T wave than the intracranial disorder in this group. Dogan et al.[8] found ischemia like ECG changes in 65% of patients, QTc interval prolongation in 26%, and arrhythmias in 44% of them.

In the study of Lindgren et al[9] transient ST,T changes were found in 54% of patients with ischemic stroke with no primary heart disease. In sub arachnoid hemorrhage patients prolong QTc interval and pathologic Q wave was observed in 50% cases each whereas ST Segment elevation and abnormal T wave was observed in 25% cases each. Similar findings were also reported by Frontera

JA et al[10], Liman T et al[11] and Mayer SA et al.[12] It was seen that in abnormal T wave (33.3%) was the most common abnormal ECG pattern in intra cerebral hemorrhage patients. It was followed by prolong QTc interval (22.2%).

Various authors have made attempt to correlate the ECG abnormalities with the location of the brain lesion but the reported results were divergent. The direct stimulation of many areas of the CNS is known to result in abnormal ECG patterns; while no relationship of the ECG changes, to the site of the bleeding aneurysm, was found by Cropp and Shuster et al,[13] Manning et al,[14] and Hunt et al.[15] In studies focused on the intracranial vascular spasm after SAH and the appearance of ECG abnormalities, Wilkins et al showed no relationship, while Stober and Kunze were able to find a correlation between cerebral arteries spasms of the left hemisphere, Twave inversion, and QT prolongation.

Conclusion: We conclude that abnormal T wave, prolong QTc interval and arrhythmia were the common ECG findings in patients of cerebrovascular accident. ECG changes in CVA are very similar to changes in myocardial ischemia and are often confused to cardiac disease. A good idea of these changes helps in effective management of CVA cases.

References:

1. Donnan GA, Fisher M, Macleod M, Davis SM (May 2008). "Stroke".

- Lancet 371 (9624): 1612–23. "Brain Basics: Preventing Stroke". National Institute of Neurological Disorders and Stroke. Retrieved 2009-10-24.
2. Cropp GJ, Manning GW. Electrocardiographic changes stimulating myocardial ischemia and interactions associated with spontaneous intracranial haemorrhage, *Circulation*, 22, 1960, 25-38.
 3. Prosser J, Mac Gregor L, R Kennedy, D Hans- Christoph, Predictors of early cardiac morbidity and mortality after ischemic stroke, *Stroke*, 38, 2007, 2295- 2302.
 4. Adams RD, Victor M, Ropper AH, Cerebrovascular diseases in principles of Neurology, USA, Me Grow Hill Health Professions Divisions, 8th Ed, 2005,660- 746.
 5. Goldstein DS. The electrocardiogram in stroke. Relationship to pathophysiological type and comparison with prior tracings. *Stroke*. 1979;10:253–9.
 6. Villa A, Bacchetta A, Milani O, Omboni E. QT interval prolongation as a predictor of early mortality in acute ischemic stroke patients. *Am J Emerg Med*. 2001;19:332–3.
 7. Dogan A, Tunc E, Ozturk M, Kerman M, Akhan G. Electrocardiographic changes in patients with ischaemic stroke and their prognostic importance. *Int J Clin Pract*. 2004;58:436–40.
 8. Lindgren A, Wohlfart A, Pahlm O, Johansson BB. Electrocardiographic changes in stroke patients without primary heart disease. *Clin Physiol*. 1994;14:223–31.
 9. Frontera JA, Parra A, Shimbo D, Fernandez A, Schmidt JM, Peter P, et al. Cardiac Arrhythmias after Subarachnoid Hemorrhage: Risk Factors and Impact on Outcome. *Cerebrovasc Dis*. 2008;26:71–8.
 10. Liman T, Endres M. Elevated troponin and ECGalterations in acute ischemic stroke and subarachnoid hemorrhage. *Nervenarzt*. 2008;79:1388–90.
 11. Mayer SA, Lin J, Homma S, Solomon RA, Lennihan L, Sherman D, et al. Myocardial injury and left ventricular performance after subarachnoid hemo rrhage. *Stroke*.1999;30:780–6.
 12. Shuster S. The electrocardiogram in subarachnoid haemorrhage. *Br Heart J*. 1960;22:316–20.
 13. Cropp GJ, Manning GW. Electrocardiographic changes stimulating myocardial ischemia and infarction associated with spontaneous intracranial hemorrhage. *Circulation*. 1960;22:25–38.
 14. Hunt D, McRae C, Zapf P. Electrocardiographic and serum enzyme changes in subarachnoid hemorrhage. *Am Heart J*. 1969;77:479–88.