

Evaluation Anemia Profile in a Tertiary Care Facility: Prospective Cross-Sectional Study

Abishek Anand¹, Sunil Kumar²

¹Assistant Professor, Department of General Medicine, BMIMS, Pawapuri, Nalanda, Bihar, India

²Senior Resident, Department of General Medicine, BMIMS, Pawapuri, Nalanda, Bihar, India

Received: 05-11-2022 / Revised: 29-11-2022 / Accepted: 28-12-2022

Corresponding author: Dr. Sunil Kumar

Conflict of interest: Nil

Abstract

Aim: The aim of the present study was to determine the clinical and laboratory profile of anaemia among patients admitted to the tertiary care facility.

Methods: This study was a hospital based prospective cross-sectional study undertaken in the Department of Medicine, BMIMS, Pawapuri, Nalanda, Bihar, India. Fifty patients with anaemia were included in the study. Duration of the study was 7 months.

Results: Easy fatiguability and generalized weakness were the most common symptoms of anemia in our study. Easy fatiguability and generalised weakness were the most common symptoms of anemia in our study. Fever was seen in 10% of cases and asymptomatic symptoms were noted in 12% patients. On systemic examination anemic murmurs on CVS examination were detected among 12 patients. Basilar crepts not attributable to other diseases were found among 3 patients. Isolated hepatomegaly was found in 8, splenomegaly in 2 and hepatosplenomegaly was found in 3 patients.

Conclusion: The most common aetiology of anemia in patients enrolled was iron deficiency, followed by anemia of chronic disease and malignancy, both hematological and non-hematological. The most commonly encountered complaints on presentation in general were subjective non-specific ones such as weakness, fatigue and lassitude whereas in females it was per vaginal bleeding.

Keywords: Anaemia, Clinical profile, Laboratory profile.

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

Introduction

Anaemia is functionally defined as insufficient Red Blood Cell (RBC) mass to adequately deliver oxygen to peripheral tissues. [1] Many references consider haemoglobin (Hb) concentrations of lower than 14g/dl in men and 12g/dl in women as lower limits of normal at sea level in industrialised world. [2] However World Health Organisation (WHO) defines the lower limit of normal for Hb concentration to be 13 g/dl and 12 g/dl in men and

women respectively at sea level. [3] Anaemia is classified into mild when Hb concentrations are between 11-12.9 g/dl in men and 11-11.9 in women. Moderate anaemia is defined as Hb of 8-10.9 in both men and women. Similarly, a patient with Hb of less than 8 is said to have severe anaemia irrespective of gender. [4]

Anaemia is an indicator of poor nutrition and poor health. It is a global public health problem affecting both developed and

developing nations. In 2002 iron deficiency anaemia was considered amongst the most important contributing factor to the global burden of disease. [5] It is assumed that 50% of cases of anaemia are due to iron deficiency. [6] The consequences of anaemia including increased maternal mortality, prematurity and decreased work productivity in adults are well documented. [6-9]

India lies partly in the tropics and partly in subtropics with extreme variations of climate. In the region where hot and humid climate prevail throughout the best part of the year, the loss of iron through sweat is appreciable. Iron is lost through sweat to the extent of 15mg per month.

Hookworm infection is endemic (prevalence 20-30% or higher) as a major cause of iron deficiency anemia, especially moderate-to-severe anemia. In populations with endemic hookworm, anthelmintic therapy should be given presumptively to anyone with severe anemia, because treatment is safe and much less expensive than diagnosing hookworm infection. [10] Improving the safety, adherence, sustainability and cost-effectiveness of iron supplementation is a continued goal for researches around the world. [11] Adults with normal intestinal function have very little risk of iron overload from dietary sources of iron. [12] However, acute intakes of more than 20 mg/kg iron from supplements or medicines can lead to gastric upset, constipation, nausea, abdominal pain, vomiting, and faintness, especially if food is not taken at the same time. [13]

The aim of the present study was to determine the clinical and laboratory profile of anaemia among patients admitted to the tertiary hospital in Bihar.

Methods

This study was a hospital based prospective cross-sectional study undertaken in the Department of Medicine, BMIMS, Pawapuri, Nalanda, Bihar, India. Fifty patients with anaemia were included in the study. Duration of the study was 7 months.

Inclusion criteria

- Patients more than or equal to 15 years of age of both sexes.
- Patients with anaemia as per WHO definition.

Exclusion criteria

Patients not willing to give informed consent.

In all the above patients thorough history was taken, general physical examination and systemic examination were done. Patients were subjected to routine blood investigations including complete blood count, peripheral smear study and serology for viral markers. Required radiological investigations were done and further studies like bone marrow examination, iron profile, vitamin B12 and folate levels were done in selected patients who did not respond to therapy started based on peripheral smear report. Stool for occult blood was done among elderly patient presenting with iron deficiency anaemia.

Results

Table 1: Symptomatology of anaemia patients

Symptoms	Number of patients (%)
Easy fatigability and generalised weakness	20 (40)
Breathlessness	10 (20)
Swelling of limbs, puffiness of face	2 (4)
Giddiness	3 (6)
Chest pain	2 (4)
Fever	5 (10)
Tinnitus	2 (4)
Asymptomatic	6 (12)

Easy fatigability and generalised weakness were the most common symptoms of anemia in our study. Fever was seen in 10% of cases and asymptomatic symptoms were noted in 12% patients.

Table 2: Signs in patients with anemia

Signs	Number of patients (%)
Pallor	50 (100)
Tachycardia	20 (40)
Platonychia/koilonychia	15 (30)
Knuckle pigmentation	10 (20)
Tachypnea	8 (16)
Pedal oedema	8 (16)
Elevated JVP	7 (14)
Icterus	3 (6)

On systemic examination murmurs on CVS examination were detected among 12 patients. Basilar crepts not attributable to other diseases were found among 3 patients. Isolated hepatomegaly was found in 8, splenomegaly in 2 and hepatosplenomegaly was found in 3 patients.

Table 3: Degree of anemia

Degree	Number of patients (%)
Mild anemia	0
Moderate anemia	10 (20)
Severe anemia	40 (80)

On laboratory examination degree of anaemia (as defined by WHO) was distributed as shown in Table 3. None of the patients admitted in the hospital had mild anaemia (defined as Hb between 11-11.9 g/dl in women and 11-12.9 g/dl in men aged 15 years or more). Moderate

anaemia (defined as Hb between 8 to 10.9 g/dl in both males and females) was seen in 20% of patients. Whereas severe anaemia (defined as Hb less than 8 g/dl in both males and females) showed highest occurrence.

Table 4: Peripheral smear study in patients with anemia

Peripheral smear	Number of patients (%)
Microcytic hypochromic anaemia	24 (48)
Dimorphic anaemia	14 (28)
Normocytic normochromic anaemia	10 (20)
Macrocytic anaemia	2 (4)

Microcytic hypochromic anaemia attributed to iron deficiency unless proved otherwise was the most common form of anemia in our study. Dimorphic anaemia was the second most common suggesting that nutritional anaemia continues to predominate in our part of world.

Discussion

Nonspecific symptoms like weakness, fatigue, are one of the commonest symptoms of presentation of anemia which can also be seen in conditions other than anemia causing diagnostic difficulties to the clinicians.

Anaemia is functionally defined as insufficient Red Blood Cell (RBC) mass to adequately deliver oxygen to peripheral

tissues. [14] Many references consider haemoglobin (Hb) concentrations of lower than 14g/dl in men and 12g/dl in women as lower limits of normal at sea level in industrialised world. [15] However World Health Organisation (WHO) defines the lower limit of normal for Hb concentration to be 13 g/dl and 12 g/dl in men and women respectively at sea level. [16]

In this study anaemia was more common among younger and middle aged persons predominantly affecting the working class of the population. Similar observations were made in a study conducted by Azad KL et al. [17] Statistically 65.1% of patients were females and rest were males depicting a female preponderance. Such female dominance was also shown in studies conducted by Alvarez-Uria G et al, and Talwelkar SR et al. [18,19]

Easy fatiguability and generalised weakness was the most common presenting symptom seen in 40% of patients and asymptomatic anaemia was detected in 6 (12%) of patients. Easy fatiguability as the predominant symptom was also noted in studies conducted by Dashratham P et al, and Gayathri BN et al. [20,21] Incidentally detected anemia constituted the second most common class. This may be explained by the fact of lack of knowledge or presence of chronic anaemia. Fever as a symptom of anaemia was also noted in study conducted by S. Selvamuthukumar. [22] Anemia causing tinnitus after ruling out neuro-otologic and other secondary causes was seen in 2 patients.

As far as signs on general physical examinations were concerned pallor was the universal finding present in 100% of patients. Such predominance of pallor as a sign on examination was noted in studies conducted by Gayathri BN et al, and Vineetha et al. [21,23]

This study noted that 80% of cases presented as severe anaemia. This may be because of the reason that mild anaemia is

neglected by people and they do not approach a doctor. Another reason may be illiteracy and lack of knowledge which makes them present to the hospital as severe anaemia cases. None of mild anaemia cases were noted in our study as they are often treated on outpatient basis and our study targeted inpatients. On peripheral smear examination microcytic hypochromic anaemia attributable to iron deficiency based on examination and observation of response to therapy was the most common cause of anaemia. Similar findings were noted by Kouli R et al, and Milman N et al. [24,25] This was followed by dimorphic anaemia as the second most common cause of anaemia. Hence nutritional anaemia continues to predominate as the most common cause of anaemia in our part of world. [26]

Conclusion

The most common aetiology of anemia in patients enrolled was iron deficiency, followed by anemia of chronic disease and malignancy, both hematological and non-hematological. The most commonly encountered complaints on presentation in general were subjective non-specific ones such as weakness, fatigue and lassitude whereas in females it was per vaginal bleeding. Most of the patients belonged to moderate anemia group. Severity of anemia was not related with total leukocyte counts and platelet counts. In spite of extensive steps taken by WHO and Government bodies in educating and treating people about the disease and the consequences of not getting treated, severe anaemia with or without failure continues as the most common mode of presentation of anaemia in medicine department of the hospital.

References

1. Means RT Jr, Glader B. Anemia general considerations. In: Glader B, Greer JP. Wintrobe's clinical hematology. 13th ed. Philadelphia:

- Lippincott Williams and Wilkins; 2014 :587.
2. Wintrobe MM. Blood of normal men and women. Erythrocyte counts, hemoglobin and volume of packed red cells of 229 individuals. Bull Johns Hopkins Hospital. 1933; 53:118-30.
 3. Khusun H, Yip R, Schultink W, Dillon DH. World Health Organization haemoglobin cut-off points for the detection of anaemia are valid for an Indonesian population. J Nutr. 1999 Sep 1;129(9):1669-74.
 4. Nutritional anaemia. Report of a WHO scientific group. World Health Org. 1968; 405:5-37.
 5. World Health Organization. The world health report 2002: reducing risks, promoting healthy life. World Health Organization; 2002.
 6. Iron deficiency anaemia: assessment, prevention, and control. A guide for programme managers. Geneva: World Health Organization; 2001.
 7. Macgregor MW. Maternal anaemia as a factor in prematurity and perinatal mortality. Scottish Med J. 1963 Apr; 8(4):134-40.
 8. Scholl TO, Hediger ML. Anemia and iron-deficiency anaemia: compilation of data on pregnancy outcome. Am J Clin Nutr. 1994 Feb 1;59(2):492S-501 S.
 9. Bothwell TH, Charlton RW. Iron Deficiency in Women: Prepared as a Report for the International Nutritional Anemia Consultative Group (INACG). Nutrition Foundation; 1981.
 10. Montresor A, Crompton DWT, Bundy DAP, et al (1998). Guidelines for the evaluation of soil-transmitted helminthiasis and schistosomiasis at community level. A guide for managers of control programmes. WHO/CTD/SIP/98.1. World Health Organization, Geneva.
 11. Gupta R, Gupta RK. Current issues for the prevention of iron deficiency anemia amongst young children. Indian Pediatr. 1992; 29:1601-3.
 12. Erdman Jr JW, Macdonald IA, Zeisel SH, editors. Present knowledge in nutrition. John Wiley & Sons; 2012 May 30.
 13. US Department of Health and Human Services. Common terminology criteria for adverse events (CTCAE) version 4.0. National Institutes of Health, National Cancer Institute. 2009 May 28;4(03).
 14. Means RT Jr, Glader B. Anemia general considerations. In: Glader B, Greer JP. Wintrobe's clinical hematology. 13th ed. Philadelphia: Lippincott Williams and Wilkins; 2014:587.
 15. Wintrobe MM. Blood of normal men and women. Erythrocyte counts, hemoglobin and volume of packed red cells of 229 individuals. Bulletin of the Johns Hopkins Hospital. 1933; 53:118-30.
 16. Khusun H, Yip R, Schultink W, Dillon DH. World Health Organization haemoglobin cut-off points for the detection of anaemia are valid for an Indonesian population. J Nutr. 1999 Sep 1;129(9):1669-74.
 17. Azad KL, Dwivedi SK. A study of profile of patients with different degrees of anemia. Int J Res Med Sci. 2017 Sep;5(9):4008-12.
 18. Alvarez-Uria G, Naik PK, Midde M, Yalla PS, Pakam R. Prevalence and severity of anaemia stratified by age and gender in rural India. Anemia. 2014 Oct;2014.
 19. Talwelkar SR, Dhruva GA, Goswami SH. Survey of different types of anaemia. Schol J Appl Med Sci, 2019;7(12):3925-8.
 20. Dasharatham P, Reddy VS. A study of etiological and clinical profile of patients with severe anaemia in a tertiary care hospital. Int J Adv Med. 2018; 5:1422-7.
 21. Gayathri BN, Rao KS. Pancytopenia: a clinico haematological study. J lab physic. 2011;3(1):15.

22. Selvamuthukumar S. A clinical study of severity of anaemia in adults attending outpatient department with regard to signs and symptoms in rural south Indian medical college. *Inter J Contempor Med Res.* 2018;5(12): L21-L4.
23. Unnikrishnan V, Dutta TK, Badhe BA, Bobby Z, Panigrahi AK. Clinico-aetiologic profile of macrocytic anemias with special reference to megaloblastic anaemia. *Ind J Hematol Blood Transfus.* 2008 Dec 1;24(4):155-65.
24. Kouli R, Pegu AK, Teronpi J, Shukla R, Bhattacharjee S, Sonowal S, et al. Study of anaemia in correlation with haematological parameters among adults in a tertiary care centre of North-East India. *Int J Med Res Prof.* 2016;2(4):84- 7.
25. Milman N. Intestinal absorption of folic acid-new physiologic and molecular aspects. *Indian J Med Res.* 2012 Nov;136(5):725.
26. Chakdoui S., Moumen A., & Guerboub A. Dyslipidemia and Diabetic Retinopathy in Moroccans Type 2 Diabetics Patients: A Cross-Sectional Study. *Journal of Medical Research and Health Sciences,* 2023; 6(3): 2471–2479.