

Prospective Study Determine Etiology, Risk Factors of Non-Traumatic Small Intestinal PerforationAsik Ikbal Mazumder¹, Faizan Ahmad², Zaid bin Afroz³¹Assistant Professor, Department of General Surgery, Silchar Medical College and Hospital, Assam, India²Senior Resident, Department of General Surgery, Silchar Medical College and Hospital, Assam, India³PG-Student, Department of General Surgery, Silchar Medical College and Hospital, Assam, India

Received: 13-08-2023 Revised: 25-09-2023 / Accepted: 21-10-2023

Corresponding author: Dr. Asik Ikbal Mazumder

Conflict of interest: Nil

Abstract**Aim:** The aim of the present study was to assess the etiology and risk factors of non-traumatic small intestinal perforation.**Material & Methods:** This prospective study was conducted in the Department of General Surgery, 50 patients were included. The patients with intestinal perforations due to trauma, mesenteric ischemia and obstruction or strangulation because of hernia, volvulus or intra- abdominal adhesions, and peptic ulcer perforations of the duodenum were excluded from the study.**Results:** Patients were of 20-30 years of age followed by 20% who belonged to 10-20 years age group. Mean age of the patients who presented with perforation peritonitis was 35.5 years that shows age was not a confounding factor. Among patients who presented with perforation peritonitis, majority (84%) of the patients were males. Only 9 out of 50 patients were females suggesting gender was not a confounding factor. Out of the 50 cases, 42% were caused by peptic ulcer, 22% by tuberculosis, 18% by typhoid, 10% by ischemic bowel disease, 2% by malignancy, 4% by intussusception, and 4% by worm infestation. The p-value cannot be determined from this table alone as it only provides the frequency distribution of the etiology of the disease. Abdominal tenderness was present in all patients, signs of peritoneal irritation were present in 38 (76%) patients. The other patients had moderate localized or generalized tenderness but relatively soft abdomen. Out of the total of 50 cases, 68% underwent primary repair, 12% underwent resection with primary anastomosis, 12% underwent resection with exteriorization of the bowel, another 6% underwent resection with anastomosis and proximal stoma, and 2% received palliative drainage.**Conclusion:** Non-traumatic small bowel perforation is a serious condition associated with significant morbidity and mortality. The etiology, clinical presentation, and outcomes can vary depending on geographical location and patient population. Prompt recognition, appropriate investigations, and early surgical intervention remain crucial in managing this condition effectively.**Keywords:** Non-Traumatic, Small Bowel Perforation, Aetiological factor, Management.

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

Introduction

Non-traumatic small bowel perforation, a rare cause of acute abdomen, is a condition that can lead to fatal complications unless timely intervention occurs. [1] Late presentation, missed diagnosis and late interventions are frequent causes of morbidity and mortality. [2] Many reasons may be counted in the etiology of non-traumatic small bowel perforation. Causes of non-traumatic perforation of the small intestine in developing countries are clearly different from those observed in developed countries. [3]. Although infectious causes, especially typhoid and tuberculosis are forefront in developing countries, Crohn's disease and malignancies are more common in developed

countries. [1,4] Patients often present late with purulent peritonitis and poor general condition. [5]

As a result, serious complications such as post-operative peritonitis caused by a leak from repaired intestine, superficial wound infection, and complete wound dehiscence are not uncommon. The causes for SBP's other than trauma and known common aetiological factors (mesenteric vascular disease, internal and external hernias, intraabdominal adhesions, inflammatory bowel diseases, and iatrogenic) are also called spontaneous or nontraumatic SBPs. [6,7] The most frequent causes include various malignancies, infections, and nonspecific inflammation. [1]

Presence of recurrent abdominal pain episodes due to underlying disease and having non-specific clinical and laboratory findings may lead to a delay in the preoperative diagnosis. Although radiological imaging procedures are helpful in diagnosis, the early diagnosis rate is low, and the majority of cases are diagnosed during laparotomy. Despite advances in surgical techniques and improvement in intensive care conditions, mortality of non-traumatic small bowel perforation is still high and can be up to 42%. [1,8]

The management of complications is particularly difficult in developing countries due to limited resources, particularly facilities for parenteral nutrition. Presentation at the hospital is generally late, and patients are in an impaired physical condition due to diffuse peritonitis. [9] This is why outcomes in the postoperative period are still poor despite developments in surgical-radiologic techniques and intensive care conditions. [9,10]

Hence the aim was to study the etiology and risk factors of non-traumatic small intestinal perforation.

Material & Methods

This prospective study was conducted in the Department of General Surgery, Silchar medical college and hospital, Assam, India for one year and 50 patients were included. The patients with intestinal perforations due to trauma, mesenteric ischemia and obstruction or strangulation because of hernia, volvulus or intra-abdominal adhesions, and peptic ulcer perforations of the duodenum were excluded from the study.

Inclusion Criteria

- Patient of all ages, both males and females
- Patient diagnosed with small bowel of perforation
- Admitted in ICU or any other department with diagnosis of Small bowel perforation

Exclusion Criteria

- Cases of esophageal, gastric and colonic perforation
- Cases of traumatic small bowel perforation.
- Cases of delayed Presentation with shock and septicaemia whose general condition did not

warrant any operative management even after resuscitate measures.

- All cases of primary peritonitis, corrosive peritonitis and post-operative peritonitis due to anastomosisleakage are excluded from the study.

History of the patients, physical examination findings, laboratory data, imaging, and operative re-ports were examined. Age, gender, complaints, duration of symptoms, comorbid disease(s), perforation location, length of stay in hospital, etiology, surgical treatment, morbidity, and mortality data were recorded. All patients were divided into two groups, survivors and non-survivors, and their features were compared.

Preoperative routine complete blood count and biochemical tests were performed in all patients. Initial imaging modalities were pulmonary and abdominal x-ray and abdominal ultrasound (USG). Abdominal computed tomography (CT) was performed in patients in whom direct abdominal X-ray and USG examination were normal. Patients who were thought to have acute abdomen according to physical examination, laboratory, and radiological findings underwent surgery after initial resuscitation. A prophylactic broad-spectrum antibiotic therapy was started for all cases preoperatively. Perforation was managed by either primary suturing or resection and anastomosis with or without ileostomy. All patients received postoperative antibiotic therapy as prophylactic or therapeutic according to the results of cultures. Tissue biopsy was performed in all patients and a segment of diseased tissue or lymph node was sent for culture for mycobacterium in suspected cases.

Statistical Analysis

Data so collected was tabulated in an excel sheet, under the guidance of statistician. The means and standard deviations of the measurements per group were used for statistical analysis (SPSS 22.00 for windows; SPSS Inc, Chicago, USA). Difference between two groups was determined using chi square test and the level of significance was set at $p < 0.05$.

Results

Table 1: Demographic data

Age (years)	Frequency (%)
10-20	10 (20)
20-30	16 (32)
30-40	7 (14)
40-50	7 (14)
50-60	3 (6)
>60	7 (14)
Mean \pm SD	35.5\pm17.3
Gender	

Female	9 (18)
Male	41 (82)

Patients were of 20-30 years of age followed by 20% who belonged to 10-20 years age group. Mean age of the patients who presented with perforation peritonitis was 35.5 years that shows age was not a confounding factor. Among patients who presented with perforation peritonitis, majority (84%) of the patients were males. Only 9 out of 50 patients were females suggesting gender was not a confounding factor.

Table 2: Etiology among the study subjects

Etiology	N=50	%
Peptic Ulcer	21	42
Tuberculosis	11	22
Typhoid	9	18
Ischaemic bowel disease	5	10
Malignancy	1	2
Intussusception	2	4
Worm infestation	1	4

Out of the 50 cases, 42% were caused by peptic ulcer, 22% by tuberculosis, 18% by typhoid, 10% by ischemic bowel disease, 2% by malignancy, 4% by intussusception, and 4% by worm infestation. The p-value cannot be determined from this table alone as it only provides the frequency distribution of the etiology of the disease.

Table 3: Signs and symptoms

Sign/Symptom	n (%)
Abdominal pain	50 (100)
Nausea/vomiting	38 (76)
Signs of peritoneal irritation	36 (72)
Abdominal distention	30 (60)
Constipation	5 (10)

Abdominal tenderness was present in all patients, signs of peritoneal irritation were present in 38 (76%) patients. The other patients had moderate localized or generalized tenderness but relatively soft abdomen.

Table 4: Types of surgical procedure performed

Procedure	N	%
Primary repair	34	68
Resection and primary anastomosis	6	12
Resection and exteriorisation of bowel	6	12
Resection and anastomosis with proximal stoma	3	6
Palliative drainage	1	2
Total	50	100

Out of the total of 50 cases, 68% underwent primary repair, 12% underwent resection with primary anastomosis, 12% underwent resection with exteriorization of the bowel, another 6% underwent resection with anastomosis and proximal stoma, and 2% received palliative drainage.

Discussion

Hollow viscus perforation leading to peritonitis is a common emergency faced by a general surgeon.¹¹ Late presentation, missed diagnosis and late interventions are frequent causes of morbidity and mortality. [12] Now a day's inadvertent use of NSAIDs and analgesics available over the counter forms one of the most common risk factors. [13] Perforation of stomach, duodenum and small bowel forms considerable proportion of emergency than colonic perforations. [14,15] Perforation of large bowel represents major surgical challenge to the clinician, because it is rapidly fatal condition, death

being caused by sepsis from peritoneal contamination by various aerobic and anaerobic pathogens. [16,17] Surgery is mainstay in management of hollow viscus perforations. High index of suspicion is essential to diagnose hollow viscus perforations, because diagnostic delay may result in significant morbidity and mortality.

Patients were of 20-30 years of age followed by 20% who belonged to 10-20 years age group. Mean age of the patients who presented with perforation peritonitis was 35.5 years that shows age was not a confounding factor. Among patients who presented with perforation peritonitis, majority (84%) of the patients were males. Only 9 out of 50 patients were females suggesting gender was not a confounding factor. Similar studies have reported a peak incidence of small bowel perforation in the 31-40 age group and a higher prevalence among males, which aligns with the findings of this study. [18] Abdominal pain was reported in all subjects, while abdominal distension and obstipation were present

in 86% and 72% of the cases, respectively. Vomiting and fever were observed in 58% and 46% of the subjects, respectively. Other studies have also noted abdominal pain as the predominant symptom, followed by constipation and fever. [19]

Out of the 50 cases, 42% were caused by peptic ulcer, 22% by tuberculosis, 18% by typhoid, 10% by ischemic bowel disease, 2% by malignancy, 4% by intussusception, and 4% by worm infestation. Similar findings have been reported in Kumar VB et al [20] and Rao et al study. [21] Abdominal tenderness was present in all patients, signs of peritoneal irritation were present in 38 (76%) patients. The other patients had moderate localized or generalized tenderness but relatively soft abdomen. These findings are consistent with previous studies where abdominal pain was the most common symptom, often accompanied by distention, constipation, and fever. [22]

Out of the total of 50 cases, 68% underwent primary repair, 12% underwent resection with primary anastomosis, 12% underwent resection with exteriorization of the bowel, another 6% underwent resection with anastomosis and proximal stoma, and 2% received palliative drainage. Etiology and the degree of peritoneal contamination should be considered in the decision of which surgical procedure to repair the perforated segments. Primary suture is recommended in patients with single perforation and no generalized peritonitis, though resection and anastomosis is preferred in patients with multiple perforations. [1,23] Because the prognosis is associated with the degree of peritonitis, ileostomy is preferred in patients with generalized peritonitis. [23] Because the degree of peritonitis is usually advanced due to the delay in diagnosis, ileostomy is required in most patients. It should be noted that anastomosis leakage rates might be high in patients with tuberculosis or lymphoma on whom primary closure or resection-anastomosis without ileostomy were performed.

The primary treatment modality for small bowel perforation in this study was surgical intervention. Exploratory laparotomy was performed in all patients, and the site of perforation was identified and repaired. The choice of surgical technique depended on the location and size of the perforation, with primary closure being the most common approach. In cases of extensive bowel involvement or contamination, bowel resection with anastomosis or stoma creation was performed. In some instances, additional procedures such as peritoneal lavage or omental patching were required. [24]

Conclusion

Non-traumatic small bowel perforation is a serious condition associated with significant morbidity and mortality. The etiology, clinical presentation, and outcomes can vary depending on geographical location and patient population. Prompt recognition, appropriate investigations, and early surgical intervention remain crucial in managing this condition effectively. Further studies are needed to explore preventive strategies and optimize treatment approaches for better patient outcomes.

References

1. Eid HO, Hefny AF, Joshi S, Abu-Zidan FM. Non-traumatic perforation of the small bowel. *Afr Health Sci* 2008; 8:36-39.
2. Dhikav V, Singh S, Pande S, Chawla A, Anand KS. Non-steroidal drug-induced gastrointestinal toxicity: mechanisms and management. *J Indian Acad Clin Med.* 2003 Oct;4(4):315-22.
3. Sharma L, Gupta S, Soin AS, Sikora S, Kapoor V. Generalized peritonitis in India. *The Tropical Spectrum. Jpn J Surg.* 1991; 21: 272-277.
4. Chaikof EL. Nontraumatic perforation of the small bowel. *Am J Surg* 1987; 153:355-358.
5. Ersumo T, W/Meskel Y, Kotisso B. Perforated peptic ulcer in Tikur Anbessa Hospital; a review of 74 cases. *Ethiop Med J.* 2005; 43:9-13.
6. Mastboom WJ, Kuypers HH, Schoots FJ, Wobbes T. Small-bowel perforation complicating the open treatment of generalized peritonitis. *Archives of Surgery.* 1989 Jun 1;124(6):689-92.
7. Mischinger HJ, Berger A, Kronberger L, Fellbaum C. Spontaneous small bowel perforation. A rare cause of acute abdomen. *Acta chirurgica scandinavica.* 1989 Nov 1;155(11-12):593-9.
8. Akyildiz HY, Akcan AC, Sözüer E, et al. Unusual causes of intestinal perforation and their surgical treatment. *Ulus Travma Acil Cerrahi Derg* 2009;15:579-583
9. Leijonmarck CE, Fenyö G, Räf L. Nontraumatic perforation of the small intestine. *Acta chirurgica scandinavica.* 1984 Jan 1;150(5):405-11.
10. Mastboom WJ, Kuypers HH, Schoots FJ, Wobbes T. Small-bowel perforation complicating the open treatment of generalized peritonitis. *Archives of Surgery.* 1989 Jun 1;124(6):689-92.
11. Langel JT, Mulvihill SJ. Gastrointestinal perforation and the acute abdomen. *Medical Clinics of North America.* 2008 May 1;92(3): 599-625.
12. Dhikav V, Singh S, Pande S, Chawla A, Anand KS. Non-steroidal drug-induced

- gastrointestinal toxicity: mechanisms and management. *J Indian Acad Clin Med.* 2003 Oct;4(4):315-22.
13. Inderbir S. Oesophagus, stomach and intestine, textbook of human histology. 2nd ed. New Delhi; Jaypee Brothers Medical Publisher; 1994:212-25.
 14. Janikalpesh A, Saxena AK. Management of large sized duodenal ulcer perforation by omental plugging, a new technique, a prospective study of 100 patients. *Ind J Surg.* 2000;62(2):134-8.
 15. Jhobta RS, Attri AK, Kaushik R, Sharma R, Jhobta A. Spectrum of perforation peritonitis in India-review of 504 consecutive cases. *World journal of Emergency surgery.* 2006 Dec;1(1):1-4.
 16. Kellog LC. A treatise on peptic perforations. *Surgery.* 1939; 6:524-30.
 17. Dhanapal V, Selvam K. Clinical study and management of hollow viscus perforation of abdomen. *Int Surg J.* 2017;4(5):1773.
 18. Sørdeide JA, Viste A. Esophagus, stomach, and duodenum. In: Brunicaardi FC, Andersen DK, Billiar TR, editors. *Schwartz's principles of surgery.* 11th ed. McGraw-Hill Education; 2019. p. 1101-46.
 19. Zafar SN, Obirize A, Adesibikan B. Surgical outcomes of peptic ulcer perforation in a national database. *Am J Surg.* 2014;207(5):721-6.
 20. Kumar VB, Mathew AS. Clinical study of abdominal hollow visceral perforation, non-traumatic. *J Evol Med Dent Sci.* 2014;3(30):8366-71.
 21. Rao M, Samee AA, Khan SM. Hollow viscus perforation: A retro spectrum study. *Int J Recent Sci Res.* 2015;6(3):3250-4.
 22. Lohsiriwat V. Tuberculous enteritis: a clinical and endoscopic analysis in 30 patients. *Indian J Gastroenterol.* 2006;25(6):284-8.
 23. Ara C, Sogutlu G, Yildiz R, Kocak O, Isik B, Yilmaz S, Kirimlioglu V. Spontaneous small bowel perforations due to intestinal tuberculosis should not be repaired by simple closure. *Journal of gastrointestinal surgery.* 2005 Apr 1;9(4):514-7.
 24. Khanna AK, Tiwary SK, Kumar P. Non-traumatic perforation of the small intestine: clinical profile and surgical experience from a tertiary care center. *Indian J Surg.* 2010;72(6):434-8.