

An Observational Assessment of the Profile of Sternalis Muscle in Cadavers: A Morphometric Study

Nimisha Madhu¹, Praveen Kumar², Rajendra Prasad³

¹Tutor, Department of Anatomy, Anugrah Narayan Magadh Medical College, Gaya, Bihar, India

²Tutor, Department of Anatomy, Anugrah Narayan Magadh Medical College, Gaya, Bihar, India

³HOD, Department of Anatomy, Anugrah Narayan Magadh Medical College, Gaya, Bihar, India

Received: 11-08-2023 / Revised: 16-09-2023 / Accepted: 21-10-2023

Corresponding Author: Dr. Praveen Kumar

Conflict of interest: Nil

Abstract

Aim: The aim of the present study was to assess the profile of sternalis muscle in cadavers.

Methods: The present study was conducted in the Department of Anatomy, ANMMC, Gaya Bihar for two years and 70 adult cadavers (55 males and 15 females) embalmed in 10% formalin were dissected in the dissection hall.

Results: Muscle was bifurcating into medial and lateral tendinous laminae after 8 cm from the distal fleshy end. The length of medial lamina was 4.5 cm which was attached to sternal angle whereas the lateral lamina was 3 cm long while was reaching to the medial end of second intercostal space merging with the deep fascia covering pectoralis major muscle. Medial border of the muscle with its medial lamina was making an angle of 350° with the midline. On left side: Distal end was fleshy while its fibres at proximal ends were tendinous. Its length was 12 cm from the fifth rib to the sternal angle. Medial border of the muscle near the sternal angle was making angle of 300° with the midline passing through centre of sternum.

Conclusion: The early detection is essential in regular mammogram screening to differentiate it from the malignant lesions. This muscle is a matter of interest for anatomists, radiologists and surgeons for doing surgeries on anterior chest wall.

Keywords: Mammography, lamina, Pectoralis major muscle, Sternalis muscle, Sternocleidomastoid muscle, Sternum

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

Introduction

The sternalis muscle is an anatomical variant of the anterior thoracic region musculature well documented and familiar to anatomists but quite unknown among clinicians and radiologists. [1] It lies superficially and perpendicular to the pectoralis major muscle and parallel to the sternum. Many more terms have been used in the literature to describe sternalis muscle such as "parasternalis" and "rectus sterni" muscle. [2,3] The muscle usually arises from the upper sternum and the infraclavicular region and can display variable insertion points such as the pectoral fascia, lower ribs, costal cartilages, rectus abdominis muscle sheath or the abdominal external oblique muscle aponeurosis. [4,5] However, there is a great variation in height (4.8±1.97 cm), width (15.1±6.84 mm), and thickness (3±0.91 mm). [6]

The sternalis muscle is a rare (strap-like) anterior chest wall muscle variant with fibers near-parallel to the sternum and perpendicular to the pectoris major. Though inadequately described in standard anatomical texts, this muscle is familiar to trained

anatomists. [7] However, it is relatively unknown to radiologists, surgeons, and other clinicians until recently. [8] Its rarity (roughly 5%-10% of the population) [9], variable morphology [10] and poorly understood embryology [7] render the muscle rather perplexing to anatomists, radiologists, and surgeons alike.

The appearance of this muscle is in the form of superficial vertical strip, which is sternal or parasternal in position. CT and MR imaging are diagnostic when they show the longitudinal extent of the muscle, which lies anterior to the medial margin of the pectoralis major muscle. If clinicians are not aware of its occasional presence, this muscle can be confused with an abnormal mass. In radiotherapy and surgery, lack of knowledge on Rectus Sternalis can cause negative changes in prognosis of patient. Sufficient knowledge on this muscle may help in making a better diagnosis and it can decrease complications. Cadaveric studies have shown the equal presence of this muscle in approximately 5-8% of both males and females. Variations in its

incidence have been reported in different ethnic groups. Rectus Sternalis is twice as often unilateral as it is bilateral. [11] Many authors assumed the Sternalis muscle to be derived from neighbouring muscles, such as pectoralis major [12], rectus abdominis, sternocleidomastoid, panniculus carnosus and external oblique. [13]

The aim of the present study was to assess the profile of sternalis muscle in cadavers.

Materials and Methods

The present study was conducted in the Department of Anatomy, ANMMC, Gaya, Bihar, India, for two years and 70 adult cadavers (55 males and 15

females) embalmed in 10% formalin were dissected in the dissection hall.

After the removal of subcutaneous tissue, the region was cleaned with the help of blunt forceps. The region was looked for presence of sternalis muscle or any other additional muscle. The region was further dissected to find out the nerve supply of muscle and whether muscle fibres were merging with any other muscle. Length of the muscle was measured with the help of vernier caliper (minimum count 01 mm). The angle of sternalis with midline was measured with Goniometer (minimum count 1°).

Results

Table 1: Morphometric profile of sternalis muscle

Features	Right side	Left side
Extent	Fifth rib to sternal angle	Fifth rib to sternal angle
Distal end	Fleshy and 3 cm wide	Fleshy and 3 cm wide
Proximal end	Tendinous with two laminae	Tendinous with single lamina
Angle of muscle with vertical plane	35°	32°
Length of muscle	8 cm from lower end till bifurcation of tendons. Lateral lamina- 3 cm Medial lamina- 4.5 cm	12 cm

Muscle was bifurcating into medial and lateral tendinous laminae after 8 cm from the distal fleshy end. The length of medial lamina was 4.5 cm which was attached to sternal angle whereas the lateral lamina was 3 cm long while was reaching to the medial end of second intercostal space merging with the deep fascia covering pectoralis major muscle. Medial border of the muscle with its medial lamina was making an angle of 35° with the midline. On left side: Distal end was fleshy while its fibres at proximal ends were tendinous. Its length was 12 cm from the fifth rib to the sternal angle. Medial border of the muscle near the sternal angle was making angle of 30° with the midline passing through centre of sternum.

Discussion

Sternalis muscle on the anterior chest wall is occasionally detected in humans. Its appearance is in the form of superficial strip of muscle, which is sternal or parasternal in position. [14] It is also known as episternalis, pre-sternalis, rectus thoracic or rectus sterni. [15] According to Turner, it was first identified by Cabrol in 1604 as a longitudinal band like muscle but its proper description was first made by Du Puy in 1726. [16] Variation in its incidence has been reported in different ethnic groups. Incidence of this muscle varies as low as 1% in Taiwanese to 11% in Asian population. [17] It is found slightly more in females than males. [18]

Developmentally, Sternalis is a part of a ventral, longitudinal column of muscles which arise at the

ventral tips of hypomere. This muscle is represented by the rectus abdominis muscle in the abdominal region and by infrahyoid muscle in the cervical region. In the thoracic region, this layer usually disappears but occasionally remains as sternalis muscle.¹³ It could be the separated fibres of pectoralis major muscle, as indicated by its nerve supply from twigs of pectoral nerves. Depending on its development it may get its nerve supply from lateral or medial pectoral nerves. During the dissection, attempts were made to find out the innervation of this muscle. It was found that it was not supplied by nerve supplying pectoralis major and pectoralis minor muscles. An assumption was made that it would have been supplied by anterior cutaneous branch of the neighboring intercostal nerves. It has been suggested that the sternalis could be an upward prolongation of the rectus abdominis or a downward extension of the sternocleidomastoid. [16,19] In this case, the sternalis muscle appeared to be continuation of the rectus abdominis muscle. Muscle was bifurcating into medial and lateral tendinous laminae after 8 cm from the distal fleshy end. The length of medial lamina was 4.5 cm which was attached to sternal angle whereas the lateral lamina was 3 cm long while was reaching to the medial end of second intercostal space merging with the deep fascia covering pectoralis major muscle. Medial border of the muscle with its medial lamina was making an angle of 35° with the midline. On left side: Distal end was fleshy while its fibres at proximal ends were tendinous. Its length was 12 cm from the fifth rib to

the sternal angle. Medial border of the muscle near the sternal angle was making angle of 300° with the midline passing through centre of sternum.

Occurrence of Rectus Sternalis is rare but it is not the rarest and it has racial and regional variations. Sarikcioglu L et al [20] reported its incidence to be 4-7% in white population, 8.4% in black population, 11% in Asian population, 1% in Taiwanese population and 9.3% in Turkish population. In Indian population, its incidence is 5-8% equal in both genders. Mehta V et al [21] discussed a more common unilateral occurrence of this muscle and lack of acquaintance of the clinicians with this muscle variant. An incidence of 4–8% has been found earlier in Indian subjects. [22,23] Reported size of Rectus Sternalis was different in different cases. In cases with a partial or complete congenital absence of Pectoralis Major muscle, a significant size of Rectus Sternalis has been observed. Muscles were wider at their caudal ends in case reports of Mehta V et al [21] (2.2 cm) and Kulkarni DU et al [24] (5.2 cm), these findings are similar to present finding. In present study, upper end of muscle was in the form of 2-3 slips and it was tendinous, while the lower end was wider (4.2 cm) and fleshy.

Conclusion

The early detection is essential in regular mammogram screening to differentiate it from the malignant lesions. This muscle is a matter of interest for anatomists, radiologists and surgeons for doing surgeries on anterior chest wall.

References

- Bailey PM, Tzarnas CD. The sternalis muscle: a normal finding encountered during breast surgery. *Plast Reconstr Surg*. 1999 Apr;10 (4) :1189-90.
- Arráez-Aybar LA, Sobrado-Perez J, Merida-Velasco JR. Left musculus sternalis. *Clin Anat*. 2003 Jul;16(4):350-4.
- Loukas M, Bowers M, Hullett J: Sternalis muscle: a mystery still. *Folia Morphol (Warsz)*. 2004, 63: 147-149.
- Raikos A, Paraskevas GK, Yusuf F, Kordali G, Ioannidis O, Brand-Saberi B: Sternalis muscle A new crossed subtype, classification, and surgical applications. *Ann Plast Surg*. 2011.
- Georgiev GP, JeleV L, Ovtsharoff VA: On the clinical significance of the sternalis muscle. *Folia Med (Plovdiv)*. 2009, 51: 53-56.
- Lee BY, Byun JY, Kim HH, Kim HS, Cho SM, Lee KH, Song KS, Kim BS, Lee JM. The sternalis muscles: incidence and imaging findings on MDCT. *Journal of Thoracic Imaging*. 2006 Aug 1;21(3):179-83.
- Sarikcioglu L, Demirel BM, Nurettin OĞ, Yaşar UÇ. Three sternalis muscles associated with abnormal attachments of the pectoralis major muscle. *Anatomy*. 2008 Jan 10;2(1):67-71.
- Raikos A, Paraskevas GK, Tzika M, Faustmann P, Triaridis S, Kordali P, Kitsoulis P, Brand-Saberi B. Sternalis muscle: an underestimated anterior chest wall anatomical variant. *Journal of cardiothoracic surgery*. 2011 Dec;6(1):1-4.
- Lee BY, Byun JY, Kim HH, Kim HS, Cho SM, Lee KH, Song KS, Kim BS, Lee JM. The sternalis muscles: incidence and imaging findings on MDCT. *Journal of Thoracic Imaging*. 2006 Aug 1;21(3):179-83.
- JeleV L, Georgiev G, Surchev L. The sternalis muscle in the Bulgarian population: classification of sternalis. *The Journal of Anatomy*. 2001 Sep;199(3):359-63.
- Loukas M, Bowers M, Hullett J. Sternalis muscle: a mystery still. *Folia Morphol (Warsz)*. 2004 May;63(2):147-9.
- Clemente CD. Gray's anatomy. In: *Muscle and fasciae*. 30th ed. Philadelphia: Lea and Febiger ; 1985.
- Sadler TW. *Langmans Medical Embryology*. 9th ed. Baltimore: Lippincott Williams & Wilkins; 2004. pp. 199–209.
- Raikos A, Paraskevas GK, Yusuf F, Kordali G , Ioannidis O, Brand saberi B. Sternalis muscle a new crossed subtype, classification and surgical applications. *Ann Plast Surg*. 2011;67(6):646-48.
- Georgiev GP, JeleV L, Ovtsharoff VA. On the clinical significance of the sternalis muscle. *Folia medica*. 2009;51(3):53-56.
- Turner WM. On the musculus sternalis. *J Anat Physiol*. 1867;1(2):246-53.
- Jeng H, Su SJ. The sternalis muscle: an uncommon anatomical variant among Taiwanese. *J anat*. 1998;193:359-63.
- Scott corner CE, Al Jurf AS. The sternalis muscle. *Clin anat*. 2002;15:67-69.
- Stranding S. *Gray's Anatomy- The Anatomical basis of clinical practice*. 40th ed. Edinburgh: Churchill livingstone; 2008.p.808.
- Sarikcioglu L, Demirel BM, Nurettin OĞ, Yaşar UÇ. Three sternalis muscles associated with abnormal attachments of the pectoralis major muscle. *Anatomy*. 2008 Jan 10;2(1):67-71.
- Mehta V, Arora J, Yadav Y, Suri RK, Rath G. Rectus thoracis bifurcatus: a new variant in the anterior chest wall musculature. *Rom J Morphol Embryol*. 2010;51(4):799-801.
- Mishra BD. The Sternalis muscle. *J Anat Soc India*. 1954;3:47–48.
- Shah AC. The Sternalis muscle. *Indian J Med Sci*. 1968;22(1):46–47.
- Kulkarni DU, Kulkarni UK. Unilateral rectus Sternalis muscle- a case report. *Al Ameen J Med Sci*. 2010;3(2):169–71.