

**Cross Sectional Study Assessing Visual Outcome Following Management of Traumatic Cataract: An Observational Study****Kumar Parmanand<sup>1</sup>, Mrityunjay Kumar<sup>2</sup>, Deepak Kumar Sinha<sup>3</sup>**<sup>1</sup>Senior Consultant, Department of Ophthalmology, Sunaina Netralaya, Biharsharif, Nalanda, Bihar, India<sup>2</sup>Senior Resident, Department of Ophthalmology, Government Medical College, Bettiah, Bihar, India<sup>3</sup>Senior resident, Department of Ophthalmology, SKMCH, Muzaffarpur, Bihar, India

Received: 05-07-2023 Revised: 12-08-2023 / Accepted: 21-09-2023

Corresponding author: Dr. Mrityunjay Kumar

Conflict of interest: Nil

**Abstract****Aim:** The aim of the present study was to assess the visual outcome following management of traumatic cataract.**Methods:** The present study was a cross sectional study carried out in Ophthalmology outpatient Department for the period of one year. Study population was all the patients presenting with cataract. 50 cases from these cases of traumatic cataract were included for this study as the remaining patients did not fulfill the inclusion criteria.**Results:** The age group ranged from 18-62 years. More number of cases was found in the age group of 21-40 years (56%). Out of 50 cases, 35 were males and 15 were females. Out of 50 cases of traumatic cataract, 15 were blunt and 35 were penetrating trauma. Pre-operative visual acuity was recorded in all cases. In 15 cases (30%), the preoperative visual acuity was PL/PR. In 22 cases (44%), the preoperative visual acuity was perception of hand movements. In 11 cases (22%), the preoperative visual acuity was 3/60 to counting fingers ½ meters. Visual acuity of the uninjured eye was recorded in all the patients and was found to be within the normal limit. Out of 50 cases of traumatic cataract, 47 cases underwent ECCE (Small Incision Cataract Surgery) with PCIOL implantation, 1 case underwent small incision cataract surgery with PCIOL implantation with corneal tear repair, 1 case underwent small incision cataract surgery with PCIOL implantation with anterior vitrectomy, 1 case underwent small incision cataract surgery with aphakia. Out of 50 cases, final visual acuity of 6/6 to 6/18 was seen in 15 (30%) cases. Final visual acuity of less than 6/18 to 6/60 was seen in 30 patients (60%). Five patients (10%) had visual acuity less than 3/60.**Conclusion:** Traumatic cataract is a serious visually challenging sequel of trauma. In cases of traumatic cataract, to have a better visual outcome after surgery, early diagnosis followed by proper management plays an important role. Therefore stress has to be given on awareness of the public and ocular safety measures should be taken at work places to prevent the ocular hazards associated with the ocular trauma. In addition, early reporting and adequate follow up especially in cases of children needs to be emphasized.**Keywords:** visual outcome, management, traumatic cataract

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

**Introduction**

Ocular trauma is an important cause of mono-ocular blindness in the world. 40% is related to it. [1,2] Traumatic cataract that may occur after various types of ocular insult is a serious visually challenging sequel of trauma. Domestic injuries are the commonest mode of injury in children. In adults sports and work related eye injuries are common. Traumatic cataract that develops during the early stages of life besides the visual impairment it can also cause amblyopia. Profound visual impairment can result due to stimulus deprivation during the early stage.

For proper management a detailed history and a pre-operative examination is a must before performing surgery in a case of traumatic cataract. Intraocular

foreign bodies and open globe injuries should be ruled out before surgery. [3] The final visual outcomes depend on the type of trauma, extend of lenticular involvement and associated damage to the ocular structures. In adults the time of intervention of cataract surgery should be carried out and completed within a year and within 6 months in children. [4] Surgery for traumatic cataract can be primary or secondary. When the lens is fragmentized, swollen causing a pupillary block or lens opacity blocking the view of posterior segment, primary cataract removal is done. However secondary cataract removal is more beneficial because of improved visibility, proper intraocular

lens power calculation, and there is less chances of post operative inflammation. [5]

Cataract is the commonest complication following ocular injury.<sup>5</sup> Traumatic cataract results most commonly from either penetrating injuries from sharp objects like stick or thorn with direct injury to lens or through blunt trauma by objects like stone, cricket ball etc. Rarely, it can occur from electrical shock, ionizing radiation or infra-red rays (glass blower's cataract). [6] Traumatic cataract following a perforating injury may be localized cataract, rosette cataract, intumescent cataract or lacerated cataract. Blunt trauma leads to concussion type of cataract due to coup or contre coup ocular injury. The lenticular opacity may be vossius ring, localized or diffuse type. [7-10]

The pathophysiology of traumatic cataract is believed to involve direct rupture of capsule or coup, countercoup and equatorial expansion due to hydraulic forces transferring the energy of trauma to the opposite side of the eye. It can be accompanied by anterior and posterior segment abnormalities depending on the force of trauma and depth of globe penetration. [11,12] Traumatic cataract poses a significant medical and surgical challenge to an ophthalmologist. A detailed history, careful examination and a clear management plan can simplify these cases and provide the best possible visual outcome. [13,14] Traumatic cataract can be treated by primary removal. The benefits of secondary removal are improved visibility and less chance of postoperative complications. [15]

The aim of the present study was to assess the visual outcome following management of traumatic cataract.

### Materials and Methods

The present study was a cross sectional study carried out in Ophthalmology outpatient Department of Sunaina Netralaya, Bihar Sharif, Nalanda, Bihar, India for the period of one year. Study population was all the patients presenting with cataract. 50 cases from these cases of traumatic cataract were included for this study as the remaining patients did not fulfill the inclusion criteria.

#### Inclusion Criteria:

All the patients presenting with cataract in the Ophthalmology outpatient department

Patients willing to participate in the study

#### Exclusion Criteria:

1. Patients presenting with any pathology other than cataract.
2. Ocular injuries without cataract.

Study was approved by ethical committee of the institute. A valid written consent was taken from the patients after explaining study to them.

All 50 cases were evaluated, with respect to a detailed history, regarding type of trauma, duration between trauma and presentation, associated ocular injury, intra-operative and post-operative complications. The visual prognosis after surgery was noted. All patients with traumatic cataract presenting at Government Medical College, Bettiah, Bihar, India fulfilling sampling criteria were selected for surgery. Detailed Systemic examination was done. Each patient was subjected to detailed examination and investigation needed. Routine blood analysis Lacrimal syringing, Intraocular pressure assessment were done. Radiological investigation, B-scan ultrasonography to rule out intraocular foreign body, vitreous hemorrhage, RD. Ocular examination included Torch light examination, slit lamp biomicroscopic examination and indirect ophthalmoscopy. Preoperative visual acuity was recorded in both eyes. Keratometry and A-scan biometry was done for intraocular power calculation, but in case of corneal scarring the power of other eye was calculated. Topical antibiotics and NSAIDS were administered hourly the day before surgery. Antiglaucoma medication was given in cases associated with raised IOP. Pupil was dilated with 0.5% tropicamide with 10% phenylephrine until full dilatation was attained. In cases associated with inflammation it was controlled with topical steroids and antibiotics before surgery. Peribulbar block was given which consisted of a mixture of 2% lignocaine with adrenaline with hyaluronidase 10 units per ml. All cases were operated under operating microscope with co-axial illumination. Eye was painted and draped. Superior rectus bridles suture was taken. Fornix based conjunctival flap was raised. A 6.5-7.00 mm partial thickness sclerocorneal tunnel was made. Side port entry was made. CCC was done. Hydro dissection was done where required. Nucleus was prolapsed in AC where possible and was expressed with viscoelastics, where it was imbedded lens matter it was directly aspirated. Epinucleus and residual cortex was aspirated with simcoe cannula. A 6mm optic single piece PMMA lens was placed in the bag, sulcus as per the case. Wound hydration was done and wound was checked for its apposition. Subconjunctival injection dexamethasone and gentamycin was given and eye was padded and bandaged. Post operatively, Systemic analgesics were administered along with systemic antibiotics. Next morning eye was examined under slit lamp to look for any postoperative complications. Patients were started on topical antibiotic steroids, NSAIDS and mydriatics eye drops. Antiglaucoma medications were given in selected cases. Patients were discharged with postoperative instructions regarding the medications and other measures. Patient were

instructed to come for follow-up after 1 week and after 6 weeks. Visual acuity was assessed with snellens chart at each postoperative visit along with slit lamp examination. Antibiotic steroid drops were gradually tapered. Refraction was done at 6th weeks postoperatively and glasses were prescribed based on patients refractive status. The clinical data of

each patient was collected in the proforma for analysis of the study.

Data was analysed with SPSS version 22.

## Results

**Table 1: Distribution of traumatic Cataract patients according to age and sex**

Age group (years)	Male	Female	Total	Percentage
11-20	4	2	6	12
21-30	11	3	14	28
31-40	9	5	14	28
41-50	9	3	12	24
>50	2	2	4	8
Total	35	15	50	100

The age group ranged from 18-62 years. More number of cases was found in the age group of 21-40 years (56%). Out of 50 cases, 35 were males and 15 were females.

**Table 2: Distribution of traumatic Cataract patients according to type of trauma**

Type of Trauma	N	%
Blunt	15	30
Penetrating	35	70
Total	50	100

Out of 50 cases of traumatic cataract, 15 were blunt and 35 were penetrating trauma.

**Table 3: Distribution of traumatic Cataract patients according to pre-operative visual acuity**

Visual acuity	N	%
3/60	4	8
CF ½mt	2	4
CF1M	2	4
CF2M	3	6
HM	22	44
PL+	2	4
PL+ PR+	15	30

Pre-operative visual acuity was recorded in all cases. In 15 cases (30%), the preoperative visual acuity was PL/PR. In 22 cases (44%), the preoperative visual acuity was perception of hand movements. In 11 cases (22%), the preoperative visual acuity was 3/60 to counting fingers ½ meters. Visual acuity of the uninjured eye was recorded in all the patients and was found to be within the normal limit.

**Table 4: Distribution of traumatic Cataract patients according to type of surgery**

Type of surgery	N	%
SICS+PCIOL	47	94
SICS+PCIOL with corneal tear repair	1	2
SICS+PCIOL with anterior vitrectomy	1	2
SICS+PCIOL with aphakia	1	2

Out of 50 cases of traumatic cataract, 47 cases underwent ECCE (Small Incision Cataract Surgery) with PCIOL implantation, 1 case underwent small incision cataract surgery with PCIOL implantation with corneal tear repair, 1 case underwent small incision cataract surgery with PCIOL implantation with anterior vitrectomy, 1 case underwent small incision cataract surgery with aphakia.

**Table 5: Distribution of traumatic Cataract according to final visual acuity**

Age groups in years	Final visual acuity		
	6/6-6/18	<6/18-3/60	Less than 3/60
11-20	2	4	0
21-30	4	8	2
31-40	4	7	3
41-50	3	9	0
>50	2	2	0
Total	15	30	5

Out of 50 cases, final visual acuity of 6/6 to 6/18 was seen in 15 (30%) cases. Final visual acuity of less than 6/18 to 6/60 was seen in 30 patients (60%). Five patients (10%) had visual acuity less than 3/60.

### Discussion

Cataract remains the commonest cause of blindness in India contributing about 81%. The incidence of ocular injuries in India is estimated to be 20.5% with 75% cases occurring among those aged less than 40 years. Males are predominantly affected than females with a male to female ratio of 9:1. Further ocular trauma is a major cause of monocular blindness and visual impairment throughout the world, although little is known about its epidemiology or associated visual outcome in developing countries. [16] Cataract is the commonest complication following ocular injury. [5] Traumatic cataract results most commonly from either penetrating injuries from sharp objects like stick or thorn with direct injury to lens or through blunt trauma by objects like stone, cricket ball. Rarely, it can occur from electrical shock, ionizing radiation or infra-red rays (glass blower's cataract). [17]

Accidental ocular trauma can occur at any age but young people are more vulnerable. Cataract is a known complication after penetrating or blunt ocular trauma occurring in around 1-15%. [18] It is estimated that 14% of all cases of cataract in children are due to ocular trauma. The type of trauma, extent of lenticular involvement and associated ocular damage determines the ultimate visual prognosis. [19] The age group ranged from 18-62 years. More number of cases was found in the age group of 21-40 years (56%). A similar age group distribution was also showed by a study by Daljith Singh et al. [20] Out of 50 cases, 35 were males and 15 were females. This study showed a male preponderance. This was because men are more exposed to ocular trauma because of occupation and they are from age earning group.

Out of 50 cases of traumatic cataract, 15 were blunt and 35 were penetrating trauma. Pre-operative visual acuity was recorded in all cases. In 15 cases (30%), the preoperative visual acuity was PL/PR. In 22 cases (44%), the preoperative visual acuity was perception of hand movements. In 11 cases (22%), the preoperative visual acuity was 3/60 to counting

fingers ½ meters. Visual acuity of the uninjured eye was recorded in all the patients and was found to be within the normal limit. Out of 50 cases of traumatic cataract, 47 cases underwent ECCE (Small Incision Cataract Surgery) with PCIOL implantation, 1 case underwent small incision cataract surgery with PCIOL implantation with corneal tear repair, 1 case underwent small incision cataract surgery with PCIOL implantation with anterior vitrectomy, 1 case underwent small incision cataract surgery with aphakia. In 1996, Marcus Blum et al [21] made a study in 148 eyes with traumatic cataract. There PCIOL was implanted in 42(66.6%) of penetrating injury group and in 72(84.7%) patients of blunt trauma. In 1998, Krishnamachary M, Rathi V et al [22] reviewed 237 children who developed traumatic cataract. In the study extra capsular cataract extraction with IOL implantation was performed in 65.67% of patients.

Out of 50 cases, final visual acuity of 6/6 to 6/18 was seen in 15 (30%) cases. Final visual acuity of less than 6/18 to 6/60 was seen in 30 patients (60%). Five patients (10%) had visual acuity less than 3/60. Renuka Srinivasan, Kumudhan et al [23] noted a final visual acuity of 6/12 or better in 88.2% patients. Eckstein M et al [24] noted a visual acuity of 6/12 or better in 67% of patients who underwent cataract extraction with PCIOL implantation. Brar et al [25], have reported a visual acuity of 20/40 in 39% eyes in penetrating trauma compared to 87% in blunt trauma after surgery.

### Conclusion

Traumatic cataract is a serious visually challenging sequel of trauma. In cases of traumatic cataract, to have a better visual outcome after surgery, early diagnosis followed by proper management plays an important role. Therefore stress has to be given on awareness of the public and ocular safety measures should be taken at work places to prevent the ocular hazards associated with the ocular trauma. In addition, early reporting and adequate follow up especially in cases of children needs to be emphasized.

### References

1. Thylefors B. Epidemiological patterns of ocular trauma. Australian and New Zealand journal of ophthalmology. 1992 May;20(2): 9 5-8.

2. Khandekar R, Mohammed AJ, Negrel AD, Al Riyami A. The prevalence and causes of blindness in the Sultanate of Oman: the Oman Eye Study (OES). *British journal of ophthalmology*. 2002 Sep 1;86(9):957-62.
3. Irvine JA, Smith RE. Lens injuries. *eye trauma* .1991:126-5.
4. Jacobs EJ, Tannen BL. Traumatic cataract: a review. *J Ocular Biol*. 2016;4(1):4-8.
5. Bhatia IM, Panda A, Sood NN. Management of traumatic cataract. *Indian Journal Of Ophthalmology*. 1983 May 1;31(3):290-3.
6. Mulrooney BC. Traumatic cataract. *E Med J*, 2002; 3(7): 21-5.
7. Wolter JR. Coup and contre coup mechanism of ocular injuries. *Am J Ophthalmol.*, 1963; 5 6:785-96.
8. Banitt MR, Malta JB, Main SI, Soong HK. Rupture of anterior lens capsule from blunt ocular injury. *J Cataract Refrac Surg.*, 2009; 3 5:943-5.
9. Shingleton BJ, Hersh PS, Kenyon KR. Lens injuries. *Eye Trauma*, 1991; 1: 126-34.
10. Weidenthal DT, Schepens CL. Peripheral fundus change associated with ocular contusion. *Am J Ophthalmol.*, 1966; 62: 465-77 .
11. Greven CM, Collins AS, Slusher MM, Weaver RG. Visual results, prognostic indicators, and posterior segment findings following surgery for cataract/lens subluxation–dislocation secondary to ocular contusion injuries. *Retina*. 2002 Oct 1;22(5):575-80.
12. Ganesh A, Al-Zuhaibi S, Mitra S, Saad Sabt BI, Ganguly SS, Bialasiewicz AA. Visual rehabilitation by scleral fixation of posterior chamber intraocular lenses in Omani children with aphakia. *Ophthalmic Surgery, Lasers and Imaging Retina*. 2009;40(4):354-60.
13. Sarikkola AU, Sen HN, Uusitalo RJ, Laatikainen L. Traumatic cataract and other adverse events with the implantable contact lens. *Journal of Cataract & Refractive Surgery*. 2005 Mar 1;31(3):511-24.
14. Kanski JJ. *Clinical Ophthalmology: A Systematic Approach*. 1989.257.
15. Kuhn F, Mester V, Morris R. Intraocular foreign body in: Kuhn F, Pieramici DJ: *Ocular trauma*.
16. Khattry SK, Lewis AE, Schein OD, et al.. The epidemiology of ocular trauma in rural Nepal. *Br J Ophthalmol*. 2004;88(4):456-60.
17. Mulrooney BC. Traumatic cataract. *EMed J.*, 2002; 3(7): 21-5.
18. Braganza A, Thomas R, George T, Mermoud A. Management of phacolytic glaucoma: experience of 135 cases. *Indian journal of ophthalmology*. 1998 Sep 1;46(3):139.
19. Ekstein M, vijayalakshmi P, Killedar M, Gilbert C, Foster A. Use of intraocular lenses in children with traumatic cataract in south India. *Br J Ophthalmol.*, 1998; 82(8): 911-5.
20. Singh D, Singh K, Singh J, Sood R. The role of intraocular lens in traumatic cataract. *Indian J Ophthalmol*. 1983.
21. Blum M, Tetz MR, Greiner C, Voelcker HE. Treatment of traumatic cataracts. *Journal of Cataract & Refractive Surgery*. 1996 Apr 1;22 (3):342-6.
22. Krishnamachary M, Rathi V, Gupta J. Management of traumatic cataract in children. *Jr. Of cat and ref surgery*, 1997; 23 (1): 681-7.
23. Kumadhan SR. Traumatic cataract-factors affecting visual outcome. *Jr. of TNOA* .;37(1): 45-8.
24. Eckstein M, Vijayalakshmi P, Killedar M, Gilbert C, Foster A. Use of intraocular lenses in children with traumatic cataract in south India. *British journal of ophthalmology*. 1998 Aug 1;82(8):911-5.
25. Brar GS, Ram J, Pandav SS, Reddy GS, Singh U, Gupta A. Postoperative complications and visual results in unocular pediatric traumatic cataract. *Ophthalmic Surgery, Lasers and Imaging Retina*. 2001 May 1;32(3):233-8.