

**A Hospital-Based Study Assessing the Efficacy of Intravenous Iron Sucrose for the Treatment of Iron Deficiency Anemia in Pregnancy**Neha Gangane<sup>1</sup>, Poonam Varma Shivkumar<sup>2</sup>, Surekha Tayade<sup>3</sup>, Shuchi Jain<sup>4</sup><sup>1</sup>Assistant Professor, Department of Obstetrics and Gynaecology, AIIMS, Nagpur, India<sup>2</sup>Professor and Head, Department of Obstetrics and Gynecology, MGIMS, Sewagram, India<sup>3</sup>Professor, Department of Obstetrics and Gynecology, MGIMS, Sewagram, India<sup>4</sup>Professor, Department of Obstetrics and Gynecology, MGIMS, Sewagram, India

Received: 17-09-2023 Revised: 20-10-2023 / Accepted: 29-11-2023

Corresponding Author: Dr. Neha Gangane

Conflict of interest: Nil

**Abstract****Aim:** The aim of the present study was to evaluate the efficacy of intravenous iron sucrose for the treatment of iron deficiency anemia in pregnancy.**Methods:** The present study carried out in Department of Obstetrics and Gynaecology, MGIMS, Sewagram, India. The duration of the study was about 6 months. 50 pregnant women were enrolled for this study.**Results:** The study results showed that the mean age of the pregnant women was 23.57±4.08 years, their mean weight was 59.61±12.28 kg, their mean gestational week was 28.72±4.08, 8 (16%) of them had gestational hypertension, 5 (10%) of them had gestational diabetes, 4 (8%) of them had hypothyroidism, 6% had asthma and 6% genitourinary infection whereas 4 (4%) of them had chronic kidney disease. The study results further showed that both the hemoglobin (p<0.001) and ferritin levels (p<0.001) of females were significantly increased at term after receiving intravenous iron sucrose as compared to the baseline. Furthermore, significant difference was observed in PCV (p<0.001) and MCV as well (p<0.001).**Conclusion:** This study concluded that the administration of iron sucrose intravenously is a secure and effective choice in the management of iron deficiency anemia in pregnant women particularly for those who had inadequate response to oral iron supplementation. Intravenous iron sucrose is well accepted along with a controllable safety profile clinically and enhanced Hemoglobin and ferritin levels thus decreasing complications during pregnancy due to iron deficiency anemia.**Keywords:** Iron deficiency anemia, intravenous iron sucrose, Efficacy, SafetyThis is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.**Introduction**

The World Health Organization [1] refers to this condition as anemia when hemoglobin is less than 11 grams/dl and hematocrit is less than 33. It is also known as the most common medical disorder in pregnancy. Anaemia is indirectly responsible for 40 percent to 50 percent of maternal deaths, especially in developing countries. [2] In well-developed countries, about 18 percent of females are affected by anemia. Moreover, in developing countries, about 35% to 75% of females are affected by anaemia. [3] All around the world, the most common cause of anaemia in pregnancy is the deficiency of iron. [4] It is reported by the World Health Organisation that women of fertile age are mostly affected by iron deficiency anaemia, representing a 50% proportion. [5] Multiple factors are involved in anemia in pregnant women in the developing countries that vary by topographical areas.<sup>6</sup> Globally, deficiency of iron is the major cause of

anemia throughout pregnancy, while the secondary cause is the constant insufficient intake and menstruation, increased requirement of the fetus and increase volume of maternal blood in pregnancy, physiologically. [6,7]

Anemia during pregnancy is associated with complications such as post-partum hemorrhage, low birth weight, premature births, stillbirths, and maternal deaths. [8] The World Health Organization (WHO) estimates that nearly 40% of pregnant women and one-third of all women of reproductive age worldwide are anemic. [9] In India, the National Family Health Survey, 2019-2021 (NFHS-5) reported that 52.2% of pregnant women in India were anemic, the prevalence being higher in rural areas (54.3%) than in urban areas (45.7%). Iron deficiency is the most common cause of anemia and is estimated to contribute to approximately 50% of

all cases of anemia among non-pregnant and pregnant women worldwide. [10] Oral iron supplementation (iron-folic acid tablets) is the therapy of choice for prophylaxis and treatment of mild and moderate iron deficiency anemia in pregnancy. [11] However, oral iron therapy requires a prolonged duration of treatment, which is often beset with poor compliance. [12] Therefore, parenteral iron therapy is an alternative treatment modality for pregnant women with moderate anemia. [13]

The determinants of birth weight and intrauterine growth are maternal weight and parental height. In 80% of the patients, the target is to achieve a haemoglobin of 11 g/dl, which can be achieved by giving intravenous iron sucrose. [14] For iron-defined patients, erythropoiesis is used to incorporate intravenous iron into haemoglobin within three to four weeks. [15] Intravenous iron is effective as it has no serious adverse effects and restores iron more effectively and faster. [16] It is a very safe, effective, and convenient therapy to treat iron deficiency anaemia in pregnancy. [17]

The aim of the present study was to evaluate the efficacy of intravenous iron sucrose for the treatment of iron deficiency anemia in pregnancy.

#### Materials and Methods

The present study was carried out in the Department of Obstetrics and Gynaecology, MGIMS, Sewagram India. The duration of the study was about 6 months. 50 pregnant women were enrolled in this study.

Pregnant women with Hb level equivalent to or <10 g/dl, Serum ferritin level equivalent to or <15 ng/l,

with age ranging from 18-40 years, gestational age of 16 weeks till 36 weeks were included in the study while identified allergic reaction to any active component, anemia not caused by lack of iron (such as hemolytic anemia), chronic or acute bacterial infection, pregnant women with gestational age <16 weeks, excess of iron or contraindication to iron (such as haemosiderosis, hemochromatosis), liver cirrhosis and hepatitis, treated with iron products intravenously or transfusion of blood in 4 weeks were excluded from the study.

Demographic data and co-morbidities were recorded at the time of registration. A two times-weekly dose of 200 mg of iron sucrose intravenously was infused to pregnant women, until the aim of total calculated dose of patient was accomplished. The total collective dose of iron sucrose, equal to the total iron deficit (mg) was calculated by the hemoglobin level (Hb) and body weight (BW). The dose of iron sucrose was individually calculated for each patient according to the total iron deficit with this formula.

#### Data Analysis

SPSS version 22 was applied to analyze the data. Frequency and percentages were calculated for categorical variables such as gender, co-morbidities, adverse effects, etc. Mean±Standard deviation was calculated for numerical variables such as age, Hb, and Ferritin level. Wilcoxon rank-test was used to compare mean Hb and ferritin levels at, baseline and at term. P<0.05 was taken as a statistically significant level.

#### Results

**Table 1: Baseline profile of pregnant females**

Variables	N (%) / Mean±SD
Age (years)	23.57±4.08
Maternal weight (kg)	59.61±12.28
Gestational week	28.72±4.08
Gestational diabetes	8 (16)
Hypertension	5 (10)
Hypothyroidism	4 (8)
Asthma	3 (6)
Chronic kidney disease	2 (4)
Genitourinary infection	3 (6)

The study results showed that the mean age of the pregnant women was 23.57±4.08 years, their mean weight was 59.61±12.28 kg, their mean gestational week was 28.72±4.08, 8 (16%) of them had gestational hypertension, 5 (10%) of them had

gestational diabetes, 4 (8%) of them had hypothyroidism, 6% had asthma and 6% genitourinary infection whereas 2 (4%) of them had chronic kidney disease.

**Table 2: Comparison of baseline and term of hematological values**

Variables	Day-zero	at Term	P value
	Mean±SD	Mean±SD	
Hb (mg/dl)	8.06±0.74	13.87±12.08	<0.001
Ferritin (ng/ml)	9.90±12.58	52.78±56.44	<0.001
Mean corpuscular volume (fl)	76.44±13.27	77.43±13.07	<0.001
Pack cell volume (%)	28.92±5.15	38.22±40.06	<0.001

The study results further showed that both the hemoglobin ( $p<0.001$ ) and ferritin levels ( $p<0.001$ ) of females were significantly increased at term after receiving intravenous iron sucrose as compared to the baseline. Furthermore, significant difference was observed in PCV ( $p<0.001$ ) and MCV as well ( $p<0.001$ ).

### Discussion

Globally, one of the most frequent nutritional deficiencies is iron deficiency anemia. Even though both genders and all ages are affected by nutritional anemia, the dilemma is more widespread in women which lead to maternal morbidity and death, in addition to a low weight of their babies at birth. [18] In developing countries, it has been predicted that about two-thirds of pregnant women are affected by nutritional anemia. However, in developing countries, most women were anemic at the time of conception with a projected occurrence of anemia of approximately 50% amongst non-pregnant women. Multiple factors are involved in anemia in pregnant women in developing countries that vary by topographical areas. [19] Globally, deficiency of iron is the major cause of anemia throughout pregnancy, the cause is the constant insufficient intake, excessive menstrual loss pre-conception, increased requirement of the fetus, and increased volume of maternal blood in pregnancy, physiologically. [19,20]

Generally, this iron is activated from iron stores. Moreover, women with already deprived stores of iron, develop deficiency of iron during pregnancy. One of the studies has revealed that Hb levels less than 8 g% (moderate to severe anemia) in pregnancy are related to higher maternal morbidity whereas Hb <5 g% is linked with cardiac de-compensation and edema of lungs. Loss of even 200 ml of blood in the third phase of labor leads to abrupt shock and fatality in these women. [21]

The study results showed that the mean age of the pregnant women was  $23.57\pm 4.08$  years, their mean weight was  $59.61\pm 12.28$  kg, their mean gestational week was  $28.72\pm 4.08$ , 8 (16%) of them had gestational hypertension, 5 (10%) of them had gestational diabetes, 4 (8%) of them had hypothyroidism, 6% had asthma and 6% genitourinary infection whereas 2 (4%) of them had chronic kidney disease.

Multiple studies have proposed that IV iron sucrose is harmless and effective substitute to oral iron in the management of Iron deficiency anemia. [22-24]

The study results further showed that both the hemoglobin ( $p<0.001$ ) and ferritin levels ( $p<0.001$ ) of females were significantly increased at term after receiving intravenous iron sucrose as compared to the baseline. Furthermore, a significant difference was observed in PCV ( $p<0.001$ ) and MCV as well ( $p<0.001$ ). A randomized control assessment reported by Neeru et al [25], utilized iron sucrose intravenously in contrast with oral iron for the management of iron deficiency anemia and observed that the efficacy of iron sucrose was more in raising hemoglobin levels significantly (23.62% vs 14.1% in oral iron) ( $p<0.05$ ). In another randomized study by Dubey et al [26] after administration of iron sucrose intravenously or oral iron in 200 pregnant women, it was observed that iron sucrose augmented hemoglobin levels and iron stores more rapidly as compared to oral iron significantly ( $p<0.001$ ). The high acceptance of the drug has been partially accredited to the sluggish discharge of iron from the iron sucrose complex and also because of a low tendency to cause allergic reactions of sucrose. [27] The finding of the above study was contradictory to the present study, where no major side effect was reported.

### Conclusion

This study concluded that the administration of iron sucrose intravenously is a secure and effective choice in the management of iron deficiency anemia in pregnant women particularly for those who had inadequate response to oral iron supplementation. Intravenous iron sucrose is well accepted along with a controllable safety profile clinically and enhanced Hemoglobin and ferritin levels and thus decreases complications during pregnancy due to iron deficiency anemia.

### References

1. Naz FA, Iqbal KI, Afreen NO, Javaid A, Khan S, Zareen A. Efficacy of intravenous iron sucrose therapy in iron deficiency anaemia of pregnancy. *Pakistan Journal of Medical and Health Sciences*. 2010;4(1):130-3.
2. Wali A, Mushtaq A. Comparative study--efficacy, safety and compliance of intravenous iron sucrose and intramuscular iron sorbitol in

- iron deficiency anemia of pregnancy. *Journal of Pakistan Medical Association*. 2002;52(9): 392.
3. Sharma J, Tiwari S. Intravenous iron sucrose therapy in iron deficiency anemia in antenatal and postnatal patients. *JNMA J Nepal Med Assoc*. 2015 Apr 1;53:104-7.
  4. Swami MB, Tiwari N, Sharma P. Intravenous iron sucrose therapy in anemia with pregnancy. *Journal of Evolution of Medical and Dental Sciences*. 2015 Aug 24;4(68):11850-7.
  5. Singh A, Singh S, Sharma A. The study of efficacy of injection iron sucrose in treatment of iron deficiency anemia in pregnancy. *International Journal of Reproduction, Contraception, Obstetrics and Gynecology*. 2019 Jun 1;8(6):2444-9.
  6. van den Broek N. Anaemia in pregnancy in developing countries. *Br J ObstetGynaecol*. 1998;105(4):385-90.
  7. Gopalan C. Current food and nutrition situation in south Asian and south-east Asian countries. *Biomed Environ Sci*. 1996;9(2-3): 102-16.
  8. Rahman MM, Abe SK, Rahman MS, Kanda M, Narita S, Bilano V, Ota E, Gilmour S, Shibuya K. Maternal anemia and risk of adverse birth and health outcomes in low-and middle-income countries: systematic review and meta-analysis, 2. *The American journal of clinical nutrition*. 2016 Feb 1;103(2):495-504.
  9. World Health Organization. Prevalence of anaemia in women aged 15–49, by pregnancy status. Geneva: Who. 2021.
  10. Chaparro CM, Suchdev PS. Anemia epidemiology, pathophysiology, and etiology in low-and middle-income countries. *Annals of the new York Academy of Sciences*. 2019 Aug;1450(1):15-31.
  11. Ministry of Health and Family Welfare, Government of India. Guidelines for control of iron deficiency anaemia. (2013).
  12. Cook JD: Diagnosis and management of iron-deficiency anaemia . *Best Pract Res Clin Haematol*. 2005, 18:319-32.
  13. Ministry of Health and Family Welfare, Government of India. Anemia Mukht Bharat operational guidelines. (2019).
  14. Gupta A, Manaktala U, Rathore AM. A randomised controlled trial to compare intravenous iron sucrose and oral iron in treatment of iron deficiency anemia in pregnancy. *Indian Journal of Hematology and Blood Transfusion*. 2014 Jun;30:120-5.
  15. Kriplani A, Mahey R, Dash BB, Kulshreshtha V, Agarwal N, Bhatla N. Intravenous iron sucrose therapy for moderate to severe anaemia in pregnancy. *The Indian journal of medical research*. 2013 Jul;138(1):78.
  16. Garg A, Agarwal M, Shankar U, Shetty S. Comparative study of oral iron and intravenous iron sucrose for the treatment of iron deficiency anemia in pregnancy. *International Journal of Reproduction, Contraception, Obstetrics and Gynecology*. 2017 Jan 1;6(1): 172-6.
  17. Kondrakunta DR. Study of efficacy and compliance of iron sucrose in iron deficiency anaemia in pregnancy. *Int J Clin Obstet Gynaecol*. 2021;5(5):170-4.
  18. WHO. Maternal Health and Safe Motherhood Programme & World Health Organization. Nutrition Programme. The prevalence of anaemia in women : a tabulation of available information, 2021.
  19. van den Broek N. Anaemia in pregnancy in developing countries. *Br J ObstetGynaecol*. 1998;105(4):385-90.
  20. Gopalan C. Current food and nutrition situation in south Asian and south-east Asian countries. *Biomed Environ Sci*. 1996;9(2-3): 102-16.
  21. Toteja GS, Singh P, Dhillon BS, Saxena BN, Ahmed FU, Singh RP et al. Prevalence of anemia among pregnant women and adolescent girls in 16 districts of India. *Food Nutr Bull*. 2006;27(4):311-5.
  22. Al RA, Unlubilgin E, Kandemir O, Yalvac S, Cakir L, Haberal A. Intravenous versus oral iron for treatment of anemia in pregnancy. *Obstet Gynecol*. 2005;106(6):1135-40.
  23. Cançado RD, Novis de Figueiredo PO, AlbeOlivatoMS, Chiattoni CS. Efficacy and safety of intravenous iron sucrose in treating adults with iron deficiency anemia. *Rev Bras HematolHemoter*. 2011;33(6):439-43.
  24. Bhandal N, Russell R. Intravenous versus oral iron therapy for postpartum anaemia. *BJOG*. 2006;113(11):1248-52.
  25. Neeru S, Nair NS, Rai L. Iron sucrose versus oral iron therapy in pregnancy anemia. *Indian J Comm Med*. 2012;37(4):214-8.
  26. Dubey S, Suri V, Aggarwal N, Das R. Is it safe to use intravenous iron sucrose during pregnancy? a randomized controlled trial. *Int J ReprodContraceptObstet Gynecol*. 2013;2(4): 544-9.
  27. Perewusnyk G, Huch R, Huch A, Breyman C. Parenteral iron therapy in obstetrics: 8 years experience with iron-sucrose complex. *Br J Nutr*. 2002;88(1):3-10.