

A Cross-Sectional Assessment of the Relaxation Techniques in Reducing Stress Levels by Measuring Heart Rate Variability

Suman¹, Robina Shamim², Jyoti Priya³, Rita Kumari⁴

¹Tutor, Department of Physiology, Nalanda Medical College, Patna, Bihar, India

²Tutor, Department of Physiology, Nalanda Medical College, Patna, Bihar, India

³Assistant Professor, Department of Physiology, Nalanda Medical College, Patna, Bihar, India

⁴Professor and HOD, Department of Physiology, Nalanda Medical College, Patna, Bihar, India

Received: 14-09-2023 Revised: 21-10-2023 / Accepted: 18-11-2023

Corresponding author: Dr. Robina Shamim

Conflict of interest: Nil

Abstract

Aim: The aim of the present study was to assess the use of Relaxation Techniques in Reducing Stress Levels by Measuring Heart Rate Variability at tertiary health care centre.

Methods: This was a cross-sectional study carried out in the apparently normal individuals to study the effect of various relaxation techniques like Breath focus, Body scan, Guided imagery, Mindfulness meditation, Yoga, taichi, and qigong, Repetitive prayer etc. on heart rate variability during the 12-month period at Nalanda Medical College, Patna, Bihar, India. There were 100 individuals who given written consent to be part of the study.

Results: The average age in both the age group was 36 ± 3.97 and 34 ± 4.26 was not statistically significant ($p > 0.05$). The baseline parameters in both the groups were comparable ($p > 0.05$) but the HRV after 12 weeks therapy High Frequency HRV increased in Group A as compared to Group B was significant ($P < 0.05$); The Low frequency waves increased in Group B as compared to Group A ($p < 0.05$) decreased Low frequency HRV and LF/HF ration after 12 weeks intervention was statistically significant ($P < 0.05$).

Conclusion: It can be concluded from our study that after 12 weeks training with various relaxations techniques the autonomic function measured by HRV i.e. High frequency Increased and low frequency decreased hence the parasympathetic dominance increased with the relaxations techniques.

Keywords: Relaxation Techniques, Heart Rate Variability (HRV), Autonomic function of heart

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

Introduction

A state of physiological and psychological imbalance is known as stress occurs when a person perceives the situation demands to be greater than their ability to satisfy it. It is the natural reaction to an unexpected incident that happens in a person's life. This reaction may be mental, physical, or emotional. [1] According to medical definitions, stress occurs when a person perceives a situation as threatening and their body responds accordingly. The body releases hormones to get ready for action. Increased blood pressure and heart rate enhanced blood flow to the muscle and heart. Reduced blood flow occurs in less critical areas, like the digestive system. Under stress, dizziness, palpitations, and nausea were reported. [2]

Stress can have both positive and negative effects on us. Positive stress can spur us to action, while negative stress can cause feelings of worry, discomfort, rage, and sadness that can worsen our mental and physical health. [3] Various relaxation

techniques that include practices such as progressive muscle relaxation, deep breathing exercises, and mindfulness meditation are used to relax and reduce a person's stress and anxiety. The short-term physiological relaxation response is regulated by two branches of the autonomic nervous system (ANS). The parasympathetic branch of the ANS is more active while resting and therefore, its activation reduces physiological arousal. The sympathetic branch of ANS is more active during emergency reactions and promotes physiological arousal. [4] Relaxation is the ability of the human body to reduce the physiological arousal that manifests itself in the human body through a less activated sympathetic branch, decreased muscle tension, diminished heart rate, lowered respiration rate, increased skin temperature, and decreased sweating. [5]

Various relaxation techniques use different instructions and training to elicit a relaxation

response. A deep breathing exercise procedure is based on breathing at controlled rates, particularly at 6 breaths/minute [6] whereas when applying a cognitive relaxation technique, people are asked to engage in relaxation strategies such as reliving one's pleasant memories or replaying one's favorite song internally for a period of 15–20 min. [7] Mindfulness meditation strategies refer to focusing one's attention in a non-judgmental manner to experiences that exist in the present moment—this may include focusing on breath, body scan, and mindful walking. [8,9]

The aim of the present study was to assess the use of Relaxation Techniques in Reducing Stress Levels by Measuring Heart Rate Variability at tertiary health care centre.

Materials and Methods

This was a cross-sectional study carried out in the apparently normal individuals to study the effect of various relaxation techniques like Breath focus, Body scan, Guided imagery, Mindfulness meditation, Yoga, taichi, and qigong, Repetitive prayer etc. on heart rate variability during the 12 month period at Nalanda Medical College, Patna, Bihar, India. There were 100 individuals who given written consent to be part of the study so these participants randomly divided into two groups Group A (n=50, Receiving relaxation techniques) and Group B (n=50, Control). All necessary details like age, sex. HRV analysis was derived by ECG machine. The parameters were compared in both group at baseline (by paired t-test) and after 12 weeks and in Group E and Group Y at the end of 12 weeks was by unpaired t-test calculated by SPSS 19 version software.

Results

Table 1: Distribution of the patient's as per age and sex

	Group Y (n=50)	Group E (n=50)	p-value
Average age (Mean ± SD)	36± 3.97	34± 4.26	>0.05
Sex			>0.05
Male	36	34	
Female	14	16	

The average age in both the age group was 36± 3.97 and 34± 4.26 was not statistically significant ($p>0.05$).

Table 2: Distribution of the study subjects as per the HRV

	Group A		Group B		p-value
	Basal	After 12 weeks	Basal	After 12 weeks	
HF (nu)	51.19 ± 17.13	57± 4.96	55.59± 8.62	42.38 ± 21.88	P<0.05
LF (nu)	43.97 ± 9.81	36.64± 8.72	38.72 ± 5.40	53.15 ± 14.86	P<0.005
LF/HF	1.22 ± 0.64	0.34 ± 0.17	1.08± 0.42	3.47 ± 1.73	P<0.01
SDNN (ms)	38.72±5.85	48± 12.72	58.22 ± 12.62	37± 3.77	p>0.05

The baseline parameters in both the groups were comparable ($p>0.05$) but the HRV after 12 weeks therapy High Frequency HRV increased in Group A as compared to Group B was significant ($P<0.05$); The Low frequency waves increased in Group B as compared to Group A ($p<0.05$) decreased Low frequency HRV and LF/HF ration after 12 weeks intervention was statistically significant ($P<0.05$).

Discussion

Yoga is an ancient science originated in India. It includes diverse practices, such as physical postures, regulated breathing, instructed relaxation and meditation. [10] Among the yoga based relaxation techniques, Deep relaxation technique (DRT), Cyclic meditation (CM), Shavasan are most popular. Practicing relaxation has earned popularity in Bangladesh. Recently, relaxation has come in limelight because of its cardiac and multisystem benefits among the practitioners. [11-14]

The average age in both the age group was 36± 3.97 and 34± 4.26 was not statistically significant ($p>0.05$). The baseline parameters in both the groups were comparable ($p>0.05$) but the HRV after 12 weeks therapy High Frequency HRV increased in Group A as compared to Group B was significant ($P<0.05$); The Low frequency waves increased in Group B as compared to Group A ($p<0.05$) decreased Low frequency HRV and LF/HF ration after 12 weeks intervention was statistically significant ($P<0.05$). A number of studies using existing stress measurement methods (e.g., psychological measures of stress) and examining biological markers (e.g., cortisol, amylase) have been performed. Moreover, studies on heart rate variability (HRV) and stress are increasing in frequency. HRV is the fluctuation of the length of heart beat intervals. [15] HRV represents the ability of the heart to respond to a variety of physiological and environmental stimuli. [16] Low HRV conveys a monotonously regular heart rate. Moreover, low HRV is associated with impaired regulatory and

homeostatic autonomic nervous system (ANS) functions, which reduce the body's ability to cope with internal and external stressors.

In view of observations of stress-associated variation in HRV and existing neurobiological evidence, HRV may be used as an objective assessment of stress and mental health. However, since psychiatric illnesses have numerous causes and symptoms, consistent biological measurements are difficult to acquire in individuals with mental illness. Thus, a patient's psychological and medical history should be equally considered when interpreting HRV results. Therefore, HRV can be considered a tool that reflects heart activity and overall autonomic health, rather than specific mental illnesses or disease states. Since the concept of stress includes biological and psychological factors, objective and physiological evaluations as well as self-reporting should be integrated when evaluating stress, using HRV in clinical practice. Many physical conditions and lifestyle habits can affect HRV results, including physiological factors (e.g., breathing, circadian rhythms, and posture), non-modifiable factors (e.g., age, sex, and genetic factors), modifiable lifestyle factors (e.g., obesity, metabolic syndrome, physical activity, smoking, and drinking), and other factors [e.g., medication (e.g., anticholinergics, stimulants, and beta-blockers)]. [17,18] Hans Seyle [19] proposed a three-stage stress response model. The first stage is the "alarm reaction stage," in which the body reacts to a stressor with the fight-or-flight response and activates the SNS. The second stage is the "resistance stage," in which the body adapts to the stressor. During this stage, the PNS restores many physiological functions to normal, while the body focuses its resources against the stressor.

Conclusion

It can be concluded from our study that after 12 weeks training with various relaxations techniques the autonomic function measured by HRV i.e. High frequency Increased and low frequency decreased hence the parasympathetic dominance increased with the relaxations techniques.

References

1. Al Rasheed F, Naqvi AA, Ahmad R, Ahmad N. Academic stress and prevalence of stress-related self-medication among undergraduate female students of health and non-health cluster colleges of a public sector university in Dammam, Saudi Arabia. *Journal of pharmacy & bioallied sciences*. 2017 Oct;9(4):251.
2. Behere SP, Yadav R, Behere PB. A comparative study of stress among students of medicine, engineering, and nursing. *Indian journal of psychological medicine*. 2011 Jul;33(2):145-8.
3. Behere SP, Yadav R, Behere PB. A comparative study of stress among students of medicine, engineering, and nursing. *Indian journal of psychological medicine*. 2011 Jul;33(2):145-8.
4. Behman PJ, Rash JA, Bagshawe M, Giesbrecht GF. Short-term autonomic nervous system and experiential responses during a labyrinth walk. *Cogent Psychology*. 2018 Dec 31;5(1):1495036.
5. Noppe G, de Rijke YB, Dorst K, van den Akker EL, van Rossum EF. LC-MS/MS-based method for long-term steroid profiling in human scalp hair. *Clin Endocrinol (Oxf)*. 2015 Aug;83(2):162-6.
6. Nuckowska MK, Gruszecki M, Kot J, Wolf J, Guminski W, Frydrychowski AF, Wtorek J, Narkiewicz K, Winklewski PJ. Impact of slow breathing on the blood pressure and subarachnoid space width oscillations in humans. *Sci Rep*. 2019 Apr 17;9(1):6232.
7. Khalsa SS, Rudrauf D, Davidson RJ, Tranel D. The effect of meditation on regulation of internal body states. *Front Psychol*. 2015 Jul 7;6:924.
8. Edenfield TM, Saeed SA. An update on mindfulness meditation as a self-help treatment for anxiety and depression. *Psychol Res Behav Manag*. 2012;5:131-41.
9. van der Zwan JE, de Vente W, Huizink AC, Bögels SM, de Bruin EI. Physical activity, mindfulness meditation, or heart rate variability biofeedback for stress reduction: a randomized controlled trial. *Appl Psychop-hysiol Biofeedback*. 2015 Dec;40(4): 257-68.
10. Subramanya T, Telles S. A review of scientific studies on cyclic meditation. *Int J Yoga*. 2009; 2(2): 46-48.
11. Benson H, Klipper MZ. *The Relaxation Response*. USA: Harper Collins; 1975.
12. Ornish D. *Reversing Heart Disease*. USA: Ballantine Books; 1990.
13. Wenneberg SR, Schneider RH, Walton KG, Maclean CR, Levitsky DK, Salerno JW, Wallace RK, Mandarino JV, Rainforth MV, Waziri R. A controlled study of the effects of the Transcendental Meditation program on cardiovascular reactivity and ambulatory blood pressure. *Int J Neurosci*. 1997; 89(1-2):15-28.
14. Paul-Labrador M, Polk D, Dwyer JH, Velasquez I, Nidich S, Rainforth M, Schneider R, Merz CN. Effects of a randomized controlled trial of transcendental meditation on components of the metabolic syndrome in subjects with coronary heart disease. *Archives of internal Medicine*. 2006 Jun 12;166(11):1218-24.
15. Malik M, Camm AJ. *Heart rate variability* (Armonk, NY, Futura Pub. Co.). Inc [Google Scholar]. 1995.

16. Rajendra Acharya U, Paul Joseph K, Kannathal N, Lim CM, Suri JS. Heart rate variability: a review. *Medical and biological engineering and computing*. 2006 Dec;44: 103 1-51.
17. Romanowicz M, Schmidt JE, Bostwick JM, Mrazek DA, Karpyak VM. Changes in heart rate variability associated with acute alcohol consumption: current knowledge and implications for practice and research. *Alcoholism: Clinical and Experimental Research*. 2011 Jun;35(6):1092-105.
18. Altuncu ME, Baspinar O, Keskin M. The use of short-term analysis of heart rate variability to assess autonomic function in obese children and its relationship with metabolic syndrome. *Cardiology journal*. 2012;19(5):501-6.
19. Selye H. *Stress in health and disease*. Butterworth-Heinemann; 2013 Oct 22.