

## A Randomized Clinical Study to Assess the Effect of Spinal Anaesthesia on Perioperative Hyperglycaemia in Diabetic Patients Undergoing Lower Limb Orthopaedic Surgeries

Deepak Kumar Maurya<sup>1</sup>, Uma Shankar Kumar<sup>2</sup>, Pramod Kumar Sinha<sup>3</sup>

<sup>1</sup>Senior Resident, Department of Anaesthesia and Critical Care, A.N. Magadh Medical College and Hospital, Gaya, Bihar, India

<sup>2</sup>Senior Resident, Department of Anaesthesia and Critical Care, A.N. Magadh Medical College and Hospital, Gaya, Bihar, India

<sup>3</sup>Associate Professor & HOD, Department of Anaesthesia and Critical Care, A.N. Magadh Medical College and Hospital, Gaya, Bihar, India

Received: 10-10-2023 / Revised: 14-11-2023 / Accepted: 20-12-2023

Corresponding Author: Dr. Uma Shankar Kumar

Conflict of interest: Nil

### Abstract

**Aim:** The aim of the present study was to assess the effect of spinal anaesthesia on perioperative hyperglycaemia in diabetic patients undergoing lower limb orthopaedic surgeries and also to state the trend of perioperative hyperglycaemia.

**Material & Methods:** A retrospective study conducted in the Department of Anaesthesia and Critical Care, A. N. Magadh Medical College and Hospital, Gaya, Bihar, India from October 2018 to August 2021, including 200 patients having either Type I or Type II Diabetes Mellitus controlled on either oral hypoglycaemic drugs or injectable insulin aged 30 to 65 years, belonging to either sex and American Society of Anaesthesiologists (ASA) physical status II and III undergoing elective lower limb orthopaedic surgeries under spinal anaesthesia.

**Results:** The mean age, weight, height and duration of anaesthesia was 52.54±14.66 years, 64.6±6.4 kg, 161.9±4.7 cm and 105.5±11.6 respectively. Mean BG value preoperatively or 10 min before induction was 112.56±12.088. Then at SI, there was statistically significant decrease in BG value to mean value 107.83±12.58. 30 min after SI, mean BG value was 109.71±16.94. This value was lower as compared to the pre-operative BG value, but not statistically significant. 1hr after SI, BG value was 111.59±13.237. This value was also lower as compared to the pre-operative BG value, but not statistically significant. 2hrs after SI, BG value increased to mean value 122.88±16.444. Even, 3 hrs after SI, BG value continued increasing and the mean value became 125.15±15.005. 4 hrs after SI, BG value was maximum with the value being 127.23±16.384. There was statistically significant difference (p=.000). Blood glucose (BG) value decreases till 1hr after surgical incision (SI), and then increases till 4th hr after SI. This change in blood glucose values is statistically significant at SI, 2nd hr after SI, 3rd hr after SI and 4th hr after SI.

**Conclusion:** Spinal anaesthesia blunts surgical stress response and hence, at SI, BG values decrease. But, BG values increase at other times in perioperative period owing to the regression of sensory analgesia.

**Keywords:** Spinal Anaesthesia, Perioperative, Hyperglycaemia, Diabetic

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

### Introduction

Spinal anaesthesia has become the preferred anaesthetic technique when providing anaesthesia for patients undergoing lower abdominal surgeries, lower limb surgeries, elective caesarean section as the risk of maternal and foetal complications associated with spinal anaesthesia is less than with general anaesthesia. [1-5] Every surgical procedure is associated with a stress response which comprises a number of endocrine, metabolic, and immunological changes triggered by neuronal activation of the hypothalamic-pituitary-adrenal axis. [6,7]

The overall metabolic effect of the stress response to surgery includes an increase in secretion of catabolic hormones, such as cortisol and catecholamine, and a decrease in secretion of anabolic hormones, such as insulin and testosterone. The increase in levels of catabolic hormones in plasma stimulates glucose production, and there is a relative lack of insulin together with impaired tissue insulin sensitivity and glucose utilization, which is called insulin resistance. Consequently, blood glucose

concentrations will increase, even in the absence of preexisting diabetes. [6-9]

Observational studies show that in the surgical patient, diabetes is associated with a higher rate of peri-operative complications such as need for transfusion, pneumonia, delayed discharge, surgical site infections, and in-hospital mortality. [10] These poorer outcomes are in part due to higher rates of comorbid conditions such as ischemic heart disease, renal impairment, and hypertension in patients with diabetes. [11] Dysglycemia, which encompasses hyperglycaemia, hypoglycaemia, stress-induced hyperglycaemia, and excessive glucose variability, is increasingly observed and associated with poorer post-operative outcomes even in those without a prior diagnosis of diabetes. In fact, several studies have shown postoperative complications occur more frequently in people with stress-induced hyperglycaemia with no prior diagnosis of diabetes than in those with diabetes. [12-14]

Stress hyperglycaemia is defined as any blood glucose concentration  $>7.8$  mmol/l (140 mg/dl) without evidence of previous diabetes by the American Diabetes Association and American Association of Clinical Endocrinologists consensus. [15] Stress-induced hyperglycaemia is common and more than 50% occurs in previously non-diabetic patients. [16,17] Perioperative stress-induced hyperglycaemia is reported in 20–40% of patients undergoing general surgical procedures. [18-20]

Hence the aim of study was to assess the effect of spinal anaesthesia on perioperative hyperglycaemia in diabetic patients undergoing lower limb orthopaedic surgeries and also to state the trend of perioperative hyperglycaemia

### Material & Methods

A retrospective study conducted in the Department of Anaesthesia and Critical Care, A. N. Magadh Medical College and Hospital, Gaya, Bihar, India from October 2018 to August 2021, including 200

patients having either Type I or Type II Diabetes Mellitus controlled on either oral hypoglycaemic drugs or injectable insulin aged 30 to 65 years, belonging to either sex and American Society of Anaesthesiologists (ASA) physical status II and III undergoing elective lower limb orthopaedic surgeries under spinal anaesthesia.

### Inclusion Criteria

Only patients having preoperative blood glucose level between 80-120mg/dl were included in the study.

### Methodology

Patients on recent intravenous or oral steroid therapy within 30 days, although inhaled steroids were permitted, known case of chronic obstructive respiratory disease and asthma on intravenous steroid therapy; having coagulation abnormalities, hypovolemia or hypotension, pre-existing severe bradycardia, or ejection fraction 20% from basal HR) and hypotension (mean atrial pressure 20% from basal BP) were recorded and managed as per the standard protocols. When blood glucose concentrations exceeded 180mg/dL, it was treated as hyperglycaemia as per continuous insulin infusion (CII) protocol. When blood glucose concentrations lowered below 60mg/dl, it was treated as hypoglycaemia as per the standard protocol.

### Statistical Analysis

Statistical testing was conducted with the statistical package for the social science system version (SPSS Statistics for Windows, Version 17.0. Chicago: SPSS Inc.). Ages, weight, height, duration of anaesthesia and blood glucose (BG) values were reported as mean  $\pm$  standard deviation. Comparison of BG before, during and after surgery was done using Student's t-test. For statistical test,  $P < 0.05$  was taken to indicate a significant difference.

### Results

**Table 1: Demographic characteristics (age, weight, and height) and duration of anaesthesia**

| Variables                     | Mean $\pm$ SD     |
|-------------------------------|-------------------|
| Age in years                  | 52.54 $\pm$ 14.66 |
| Weight (kg)                   | 64.6 $\pm$ 6.4    |
| Height (cm)                   | 161.9 $\pm$ 4.7   |
| Duration of anaesthesia (min) | 105.5 $\pm$ 11.6  |

The mean age, weight, height and duration of anaesthesia was 52.54 $\pm$ 14.66 years, 64.6 $\pm$ 6.4 kg, 161.9 $\pm$ 4.7 cm and 105.5 $\pm$ 11.6 respectively.

**Table 2: Blood glucose (BG) values (mg/dl)**

| Time                    | Mean $\pm$ SD       |
|-------------------------|---------------------|
| 10 min before induction | 112.56 $\pm$ 12.088 |
| SI                      | 107.83 $\pm$ 12.58  |
| 30 min after SI         | 109.71 $\pm$ 16.94  |

|              |               |
|--------------|---------------|
| 1hr after SI | 111.59±13.237 |
| 2hr after SI | 122.88±16.444 |
| 3hr after SI | 125.15±15.005 |
| 4hr after SI | 127.23±16.384 |

Mean BG value preoperatively or 10 min before induction was 112.56±12.088. Then at SI, there was statistically significant decrease in BG value to mean value 107.83±12.58. 30 min after SI, mean BG value was 109.71±16.94. This value was lower as compared to the pre-operative BG value, but not statistically significant. 1hr after SI, BG value was 111.59±13.237. This value was also lower as

compared to the pre-operative BG value, but not statistically significant. 2hrs after SI, BG value increased to mean value 122.88±16.444. Even, 3 hrs after SI, BG value continued increasing and the mean value became 125.15±15.005. 4 hrs after SI, BG value was maximum with the value being 127.23±16.384. There was statistically significant difference (p=.000).

**Table 3: Trend of blood glucose (BG) values taking BG value at 10min before induction as reference value (mg/dl)**

| Time            | Mean±SD        | P Value |
|-----------------|----------------|---------|
| SI              | 4.746±7.843    | .007    |
| 30 min after SI | .903±12.488    | .743    |
| 1hr after SI    | .929±15.275    | .740    |
| 2hr after SI    | -10.234±23.097 | .014    |
| 3hr after SI    | -12.512±14.076 | .000    |
| 4hr after SI    | -17.633±14.636 | .000    |

Blood glucose (BG) value decreases till 1hr after surgical incision (SI), and then increases till 4th hr after SI. This change in blood glucose values is statistically significant at SI, 2nd hr after SI, 3rd hr after SI and 4th hr after SI.

### Discussion

Spinal anaesthesia (SA) is a commonly performed regional anaesthesia technique in current practice. Drops of cerebrospinal fluid during SA provide an objective criterion for application field compared to other neuraxial anaesthesia procedures. Even so, failure of SA is observed depending different causes (failed lumbar puncture, dose selection, drug solution error, anatomical abnormalities, solution=density, inactive local anaesthetic solution and local anaesthetic resistance). [21]

The stress of surgery results in increased levels of glucose-regulatory hormones (catecholamines, cortisol, glucagon, and growth hormone) and excessive release of interleukin-6 and interleukin-1. The counter-regulatory response produces alterations in carbohydrate metabolism, including insulin resistance, increased hepatic glucose production, impaired peripheral glucose utilization, and relative insulin deficiency. [22,23] The mean age, weight, height and duration of anaesthesia was 52.54±14.6 6 years, 64.6±6.4 kg, 161.9±4.7 cm and 105.5±11.6 respectively. Mean BG value preoperatively or 10 min before induction was 112.56±12.088. Then at SI, there was statistically significant decrease in BG value to mean value 107.83±12.58. 30 min after SI, mean BG value was 109.71±16.94. This value was lower as compared to

the pre-operative BG value, but not statistically significant. 1hr after SI, BG value was 111.59±13.237. This value was also lower as compared to the pre-operative BG value, but not statistically significant. 2hrs after SI, BG value increased to mean value 122.88±16.444. Even, 3 hrs after SI, BG value continued increasing and the mean value became 125.15±15.005. 4 hrs after SI, BG value was maximum with the value being 127.23±16.384. There was statistically significant difference (p=.000). During anaesthesia induction and surgery, insulin concentration may decrease due to  $\alpha$  adrenergic inhibition of  $\beta$ -cell secretion. Plasma glucose concentrations increase in perioperative period. In fact, anaesthesia itself results in hyperglycaemia, which is then further aggravated by the surgical procedure. The initial increase in plasma glucose after injury is due to activation of glycogenolysis. But later hepatic gluconeogenesis becomes the major factor in liver glucose release because liver glycogen stores are limited. The usual mechanisms that maintain glucose homeostasis are ineffective in the perioperative period and catabolic hormones promote the production of glucose, thereby resulting in hyperglycemia. [24-26]

Blood glucose (BG) value decreases till 1hr after surgical incision (SI), and then increases till 4th hr after SI. This change in blood glucose values is statistically significant at SI, 2nd hr after SI, 3rd hr after SI and 4th hr after SI. Poon et al [27] achieved better stress response control by combining epidural anaesthesia with general anaesthesia. Opioids also suppress the stress response by inhibiting hypothalamic pituitary gland function. In a study of

lower abdominal surgery, 50 µg/kg fentanyl suppressed the stress response by reducing growth hormone, cortisol, and glucose concentrations. But, systemic opioids may be insufficient to suppress this response in upper abdominal surgeries. In other study of cholecystectomies using 100 µg/kg fentanyl, the stress response was suppressed; however, patients also required postoperative ventilator support. Most studies of neural blocks have assessed the effect of epidural anaesthesia, but, few have addressed spinal anaesthesia and stress. Moller et al [28] (1984) compared stress responses following spinal and general anaesthesia in abdominal hysterectomies, and reported that spinal anaesthesia had a temporary inhibitory effect, which was correlated with the sensorial block level. According to Basem et al [29] (2013) for all patients combined, mean glucose increased slightly from preoperative to incision, substantially from incision to surgery midpoint, and then remained high and fairly stable through emergence, with nondiabetic patients showing a greater increase. For nondiabetics, the mean increase in glucose concentration was more in patients given dexamethasone than placebo. However, there was no dexamethasone effect in diabetics. They assessed this response in patients undergoing non-cardiac surgery under general anaesthesia.

### Conclusion

Spinal anaesthesia blunts surgical stress response and hence, at SI, BG values decrease. But, BG values increase at other times in perioperative period owing to the regression of sensory analgesia.

### References

1. F. Reynolds, "General anesthesia is unacceptable for elective cesarean section," *International Journal of Obstetric Anesthesia*, vol. 19, no. 2, pp. 212–217, 2010.
2. Wong CA. General anesthesia is unacceptable for elective cesarean section. *International journal of obstetric anesthesia*. 2010 Apr 1;19(2):209-12.
3. Bucklin BA, Hawkins JL, Anderson JR, Ullrich FA. Obstetric anesthesia workforce survey: twenty-year update. *The Journal of the American Society of Anesthesiologists*. 2005 Sep 1;103(3):645-53.
4. Heesen M, Stewart A, Fernando R. Vasopressors for the treatment of maternal hypotension following spinal anaesthesia for elective caesarean section: past, present and future. *Anaesthesia*. 2015 Mar;70(3):252-7.
5. Sia AT, Tan KH, Sng BL, Lim Y, Chan ES, Siddiqui FJ. Hyperbaric versus plain bupivacaine for spinal anesthesia for cesarean delivery. *Anesthesia & Analgesia*. 2015 Jan 1; 120(1):132-40.
6. Hadimioglu N, Ulugol H, Akbas H, Coskunfirat N, Ertug Z, Dinckan A. Combination of epidural anesthesia and general anesthesia attenuates stress response to renal transplantation surgery. In *Transplantation proceedings 2012 Dec 1* (Vol. 44, No. 10, pp. 2949-2954). Elsevier.
7. Desborough JP. The stress response to trauma and surgery. *British journal of anaesthesia*. 2000 Jul 1;85(1):109-17.
8. Demirbilek S, Ganidağlı S, Aksoy N, Becerik C, Baysal Z. The effects of remifentanyl and alfentanil-based total intravenous anesthesia (TIVA) on the endocrine response to abdominal hysterectomy. *Journal of clinical anesthesia*. 2004 Aug 1;16(5):358-63.
9. Geisser W, Schreiber M, Hofbauer H, Lattermann R, Füssel S, Wachter U, Georgieff M, Schrickler T. Sevoflurane versus isoflurane—anaesthesia for lower abdominal surgery. Effects on perioperative glucose metabolism. *Acta anaesthesiologica scandinavica*. 2003 Feb;47(2):174-80.
10. Browne JA, Cook C, Pietrobon R, et al. Diabetes and early postoperative outcomes following lumbar fusion. *Spine*. 2007;32(20): 2214–9.
11. Wukich DK, Joseph A, Ryan M, Ramirez C, Irrgang JJ. Outcomes of ankle fractures in patients with uncomplicated versus complicated diabetes. *Foot Ankle Int*. 2011;32(2):120–30.
12. Kerby JD, Griffin RL, MacLennan P, Rue III LW. Stress-induced hyperglycemia, not diabetic hyperglycemia, is associated with higher mortality in trauma. *Ann Surg*. 2012;256(3):446–52.
13. Karunakar, Madhav A, Staples, Kurtis S. Does stress-induced hyperglycemia increase the risk of peri-operative infectious complications in orthopedic trauma patients? *J Orthop Trauma*. 2010;24(12):752–6.
14. Lauruschkat AH, Arnrich B, Albert AA, Walter JA, Amann B, Rosendahl UP, Alexander T, Ennker J. Prevalence and risks of undiagnosed diabetes mellitus in patients undergoing coronary artery bypass grafting. *Circulation*. 2005 Oct 18;112(16):2397–402.
15. Moghissi ES, Korytkowski MT, DiNardo M, Einhorn D, Hellman R, Hirsch IB, Inzucchi SE, Ismail-Beigi F, Kirkman MS, Umpierrez GE. American Association of Clinical Endocrinologists and American Diabetes Association consensus statement on inpatient glycemic control. *Diabetes care*. 2009 Jun;32(6):1119.
16. Knaak C, Wollersheim T, Mörgeli R, Spies C, Vorderwülbecke G, Windmann V, Kuenz S, Kurpanik M, Lachmann G. Risk factors of intraoperative dysglycemia in elderly surgical

- patients. *International Journal of Medical Sciences*. 2019;16(5):665.
17. Dave JA, Engel ME, Freercks R, Peter J, May W, Badri M, Van Niekerk L, Levitt NS. Abnormal glucose metabolism in non-diabetic patients presenting with an acute stroke: prospective study and systematic review. *QJM: An International Journal of Medicine*. 2010 Jul 1;103(7):495-503.
  18. Kwon S, Thompson R, Dellinger P, Yanez D, Farrohki E, Flum D. Importance of perioperative glycemic control in general surgery: a report from the Surgical Care and Outcomes Assessment Program. *Annals of surgery*. 2013 Jan;257(1):8.
  19. Maitra S, Kirtania J, Pal S, Bhattacharjee S, Layek A, Ray S. Intraoperative blood glucose levels in nondiabetic patients undergoing elective major surgery under general anaesthesia receiving different crystalloid solutions for maintenance fluid. *Anesthesia, essays and researches*. 2013 May;7(2):183.
  20. Shi Z, Tang S, Chen Y, Yang J, Jiang B, Liu X, Zhou X, Pan X, Yang J, Wu J, Hu H. Prevalence of stress hyperglycemia among hepatopancreatobiliary postoperative patients. *International Journal of Clinical and Experimental Medicine*. 2013;6(9):799.
  21. Fettes PD, Jansson JR, Wildsmith JA. Failed spinal anaesthesia: mechanisms, management, and prevention. *British journal of anaesthesia*. 2009 Jun 1;102(6):739-48.
  22. Alberti KG. Role of counterregulatory hormones in the catabolic response to stress. *J Clin Invest* 1984; 74(6):2238-48.
  23. Weismann C. The metabolic response to stress: an overview and update. *Anesthesiology* 1990; 73:308–27.
  24. Burton D, Nicholson G, Hall G. Endocrine and metabolic response to surgery. *Continu Educ Anaesthesia Crit Care Pain* 2004;4:144–7.
  25. Singh M. Stress response and anaesthesia. *Indian Journal of Anaesthesia* 2003; 47:427–434.
  26. Fereshteh Amiri, Ali Ghomeishi, Seyed Mohammad, Mehdi Aslani, Sholeh Nesioonpour, Sara Adarvishi. Comparison of Surgical Stress Responses During Spinal and General Anesthesia in Curettage Surgery. *Anesth Pain Med* 2014 August; 4(3): e20554.
  27. Poon KS, Chang WK, Chen YC, Chan KH, Lee TY. Evaluation of stress response to surgery under general anesthesia combined with spinal analgesia. *Acta Anaesthesiologica Sinica*. 1995 Jun 1;33(2):85-90.
  28. Møller IW, Hjørtso E, Krantz T, Wandall E, Kehlet H. The modifying effect of spinal anaesthesia on intra-and postoperative adrenocortical and hyperglycaemic response to surgery. *Acta anaesthesiologica scandinavica*. 1984 Jun;28(3):266-9.
  29. Abdelmalak BB, Bonilla AM, Yang D, Chowdary HT, Gottlieb A, Lyden SP, Sessler DI. The hyperglycemic response to major noncardiac surgery and the added effect of steroid administration in patients with and without diabetes. *Anesthesia & Analgesia*. 2013 May 1;116(5):1116-22.