

A Hospital Based Observational Assessment of the Post-Operative Morbidity with Relation to Pre-Operative Albumin Status, in Patients Undergoing Emergency Abdominal Surgery

Manoj Kumar Shaw¹, Shanker Bharti², Anjani kumar Anjan³, Rajesh Narayan⁴

¹Assistant Professor, Department of General Surgery, Bhagwan Mahavir Institute of Medical Science, Pawapuri, Nalanda, Bihar

²Senior Resident, Department of General Surgery, Bhagwan Mahavir Institute of Medical Science, Pawapuri, Nalanda, Bihar

³Assistant professor, Department of General Surgery, Bhagwan Mahavir Institute of Medical Science, Pawapuri, Nalanda, Bihar

⁴Associate professor and HOD, Department of General Surgery, Bhagwan Mahavir Institute of Medical Science, Pawapuri, Nalanda, Bihar

Received: 11-02-2023/ Revised: 21-03-2023 / Accepted: 15-04-2023

Corresponding author: Dr. Shanker Bharti

Conflict of interest: Nil

Abstract

Aim: The aim of study was to assess the post-operative morbidity with relation to pre-operative albumin status, in patients undergoing emergency abdominal surgery.

Material & Methods: In this hospital based observational study, 100 patients undergoing emergency abdominal surgery at Department of General Surgery, Bhagwan Mahavir Institute of Medical Science, Pawapuri, Nalanda, Bihar, India were included. Hypoalbuminemia was diagnosed if serum albumin level was less than 3.5 gm/dl. Types of surgery and postoperative complications were included in the study. Any association between preoperative serum albumin level and postoperative morbidity was assessed. Study period was of 2 years. Pre-operatively, 50 patients had hypoalbuminemia (serum albumin <3.5 g/dl) and 50 patients had normal albumin levels (≥ 3.5 g/dl).

Results: Of the 100 patients studied, 65% were male and 35% were female. Pre-operatively, 50 patients had hypoalbuminemia (serum albumin <3.5 g/dl) and 50 patients had normal albumin levels (≥ 3.5 g/dl). Most of the patients belonged to 38-57 years in the present study. The most common indication for emergency abdominal surgery was peptic ulcer perforation 35 (35%), followed by acute intestinal obstruction 30 (30%). Surgical site infection was commonest complication found in 29 (29%) cases. Wound dehiscence was found in 11 (11%) cases. Mean length of hospital stay was found to be higher i.e. 8.72 ± 5.60 days in patients with hypoalbuminemia, compared to 7.35 ± 4.16 days with those having albumin level ≥ 3.5 g/dl.

Conclusion: Preoperative albumin is a useful low-cost prognostic predictor for predicting surgery outcome.

Keywords: Serum albumin, Emergency surgery, Morbidity

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

Introduction

Abdominal surgery is a common surgical procedure. It is performed as either elective surgery or on an emergency basis. Underlying diseases may cause stress to body tissues and surgical trauma adds additional stress that triggers a corrective response by the body.[1] The magnitude of the metabolic stress response with the long duration of surgery may contribute to the development of postoperative complications. Early identification of patients with nutritional deficiencies and comorbidities and adequate initial resuscitation may help in reducing the postoperative mortality and morbidity of patients. [2,3,4] Albumin is the major protein of human plasma. It constitutes approximately 60% of the total plasma protein and its normal serum concentration is 3.5–5.0 g/dl. A serum level of less than 3.4 g/dl is considered hypoalbuminemia (citation). Plasma albumin has three primary functions: osmotic, transportation and nutritional and it accounts for more than 75–80% of total plasma osmotic pressure (25 mmHg). During physiological stress, decreases in serum albumin levels to hypoalbuminemia levels causes a fall in oncotic pressure, which in turn leads to interstitial oedema. [5] Albumin, a crucial protein, transports hormones, fatty acids, and exogenous medications while also regulating plasma oncotic pressure. Because albumin levels drop during injuries and infection, albumin is referred to as a negative acute-phase protein.[6] A maintenance protein called serum albumin is quickly downregulated by inflammatory signals. Low levels of serum albumin are mostly brought on by inflammatory conditions, specifically by high levels of cytokines interleukin-6 (IL-6) and tumor necrosis factor-alpha (TNF-alpha). A common finding in both acute and chronic diseases is hypoalbuminemia. The mechanisms causing hypoalbuminemia in acute conditions are different from those in chronic diseases because capillary leakage into the interstitial space as a result of

inflammatory processes is the main source of hypoalbuminemia in acute conditions. Additionally, reduced synthesis, dilution of blood due to fluid administration, renal and intestinal losses due to congestion, and increased catabolism also play a role. [7,8,9]. A preoperative serum albumin level in patients without certain comorbid diseases is associated with postoperative infectious and surgical complications. Low serum albumin levels can be used as a marker for nutritional deficiency. [10] A serum albumin level >3.5 g/dL suggests adequate protein stores and confers a protective effect through several biological mechanisms. It predicts perioperative morbidity and mortality. Serum albumin level <3.5 g/dL in a stable and well-hydrated patient suggests malnutrition. [11] The preoperative serum albumin is a good predictor of surgical outcomes in elective gastrointestinal surgeries as well as many other surgeries except cardiac surgeries.[12,13] This study is an attempt to analyze the correlation between preoperative hypoalbuminemia and surgical complications encountered in patients undergoing emergency abdominal surgery.

Material & Methods

In this hospital based observational study, 100 patients undergoing emergency abdominal surgery at Department of General Surgery, Bhagwan Mahavir Institute of Medical Science, Pawapuri, Nalanda, Bihar, India were included. Hypoalbuminemia was diagnosed if serum albumin level was less than 3.5 gm/dl. Types of surgery and postoperative complications were included in the study. Any association between preoperative serum albumin level and postoperative morbidity was assessed. Study period was of 2 years. Pre-operatively, 50 patients had hypoalbuminemia (serum albumin <3.5 g/dl) and 50 patients had normal albumin levels (≥ 3.5 g/dl).

A detailed history was taken, and a thorough clinical examination was performed. Relevant biochemical and radiological investigations were done. Data on type of surgery, post-operative complications were collected. Patients with no significant comorbidity admitted and operated within 24 hours were included in the study. Serum albumin level of 3.5 g/dl was considered the standard baseline in the study. Serum albumin level was less than 3.5 gm/dl was considered as hypoalbuminemia. The association between preoperative serum albumin level and postoperative morbidity was assessed. Ethical clearance was obtained from the institutional ethics committee (human) prior to the conduction of the study.

Exclusion criteria

Patients with chronic debilitating disease like patients having anemia with hemoglobin <7 g/dl, liver failure, nephrotic syndrome, diabetic patients; immunocompromised patients; femoral

hernias; perforation due to medical causes like typhoid ulcer perforation; patients with associated head injury, orthopedic injury, renal injury, chest trauma

Statistical Analysis

Statistical testing was conducted using SPSS Statistics version 21.0 (IBM Corp., Armonk, NY, USA). Continuous variables were presented as mean \pm standard deviation (SD) or median (interquartile range, IQR) for non-uniformly distributed data. Categorical variables were expressed as frequencies and percentages. Uniformly distributed continuous variables between the groups were compared using Student's t-test. Nominal categorical data between the groups were compared using the chi-square test or Fisher's exact test as appropriate. Non-normally distributed continuous variables were compared using the Mann-Whitney U test. For all statistical tests, a p-value less than 0.05 indicated a statistically significant difference.

Results

Table 1: Demographic details

Age (years)	Total no.
18-27	12
28-37	25
38-47	25
48-57	15
58-67	10
68-77	9
≥ 78	4
Gender	
Male	65
Female	35
Patients with albumin	
Hypoalbuminemia (serum albumin <3.5 g/dl)	50
Normal albumin levels (≥ 3.5 g/dl).	50

Of the 100 patients studied, 65% were male and 35% were female. Pre-operatively, 50 patients had hypoalbuminemia (serum albumin <3.5 g/dl) and 50 patients had normal albumin levels (≥ 3.5 g/dl). Most of the patients belonged to 38-57 years in the present study.

Table 2: Indications for emergency laparotomy

Indications	Number of patients	Percentage
Perforated peptic ulcer	35	35
Acute intestinal obstruction		
Malignant	25	30
Non malignant	5	
Appendiceal perforations	15	15
Incarcerated ventral hernia	8	8
Strangulated hernia	12	12

The most common indication for emergency abdominal surgery was peptic ulcer perforation 35 (35%), followed by acute intestinal obstruction 30 (30%).

Table 3: Post-operative morbidity associated with hypoalbuminemia

Complications	S. albumin <3.5 g/dl (50)	S. albumin >3.5 g/dl (50)	Total (100), p-value
Surgical site infection (%)	19 (38)	10 (20)	P<0.025
Wound dehiscence (%)	8 (16)	3 (3)	P<0.028
Mean length of hospital stays (days)	8.72±5.60	7.35±4.16	P<0.05

Surgical site infection was commonest complication found in 29 (29%) cases. Wound dehiscence was found in 11 (11 %) cases. Mean length of hospital stay was found to be higher i.e. 8.72±5.60 days in patients with hypoalbuminemia, compared to 7.35±4.16 days with those having albumin level ≥ 3.5 g/dl.

Discussion

Malnutrition is prevalent in roughly 30% of surgical patients with gastrointestinal illnesses, and up to 60% of those whose hospital stay has been extended due to post-operative complications. There is substantial proof that patients with malnutrition have a higher risk of complications and death than individuals with adequate nutrition.[14,15] Initially, serum albumin was misinterpreted, and the significance of serum albumin in assessing a person's nutritional makeup was overlooked. When there is acute illness or stress, a reduction in serum albumin occurs due to alterations in hepatic metabolism, and loss of albumin into the interstitium. It also shows an immediate response to surgical stress. As a result, it may meet the criteria for predicting a difficult post-operative course. Abdominal surgeries in

the emergency are very commonly done across hospitals in India. Although surgical and perioperative improvements have reduced postoperative mortality over the last few decades, postoperative morbidity remains high. In addition to the morbidity patients are exposed to, postoperative complications pose a significant financial burden.[16] The magnitude of the metabolic stress response with the long duration of surgery may contribute to the development of postoperative complications. Early identification of patients with nutritional deficiencies and comorbidities and adequate initial resuscitation may help in reducing the postoperative mortality and morbidity of patients.[2,17,18]

Of the 100 patients studied, 65% were male and 35% were female. Pre-operatively, 50 patients had hypoalbuminemia (serum albumin <3.5 g/dl) and 50 patients had normal albumin levels (≥ 3.5 g/dl). Most of the patients belonged to 38-57 years in the present study. Similar results were reported in a previous study by Sharath Kumar et al[19], in which 61.5% were males and 38.5% were females. In another study by

Bhandari et al[20], 66% were males and 34% were females. Males were more commonly involved in road traffic accidents compared to females because males are at high risk while traveling compared to females who mostly stay at home in a developing country like India. This could be the reason for more male patients undergoing exploratory laparotomy in trauma cases. Another reason could be the presence of risk factors usually found in males rather than females, such as alcoholism, smoking, and drug abuse, which are usually responsible for increased trauma and conditions such as pre-pyloric perforation, hollow viscus perforation, and ruptured liver abscess.

The most common indication for emergency abdominal surgery was peptic ulcer perforation 35 (35%), followed by acute intestinal obstruction 30 (30%). The preoperative albumin values had an inverse relation with the length of hospital stay, whereas the reduction in albumin levels had a positive correlation with the length of hospital stay.[21] Truong et al performed a systematic literature review through an electronic search of MEDLINE from PubMed and the Cochrane library found that hypoalbuminemia has a major impact on the length of stay in the hospital, the rate of surgical site infections, the likelihood of enterocutaneous fistulas, and the development of deep vein thrombosis.[22]

Surgical site infection was commonest complication found in 29 (29%) cases. Wound dehiscence was found in 11 (11 %) cases. Mean length of hospital stay was found to be higher i.e. 8.72 ± 5.60 days in patients with hypoalbuminemia, compared to 7.35 ± 4.16 days with those having albumin level ≥ 3.5 g/dl. Hu et al, on 42,483 patients undergoing gastrointestinal surgery found that serum albumin < 3.5 g/dl was associated with wound dehiscence ($p < 0.0001$).[23] Perioperative levels of serum albumin have been shown to be powerful predictors of morbidity and

mortality.[24] Hypoalbuminemia was linked to poor tissue healing, reduced collagen production in surgical wounds or anastomoses, and impaired immunological response.[25] Preoperative serum albumin plays a critical role in determining a patient's postoperative outcome after major surgery.[26]

Conclusion

Preoperative albumin is a useful prognostic predictor for predicting surgical outcome, especially in emergency surgery. It is a simple-to use test. Post-operative morbidity in surgical patients could be decreased by assessing albumin levels in preoperative period, as hypoalbuminemia has adverse relation with surgical outcome. So, it is preferable to have normal albumin perioperatively.

References

1. Beutner KR, Friedman DJ, Forszpaniak C, Andersen PL, Wood MJ. Valaciclovir compared with acyclovir for improved therapy for herpes zoster in immunocompetent adults. *Antimicrobial Agents and Chemotherapy*. 1995 Jul;39(7):1546-53.
2. Thorell A, Nygren J, Ljungqvist O. Insulin resistance: a marker of surgical stress. *Current Opinion in Clinical Nutrition & Metabolic Care*. 1999 Jan 1;2(1):69-78.
3. Rohith VN, Arya SV, Rani A, Chejara RK, Sharma A, Arora JK, Kalwaniya DS, Tolat A, Pawan G, Singh A, Arya SV. Preoperative Serum Albumin Level as a Predictor of Abdominal Wound-Related Complications After Emergency Exploratory Laparotomy. *Cureus*. 2022 Nov 28;14(11).
4. Hall R. Identification of inflammatory mediators and their modulation by strategies for the management of the systemic inflammatory response during cardiac surgery. *Journal of Cardiothoracic and Vascular*

- Anesthesia. 2013 Oct 1;27(5):983-1033.
5. Robert M, Daryl G, Mayes Peter RV. Harper's Illustrated Biochemistry. 26th ed. New York: Mc Graw- Hill Medical; 2003.
 6. Issangya CE, Msuya D, Chilonga K, Herman A, Shao E, Shirima F, Naman E, Mkumbi H, Pyuza J, Mtui E, Sanga LA. Perioperative serum albumin as a predictor of adverse outcomes in abdominal surgery: prospective cohort hospital-based study in Northern Tanzania. BMC surgery. 2020 Dec;20(1):1-7.
 7. Labгаа I, Joliat GR, Kefleyesus A, Mantziari S, Schäfer M, Demartines N, Hübner M. Is postoperative decrease of serum albumin an early predictor of complications after major abdominal surgery? A prospective cohort study in a European centre. BMJ open. 2017 Apr 1;7(4):e013966.
 8. Keller U: Nutritional laboratory markers in malnutrition. J Clin Med. 2019, 8:775. 1
 9. Jäntti T, Tarvasmäki T, Harjola VP, et al.: Hypoalbuminemia is a frequent marker of increased mortality in cardiogenic shock. PLoS One. 2019, 14: e0217006
 10. Paocharoen V, Mingphreudhi S, Lertsithichai P, Euanorasetr C. Preoperative serum albumin level and postoperative septic complications. The Thai Journal of Surgery. 2003 Mar 31;24(1):29-32.
 11. Sharma L, Purohit PM, Pipal D, Kothari S, Srivastava H, Soni A. Serum albumin as predictor of post-operative morbidity and mortality. International Surgery Journal. 2017 Feb 25;4(3):1078-81.
 12. Bhandari TR, Shahi S, Bhandari RS, Lakhey PJ. Preoperative serum albumin level as a predictor of perioperative outcome in patient undergoing major gastrointestinal surgery. Journal of Society of Surgeons of Nepal. 2016 Dec 31;19(2):13-20.
 13. Gibbs J, Cull W, Henderson W, Daley J, Hur K, Khuri SF. Preoperative serum albumin level as a predictor of operative mortality and morbidity: results from the National VA Surgical Risk Study. Archives of surgery. 1999 Jan 1;134(1):36-42.
 14. Sungurtekin H, Sungurtekin U, Balci C, Zencir M, Erdem E. The influence of nutritional status on complications after major intraabdominal surgery. J Am Coll Nutr. 2004;23(3):227-32.
 15. MacFie J. Nutrition and fluid therapy. In Williams N, O'Connell PR, editors. Bailey & Love's Short Practice of Surgery. 25th edition. London: Hodder Arnold. 2008;223.
 16. Vonlanthen R, Slankamenac K, Breitenstein S, Puhan MA, Muller MK, Hahnloser D, Hauri D, Graf R, Clavien PA. The impact of complications on costs of major surgical procedures: a cost analysis of 1200 patients. Annals of surgery. 2011 Dec 1;254(6):907-13.
 17. Mantziari S, Hübner M, Coti-Bertrand P, Pralong F, Demartines N, Schäfer M. A novel approach to major surgery: tracking its pathophysiologic footprints. World journal of surgery. 2015 Nov; 39:2641-51.
 18. Hall R. Identification of inflammatory mediators and their modulation by strategies for the management of the systemic inflammatory response during cardiac surgery. Journal of Cardiothoracic and Vascular Anesthesia. 2013 Oct 1;27(5):983-1033.
 19. Kumar S, Prakash DG, Pottendla VK. Preoperative serum albumin level as a predictor of surgical complications after emergency abdominal surgery. International Surgery Journal. 2019 Jan 28;6(2):361-4.
 20. Bhandari TR, Shahi S, Bhandari RS, Lakhey PJ. Preoperative serum albumin level as a predictor of

- perioperative outcome in patient undergoing major gastrointestinal surgery. *Journal of Society of Surgeons of Nepal*. 2016 Dec 31;19(2):13-20.
21. Kumar DM, Sivakumar S. Evaluation of the role of pre-operative albumin and its post-operative drop in the prediction of outcomes of emergency laparotomy. *International Surgery Journal*. 2020 Mar 26;7(4):1234-7.
 22. Truong A, Hanna MH, Moghadamyeghaneh Z, Stamos MJ. Implications of preoperative hypoalbuminemia in colorectal surgery. *World J Gastrointest Surg*. 2016;8(5):353-62.
 23. Hu WH, Cajas-Monson LC, Eisenstein S, Parry L, Cosman B, Ramamoorthy S. Preoperative malnutrition assessments as predictors of postoperative mortality and morbidity in colorectal cancer: an analysis of ACS-NSQIP. *Nutr J*. 2015;14(1):91.
 24. Anand S, Panda C, Behera MR, Mohanty NK, Thatei C. Role of serum albumin as a prognostic indicator. *Int J Med Res Professionals*. 2016;2(2):123-8.
 25. Lohsiriwat V, Chinswangwatanakul V, Lohsiriwat S, Akaraviputh T, Boonnuch W, Methasade A, et al. Hypoalbuminemia is a predictor of delayed postoperative bowel function and poor surgical outcomes in right-sided colon cancer patients. *Asia Pac J Clin Nutr*. 2007;16(2):213-7.
 26. Garcia-Martinez R, Caraceni P, Bernardi M, Gines P, Arroyo V, Jalan R. Albumin: pathophysiologic basis of its role in the treatment of cirrhosis and its complications. *Hepatology*. 2013;58(5):1836-46.