

A Retrospective Observational Clinicopathological Assessment Post-dated Pregnancy

Priti Kumari¹, Rashmi Kumari², Anupma Sinha³

¹Senior Resident, Department of Obstetrics and Gynaecology, JLNMCH, Bhagalpur, Bihar, India

²Senior Resident, Department of Obstetrics and Gynaecology, JLNMCH, Bhagalpur, Bihar, India

³Associate Professor, Department of Obstetrics and Gynaecology, JLNMCH, Bhagalpur, Bihar, India

Received: 20-03-2023 / Revised: 08-04-2023 / Accepted: 18-05-2023

Corresponding author: Dr. Rashmi Kumari

Conflict of interest: Nil

Abstract

Aim: The aim of the present study was to find out the incidence of maternal complications, perinatal mortality and morbidity in post-dated pregnancies.

Methods: This was a retrospective observational study conducted in the Department of Obstetrics and Gynaecology, JLNMCH, Bhagalpur, Bihar, India for 11 months. Total 100 patients in the antenatal ward and labour room were selected for the study and they were divided into two groups, Control group with Gestational age 37-40 weeks and Study group with Gestational age >40 weeks.

Results: Maximum number of patients belonged to the age group of 25-30 years both in cases (50%) and control group (50%). The maximum number of patients in the study group (80%) belonged to the gestational age of 40-41 weeks while all the controls belonged to 37-40 weeks gestational age. 64% of the patients in study group were primigravida and in the control group 54% were primigravida. The percentage of LSCS was 32% which was higher than in the control group where it was 14%. Incidence of instrumental delivery was also higher in the study group as compared to control group (10% as compared to 6%). Among the indications for LSCS, the most common indication among the study group was acute foetal distress which includes meconium-stained liquor (10%) followed by cephalopelvic disproportion (8%). In the control group, most common indication was non progress of labour (2%) followed by acute foetal distress (0.5%) and non-reactive CTG (1%).

Conclusion: Post-dated pregnancy remains a clinical dilemma for an obstetrician. The choice is between watchful expectancy for labour to start or induction in post-dated patients. According to our study, post-dated pregnancies are related with higher rate of LSCS and instrumental deliveries.

Keywords: Maternal complications, Post datism, Perinatal morbidity, clinic-pathological.

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

Introduction

Post term pregnancy is defined as pregnancy that has extended to or beyond 42 weeks of gestation (294 days), or estimated date of delivery (edd) + 14 days. The terms prolonged pregnancy, postdates and postdatism are synonymously used to describe the same condition. The terms postdate and prolonged pregnancy is ill-defined and best avoided. [1] Post maturity, post maturity syndrome and dysmaturity are not synonymous terms to post term pregnancy. They are often used to describe the features of a neonate who appears to have been in utero longer than 42 weeks of gestation. They describe the effects of intrauterine growth restriction (IUGR) secondary to utero-placental insufficiency encountered in a post term pregnancy. [2]

Several retrospective and relatively small studies have concluded that prolongation of pregnancy beyond term is accompanied by a rise in perinatal morbidity and mortality. [3,4] Abnormalities such as congenital anomalies, oligohydramnios, meconium aspiration, fetal asphyxia, shoulder dystocia, and fetal dysmaturity are commonly observed in these pregnancies. [5,6] The presumed etiology for this rise in perinatal morbidity and mortality is "placental insufficiency." Cunningham et al. [4] concluded that the postterm fetus may outgrow the ability of its placenta to supply nutrients and provide adequate gas exchange and is therefore at risk for adverse outcome resulting from either malnutrition or asphyxia. Indeed, studies of placental histologic features in these pregnancies revealed an increased incidence of placental infarcts, calcification, intervillous thrombosis,

perivillous fibrin deposits, arterial thrombosis, and arterial endarteritis. [7,8] Although the increased morbidity and mortality associated with postterm pregnancy has long been appreciated, most authors have studied gestational age as the only contributing factor. [9-11] The influence of other factors such as maternal age, parity, maternal smoking history, fetal gender, or birth weight has not been adequately evaluated. Additionally, the accuracy of most these earlier studies is limited by the fact that they predate the widespread use of both ultrasonography (for accurate gestational dating) and intensive fetal testing (to establish fetal well-being).

As there is fetal and maternal risk associated with post dated pregnancy, need of induction is more with postdated pregnancy. There is general consideration that perinatal mortality and morbidity controversy is centered on adequacy of detecting different methods for the fetus at risk, the time when testing should be done, the method of monitoring, optimum time for delivery and mode of delivery. The availability of biophysical profile and electronic fetal monitoring can affect the outcome of a given pregnancy. One recent systematic review showed that a policy of labor induction for women with post-dated pregnancy compared with expectant management is associated with fewer perinatal deaths and fewer Caesarean sections. [12]

The aim of the present study was to find out the incidence of maternal complications, perinatal mortality and morbidity in post-dated pregnancies.

Materials and Methods

This was a retrospective observational study conducted in the Department of Obstetrics and gynaecology, JLNMCH, Bhagalpur, Bihar, India for 11 months (Jan 2021 to November 2021) Total 100 patients in the antenatal ward and labour room were selected for the study and they were divided into two groups, Control group with Gestational age 37-40 weeks and Study group with Gestational age >40 weeks.

Inclusion criteria

- Singleton pregnancy
- Cephalic presentation
- Absence of any other maternal complication

Exclusion criteria

- Previous cesarean section
- Gestational hypertension
- Malpresentation
- Abruption
- Placenta previa

All the data regarding the age, parity gestational age, any maternal complications like oligohydromnios, intrauterine growth restriction etc was collected. The maternal outcome was noted in terms of need for cesarean section, postpartum haemorrhage and sepsis. Foetal outcome was noted in terms of intrapartam asphyxia, intrauterine foetal death, admission to neonatal intensive care unit etc.

Results

Table 1: Demographic details and mode of delivery

Age (Years)	Number of Cases (%)	Number of Controls (%)
Below -25	14 (28%)	16 (32%)
25-30	24 (48%)	25 (50%)
Above 30	12 (24%)	9 (18%)
Total	50	50
Mean ± SD	26.4 ± 3.48	28.2 ± 3.80
Period of gestation		
37-40 weeks	0	50 (100%)
40-41 weeks	40 (80%)	0
41-42 weeks	10 (20%)	0
Total	50	50
Parity		
Primigravida	32 (64%)	27 (54%)
Multigravida	18 (36%)	23 (46%)
Total	50	50
Type of delivery		
NVD	29 (58%)	40 (80%)
Instrumental delivery	5 (10%)	3 (6%)
LSCS	16 (32%)	7 (14%)
Total	50	50

Maximum number of patients belonged to the age group of 25-30 years both in cases (50%) and

control group (50%). The maximum number of patients in the study group (80%) belonged to the

gestational age of 40-41 weeks while all the controls belonged to 37-40 weeks gestational age. 64% of the patients in study group were primigravida and in the control group 54% were primigravida. The percentage of LSCS was 32%

which was higher than in the control group where it was 14%. Incidence of instrumental delivery was also higher in the study group as compared to control group (10% as compared to 6%).

Table 2: Distribution of cases and controls according to the indication of LSCS

Indication of LSCS	Number of Cases (%)	Number of Controls (%)
Acute foetal distress/MSL	5 (10%)	1 (0.5%)
Failed induction	3 (6%)	0
Nonprogress of labour	2 (4%)	4 (2%)
Non-reactive CTG	2 (4%)	2 (1%)
CPD	4 (8%)	0
Total	16	7

Among the indications for LSCS, the most common indication among the study group was acute foetal distress which includes meconium-stained liquor (10%) followed by cephalopelvic disproportion (8%). In the control group, most common indication was non progress of labour (2%) followed by acute foetal distress (0.5%) and non-reactive CTG (1%).

Table 3: Distribution of cases and controls according to maternal complications

Maternal complication	Number of cases	Number of controls
LSCS	13 (26%)	7 (14%)
PPH	7 (14%)	2 (4%)
Sepsis	4 (8%)	1 (2%)
Total	24	10

Maternal complications like LSCS, PPH and sepsis all were higher in the study group as compared to the control group.

Table 4: Distribution of cases and controls according to the foetal outcome

Foetal outcome	Number of Cases (%)	Number of Controls (%)
No asphyxia	33 (66%)	41 (82%)
Fetal asphyxia (APGAR score<6/10)	6 (12%)	3 (6%)
Admission to NICU	7 (14%)	6 (12%)
IUD	4 (2%)	0
Total	50	50

12% of infants in the study group had asphyxia as compared to only 6% in the control group. 14% infants of the study group had to be admitted to the NICU as compared to 12% in the control group. 2% was the percentage of intrauterine deaths in the study group as compared to none in the control group.

Discussion

Fernandos Arias defined prolonged pregnancy as those pregnancies advancing beyond the expected date of delivery (EDD). [13] Prolongation of pregnancy complicates up to 10% of all pregnancies and carries increased risk to mother and fetus. [14,15] Post term perinatal mortality is greater than that of term pregnancy in almost all studies reviewed. [16] The growth and survival of most post-dated infants suggests that the placenta uncommonly deteriorates with increasing length of gestation; thus the changes seen in fetuses afflicted with post maturity syndrome may not be explained by placental findings alone. Vorherr described critical reductions of fetal oxygen supply after 43rd

week of gestation by cord blood oxygen content determinations. [16] The combination of continued fetal growth and arrested placental growth may lead to situation of decreasing placental nutrient reserve, compromised fetal circulation and eventually fetal distress. However, a recent electron microscopy study of placental changes in prolonged pregnancy suggests that the uteroplacental ischemia and not placental aging may be more important in genesis of post maturity syndrome. [17]

Maximum number of patients belonged to the age group of 25-30 years both in cases (50%) and control group (50%). The maximum number of patients in the study group (80%) belonged to the gestational age of 40-41 weeks while all the controls belonged to 37-40 weeks gestational age. 64% of the patients in study group were primigravida and in the control group 54% were primigravida. Similar studies by Mahapatro [18] and Eden et al [19] have shown the mean age to be 24.19 ± 3.30 and 25.8 years respectively. 62% of the patients in study group were primigravida

which is similar to Mahapatro¹⁸ and Alexander et al's study. [20] The percentage of LSCS was 32% which was higher than in the control group where it was 14%. Incidence of instrumental delivery was also higher in the study group as compared to control group (10% as compared to 6%). In a similar study by Mahapatro¹⁸ the rate of LSCS was found to be 28.9% and that of instrumental delivery was 5.72%. In study by Singhal et al [21] the rate of LSCS was found to be 14.7% and that of instrumental delivery was 8.6%. Davinder et al [22] study showed the rate of instrumental delivery as 10.35%.

Among the indications for LSCS, the most common indication among the study group was acute foetal distress which includes meconium stained liquor (10%) followed by cephalopelvic disproportion (8%). In the control group, most common indication was non progress of labour (2%) followed by acute foetal distress (0.5%) and non-reactive CTG (1%). Bhriegu R et al [23] in their study also found that Meconium stained liquor with fetal distress was the most common indication for LSCS (23.5%) and in Mahapatro's study [18], again fetal distress was found to be the most common indication for LSCS (65.5%).

Maternal complications like LSCS, PPH and sepsis all were higher in the study group as compared to the control group. 12% of infants in the study group had asphyxia as compared to only 6% in the control group. 14% infants of the study group had to be admitted to the NICU as compared to 12% in the control group. 2% was the percentage of intrauterine deaths in the study group as compared to none in the control group. According to study done by Aaron, estimated rates of maternal complications increase beyond 40 weeks of gestation. Beyond that the rates of operative vaginal delivery, 3rd or 4th degree perineal laceration and chorioamnionitis all increases. ($p < 0.001$), and rates of postpartum hemorrhage, endometritis and primary caesarean delivery increased at 41 weeks of gestation. [24]

Conclusion

Postdated pregnancy remains a clinical dilemma for an obstetrician. The choice is between watchful expectancy for labour to start or induction in postdated patients. According to our study, postdated pregnancies are related with higher rate of LSCS and instrumental deliveries. Foetal distress and meconium stained liquor are significantly higher in post-dated pregnancies. Similarly, maternal and perinatal complications like postpartum haemorrhage, sepsis, admission to NICU, low Apgar scores, foetal asphyxia and intrauterine deaths are also higher in post-term pregnancies. Therefore it is recommended to not prolong the post-dated pregnancies and their

induction must be soon after the expected date is crossed to prevent the above mentioned complications.

References

1. aCOG (american College of Obstetricians and Gynecologists). Management of Postterm Pregnancy. aCOG Practice bulletin no. 55. Obstet Gynecol. 2004; 104:639-46.
2. Shime J, Librach CL, Gare DJ, Cook CJ. The influence of prolonged pregnancy on infant development at one and two years of age: a prospective controlled study. American journal of obstetrics and gynecology. 1986 Feb 1;154 (2):341-5.
3. NAEYE RL. Causes of perinatal mortality excess in prolonged gestations. American journal of epidemiology. 1978 Nov 1;108(5) 429-33.
4. Clifford SH. Postmaturity—with placental dysfunction: Clinical syndrome and pathologic findings. The Journal of pediatrics. 1954 Jan 1;44(1):1-3.
5. Chamberlain PF, Manning FA, Morrison I, Harman CR, Lange IR. Ultrasound evaluation of amniotic fluid volume: I. The relationship of marginal and decreased amniotic fluid volumes to perinatal outcome. American journal of obstetrics and gynecology. 1984 Oct 1;150(3): 245-9.
6. Cunningham FG, McDonald PC, Gant NF, et al., editors. Williams' obstetrics. 20th ed. Stamford (CT): Appleton & Lange; 1997. p. 827-37.
7. Vorherr H. Placental insufficiency in relation to postterm pregnancy and fetal postmaturity: evaluation of fetoplacental function; management of the postterm gravida. American journal of obstetrics and gynecology. 1975 Sep 1;123(1):67-103.
8. Thliveris JA, Baskett TF. Fine structure of the human placenta in prolonged pregnancy. Obstetrical & Gynecological Survey. 1979 Aug 1;34(8):573-5.
9. Browne JM. Postmaturity. JAMA. 1963 Dec 21;186(12):1047-52.
10. Stewart Jr HL. Duration of pregnancy and postmaturity. Journal of the American Medical Association. 1952 Mar 29;148(13):1079-83.
11. Saito M, Yazawa K, Hashiguchi A, Kumasaka T, Nishi N, Kato K. Time of ovulation and prolonged pregnancy. American journal of obstetrics and gynecology. 1972 Jan 1;112 (1):31-8.
12. Gülmezoglu AM, Crowther CA, Middleton P, Heatley E. Induction of labour for improving birth outcomes for women at or beyond term. Cochrane database of systematic reviews. 2012(6).

13. Arias F, Bhide AG, Arulkumaran S, Damania K, Daftary SN, editors. Practical Guide to High-Risk Pregnancy and Delivery-E-Book: A South Asian Perspective. Elsevier health sciences; 2008 Jul 15.
14. Olesen AW, Olsen J, Westergaard JG. Perinatal and maternal complications related to post-term delivery: A national register-based study 1978-93.
15. Norwitz ER, Snegovskikh VV, Caughey AB. Prolonged Pregnancy. When Should We Intervene? Clinical obstetrics and gynecology. 2007 Jun 1;50(2):547-57.
16. Vorherr H. Placental insufficiency in relation to postterm pregnancy and fetal postmaturity: evaluation of fetoplacental function; management of the postterm gravida. American journal of obstetrics and gynecology. 1975 Sep 1;123(1):67-103.
17. Jones JP, Fox H. Ultrastructure of the placenta in prolonged pregnancy. J Pathol. 1978; 126 (3):173-9.
18. Mahapatro A, Samal S. Fetomaternal outcome in pregnancy beyond 40 weeks. Int J Pharm Bio Sci. 2015;6(2):53-8.
19. Eden RD, Gergely RZ, Schifrin BS, Wade ME. Comparison of antepartum testing schemes for the management of the postdate pregnancy. American journal of obstetrics and gynecology. 1982 Nov 15;144(6):683-92.
20. Alexander JM, McIntire DD, Leveno KJ. Forty weeks and beyond: pregnancy outcomes by week of gestation. Obstetrics & Gynecology. 2000 Aug 1;96(2):291-4.
21. Singhal P, Sharma A, Jain D, Panday V. Fetomaternal outcome following postdate pregnancy. A prospective study. J Obstet Gynecol India. 2001; 51:89-93.
22. Kaur D, Saini AS, Kaur J. Maternal and fetal outcome in postdated pregnancies. J Obstet Gynecol India. 1997; 47:331-4.
23. Bhriegu R, Agrawal M, Hariharan C. Assessment of maternal and perinatal outcome in postdated pregnancy. Journal of Datta Meghe Institute of Medical Sciences University. 2017 Jan 1;12(1):35-40.
24. Caughey AB, Stotland NE, Washington AE, Escobar GJ. Maternal Complications of Pregnancy Increase Beyond 40 Weeks' Gestation. Am J Obstet Gynecol. 2007; 196:155.