

A Hospital Based 1-Year Prospective Study to Estimate Prevalence and Severity of DR in Recently Diagnosed Type 2 DM Patients**Rajnee Sinha¹, Madhulika Sinha², Rajeshwar Rao³, Gyan Bhaskar⁴**¹Senior Resident, Department of Ophthalmology, IGIMS, Patna, Bihar, India²Senior Resident, Department of Ophthalmology, IGIMS, Patna, Bihar, India³Senior Resident, Department of Nephrology, IGIMS, Patna, Bihar, India⁴Professor, Department of Ophthalmology, IGIMS, Patna, Bihar, India

Received: 10-12-2023 Revised: 20-01-2024. Accepted: 17-02-2024

Corresponding Author: Dr. Madhulika Sinha

Conflict of interest: Nil

Abstract**Aim:** The aim of the present study was to estimate prevalence and severity of DR in recently diagnosed type 2 DM patients.**Material & Methods:** A study was conducted in the Department of Ophthalmology for 1 year. This observational, prospective study was conducted in patients who visited the diabetic clinic and recently diagnosed (less than 3 months from first diagnosis) for type 2 DM. In present study total 100 patients were evaluated for diabetic retinopathy.**Results:** In present study total 100 patients were evaluated for diabetic retinopathy. Most of patients were from 61-70 years (25%), followed by 51-60 years (22%) and 41-50 years (20%) age group. Male patients (58%) were more than female patients (42%). We noted 10% prevalence of diabetic retinopathy in study patients. Prevalence was more in males (66.66%) as compared to females (33.33%). In present study diabetic retinopathy was noted in 15 patients, divided as mild nonproliferative diabetic retinopathy (66.66%), Moderate nonproliferative diabetic retinopathy (20%), Severe nonproliferative diabetic retinopathy (6.66%) and proliferative diabetic retinopathy (PDR) (6.66%). Macular edema was noted in 4 patients. 60% of them had mild macular edema while 20% each had moderate and severe macular edema.**Conclusion:** Screening for diabetic retinopathy is important for newly diagnosed diabetic patients. A systematic screening program in the community is needed for early detection and to reduce blindness in diabetic patients.**Keywords:** Diabetic Retinopathy, Type 2 Diabetes Mellitus, Non-proliferative, Macular EdemaThis is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.**Introduction**

Diabetic retinopathy (DR) is the leading cause of visual impairment in the Western world, particularly among persons of working age. [1,2] It is estimated that DR develops in more than 75% of diabetic patients within 15 to 20 years of diagnosis of diabetes. [3,4] Several epidemiologic studies have provided valuable information on the prevalence of DR in Western countries that is useful for identifying subgroups at risk and for the planning of public health policies. [5] The Eye Diseases Prevalence Research Group collates data on eye diseases in the United States, and provides information on the health services burden due to eye diseases, including DR. [5] However, there is a paucity of data on the prevalence of diabetes-related eye diseases in developing countries such as India, which in fact has the largest number of diabetic individuals in the world.

According to the latest World Health Organization (WHO) report, India has 31.7 million diabetic subjects, and the number is expected to increase to a staggering 79.4 million by 2030. [6] Furthermore, type 2 diabetes in Indians differs from that in Europeans in several aspects: The onset is at a younger age, [7] obesity is less common, [8] and genetic factors appear to be stronger. [9] These clinical differences and the rising prevalence of diabetes in India [10] warrant well-conducted epidemiologic studies on diabetes-related complications in this population, to assess the health services burden due to diabetes.

Diabetic retinopathy (DR) is the leading cause of visual impairment among working-age adults; however, increasing awareness and the early identification of DM has ameliorated the problem somewhat. [11] Most standard protocols for the

screening of DR recommend annual or biannual dilated retinal examinations for all patients with diabetes. [12]

Therefore, in present study we aimed to estimate prevalence and severity of DR in recently diagnosed type 2 DM patients

Material & Methods

A study was conducted in the Department of Ophthalmology, IGIMS, Patna, Bihar, India for 1 year. This observational, prospective study was conducted in patients who visited the diabetic clinic and recently diagnosed (less than 3 months from first diagnosis) for type 2 DM. In present study total 100 patients were evaluated for diabetic retinopathy.

Inclusion Criteria:

- Patients who were recently diagnosed (less than 3 months from first diagnosis) for type 2 DM, willing to participate

Exclusion Criteria:

- Patients with mature cataracts and hazy media, whose fundi could not be examined.
- Patients with a history of exposure to radiation, hypertensive retinopathy without DM, sickle cell disease and pheochromocytoma.

- Patients who were not willing to participate.

Methodology

Study was explained and a written informed consent was taken. Patients underwent history taking (age, sex, medical history, smoking), general physical examination and routine ophthalmological examination was done. The pupils of both eyes were dilated by using a mydriatic agent (Tropicamide with phenylephrine eye drops). Distant direct ophthalmoscopy, direct ophthalmoscopy and binocular indirect ophthalmoscopy were done. Binocular indirect ophthalmoscopy was done with a 20 D lens with the patient in supine position. Findings were noted and patients were categorized according to findings; whether diabetic retinopathy was present or absent. If present, retinopathy was classified according to early treatment of diabetic retinopathy study (ETDRS) classification. [13] Presence of diabetic macular oedema was noted. If present, it was further classified into clinically significant (CSME) or non-significant. [13]

Statistical Analysis

Data was collected, entered in Microsoft excel sheet and analysed by descriptive statistics

Results

Table 1: Age and gender distribution

Age in years	Total N=100	Diabetic Retinopathy N=15
≤ 40	10 (10%)	2 (13.33%)
41-50	20 (20%)	2 (13.33%)
51-60	22 (22%)	3 (20%)
61-70	25 (25%)	3 (20%)
71-80	17 (17%)	3 (20%)
>80	6 (6%)	2 (13.33%)
Total	100	15
Gender		
Male	58 (58%)	10 (66.66%)
Female	42 (42%)	5 (33.33%)

In present study total 100 patients were evaluated for diabetic retinopathy. Most of patients were from 61-70 years (25%), followed by 51-60 years (22%) and 41-50 years (20%) age group. Male patients (58%) were more than female patients (42%). We noted 10% prevalence of diabetic retinopathy in study patients. Prevalence was more in males (66.66%) as compared to females (33.33%).

Table 2: Diabetic retinopathy and macular edema

	N	%
Diabetic retinopathy (N=15)		
Mild NPDR	10	66.66
Moderate NPDR	3	20
Severe NPDR	1	6.66
PDR	1	6.66
Total	15	100
Macular edema (N=5)		
Mild	3	60
Moderate	1	20

Severe	1	20
--------	---	----

In present study diabetic retinopathy was noted in 15 patients, divided as mild nonproliferative diabetic retinopathy (66.66%), Moderate nonproliferative diabetic retinopathy (20%), Severe nonproliferative diabetic retinopathy (6.66%) and proliferative diabetic retinopathy (PDR) (6.66%). Macular edema was noted in 4 patients. 60% of them had mild macular edema while 20% each had moderate and severe macular edema.

Discussion

Diabetes mellitus (DM) has exemplified its presence globally. In India, 69.1 million are diabetics as of 2015 compared to 18 million in 1995. [14,15] At this rate, crossing one billion as estimated by the International Diabetes Federation Association and World health organization seems obvious. [15] Pan India studies on the prevalence of Diabetic retinopathy (DR) are limited, few initial studies, such as The Chennai Urban Rural Epidemiology Study and Arvind comprehensive eye study in rural population, concluded the prevalence of DR as 17.8% and 10.8% in diabetics, respectively. [16,17] DR is a common microvascular complication, an important complication among adults and elderly. DR is of two types, nonproliferative diabetic retinopathy (NPDR) and proliferative diabetic retinopathy (PDR). Severity of NPDR depends on microaneurysms, hemorrhages, cotton wool spots, and beading of veins and can progress to PDR. Inherent characteristic of PDR is neovascularization. Its either PDR or diabetic maculopathy that is responsible for vision loss. Chronic hyperglycemia inflicts profoundly in development and progression of DR by stimulating multiple mechanisms such as polyol pathway, enzymatic glycation, protein kinase C activation, hemodynamic changes, and renin angiotensin aldosterone pathway [18] and Type 2 DM accounts for more than 90% of the diabetic population worldwide. [19]

Proliferative diabetic retinopathy (PDR) is a treatable cause of severe visual loss in people with diabetes. If left untreated, most eyes with low-risk PDR characterized by mild to moderate retinal or optic disc neovascularization progress to high-risk PDR with increasing retinal or disc neovascularization. Risk factors for the development of retinopathy and visual loss include type of diabetes, duration of diabetes, poor glycemic control, poor blood pressure control, deranged lipid profile, obesity, obstructive sleep apnea (OSA), pregnancy and anaemia. The duration of diabetes is probably the strongest predictor for development and progression of retinopathy. [4] In present study total 100 patients were evaluated for diabetic retinopathy. Most of patients were from 61-70 years (25%), followed by 51-60 years (22%) and 41-50 years (20%) age group. Male patients (58%) were

more than female patients (42%). We noted 10% prevalence of diabetic retinopathy in study patients. Prevalence was more in males (66.66%) as compared to females (33.33%). Sosale A et al [20] studied 4600 (males 67%, females 33%) newly diagnosed patients with T2D, majority were from the age group 41-50 years (40%).

In present study diabetic retinopathy was noted in 15 patients, divided as mild nonproliferative diabetic retinopathy (66.66%), Moderate nonproliferative diabetic retinopathy (20%), Severe nonproliferative diabetic retinopathy (6.66%) and proliferative diabetic retinopathy (PDR) (6.66%). Macular edema was noted in 4 patients. 60% of them had mild macular edema while 20% each had moderate and severe macular edema. Hao Z et al [21] enrolled 947 patients in their study and were divided into two groups according to whether they were diagnosed with DR. There was no statistically significant difference between the two groups in sex, age, hypertension, DM diagnosed age, family history of diabetes, drinking history and HbA1c. BMI was significantly higher in DR patients. newly diagnosed T2DM subjects, DR is associated with reduced beta-cell responsiveness, resulting from beta-cell failure rather than insulin resistance, leading to fasting and postprandial hyperglycemia and hypoinsulinemia. [22] The most effective way of preventing the risk of vision loss from diabetes mellitus is patient education about the need for screening for retinopathy even in the absence of any visual complaints. Therefore, early screening is strongly recommended for all newly diagnosed T2DM patients [23]

Conclusion

Screening for diabetic retinopathy is important for newly diagnosed diabetic patients. A systematic screening program in the community is needed for early detection and to reduce blindness in diabetic patients.

References

1. Klein R, Klein BE, Moss SE. Visual impairment in diabetes. *Ophthalmology*. 1984 Jan 1;91(1):1-9.
2. Vision Problems in the United States. Data Analysis: Definitions, Data Sources, Detailed Data Tables, Analysis, Interpretation. New York: National Society to Prevent Blindness; 1989;1-46.
3. Dwyer MS, Melton III LJ, Ballard DJ, Palumbo PJ, Trautmann JC, Chu CP. Incidence of diabetic retinopathy and blindness: a population-based study in Rochester, Minnesota. *Diabetes Care*. 1985 Jul 1;8(4): 316-22.

4. Klein R, Klein BE, Moss SE, Davis MD, DeMets DL. The Wisconsin Epidemiologic Study of Diabetic Retinopathy: III. Prevalence and risk of diabetic retinopathy when age at diagnosis is 30 or more years. *Archives of ophthalmology*. 1984 Apr 1;102(4):527-32.
5. Kempen JH, O'Colmain BJ, Leske MC, Haffner SM, Klein R, Moss SE, Taylor HR, Hamman RF. The prevalence of diabetic retinopathy among adults in the United States. *Archives of ophthalmology (Chicago, Ill.: 1960)*. 2004 Apr 1;122(4):552-63.
6. Wild S, Roglic G, Green A, Sicree R, King H. Global prevalence of diabetes: estimates for the year 2000 and projections for 2030. *Diabetes care*. 2004 May 1;27(5):1047-53.
7. Mohan V, Ramachandran A, Snehalatha C, Mohan R, Bharani G, Viswanathan M. High prevalence of maturity-onset diabetes of the young (MODY) among Indians. *Diabetes care*. 1985 Jul 1;8(4):371-4.
8. Joshi SR. Metabolic syndrome-emerging clusters of the Indian phenotype. *Journal-Association of Physicians of India*. 2003 May 1;51:445-6.
9. Viswanathan M, Mohan V, Snehalatha C, Ramachandran A. High prevalence of type 2 (non-insulin-dependent) diabetes among the offspring of conjugal type 2 diabetic parents in India. *Diabetologia*. 1985 Dec;28:907-10.
10. Mohan V, Shanthirani CS, Deepa R. Glucose intolerance (diabetes and IGT) in a selected South Indian population with special reference to family history, obesity and lifestyle factors--the Chennai Urban Population Study (CUPS 14). *The Journal of the Association of Physicians of India*. 2003 Aug 1;51:771-.
11. Tien YW, MWAMBURI M, KLEIN R, LARSEN M, FLYNN H, HERNANDEZ-MEDINA M, RANGANATHAN G, WIROSTKO B, PLEIL A, MITCHELL P. Rates of Progression in Diabetic Retinopathy During Different Time Periods: A systematic review and meta-analysis. *Diabetes care*. 2009;32(12):2307-13.
12. Flaxel CJ, Adelman RA, Bailey ST, Fawzi A, Lim JJ, Vemulakonda GA, Ying GS. Diabetic retinopathy preferred practice pattern®. *Ophthalmology*. 2020 Jan 1;127(1):P66-145.
13. Ryan SJ. *Medical retina*. 4th ed. Baltimore: Mosby; Chapter 67:1276.
14. Raman R, Ganesan S, Pal SS, Kulothungan V, Sharma T. Prevalence and risk factors for diabetic retinopathy in rural India. *Sankara Nethralaya Diabetic Retinopathy Epidemiology and Molecular Genetic Study III (SN-DREAMS III)*, report no 2. *BMJ Open Diabetes Research and Care*. 2014 Jun 1;2(1): e000005.
15. Raman R, Rani PK, Racheppalle SR, Gnanamoorthy P, Uthra S, Kumaramanickavel G, Sharma T. Prevalence of diabetic retinopathy in India: Sankara Nethralaya diabetic retinopathy epidemiology and molecular genetics study report 2. *Ophthalmology*. 2009 Feb 1;116(2):311-8.
16. Deepa M, Pradeepa R, Rema M, Anjana RM, Deepa R, Shanthirani S, Mohan V. The Chennai Urban Rural Epidemiology Study (CURES)--study design and methodology (urban component)(CURES-I). *The journal of the association of physicians of India*. 2003 Sep 1;51:863-70.
17. Nirmalan PK, Katz J, Robin AL, Tielsch JM, Namperumalsamy P, Kim R, Narendran V, Ramakrishnan R, Krishnadas R, Thulasiraj RD, Suan E. Prevalence of vitreoretinal disorders in a rural population of southern India: The aravind comprehensive eye study. *Archives of ophthalmology*. 2004 Apr 1;122(4):581-6.
18. Tarr JM, Kaul K, Chopra M, Kohner EM, Chibber R. Pathophysiology of diabetic retinopathy. *International Scholarly Research Notices*. 2013;2013.
19. Jain R, Jain P, Jain P. A review on treatment and prevention of diabetes mellitus. *Int J Curr Pharm Res*. 2016;8(3):16-8.
20. Sosale A, Kumar KP, Sadikot SM, Nigam A, Bajaj S, Zargar AH, Singh SK. Chronic complications in newly diagnosed patients with type 2 diabetes mellitus in India. *Indian journal of endocrinology and metabolism*. 2014 May;18(3):355.
21. Yadav, S. ., Gupta, D. K., Patil, P. H., Tiwari, A. ., & soni, P. . (2023). To Study the Pharmacotherapy of Diabetes Mellitus Type 2 Patient in Echo Health Care & Research Centre, Indore". *Jour Med Resh and Health Sci*, 6(2),2389–2397. <https://doi.org/10.52845/JMRHS/2023-6-2-1>
22. Hao Z, Huang X, Qin Y, Li H, Tian F, Xu R, Chang B, Shao H. Analysis of factors related to diabetic retinopathy in patients with newly diagnosed type 2 diabetes: a cross-sectional study. *BMJ open*. 2020 Feb 1;10(2):e032095.
23. Roy Chowdhury S, Thomas RL, Dunseath GJ, Peter R, Rees DA, North RV, Luzio SD, Owens DR. Diabetic retinopathy in newly diagnosed subjects with type 2 diabetes mellitus: contribution of β -cell function. *The Journal of Clinical Endocrinology & Metabolism*. 2016 Feb 1;101(2):572-80.