

A Retrospective Assessment of Common Bacteria Associated with UTI Cases and Their Antibiotic Susceptibility Pattern

Manju Kumari¹, Jitendra Kumar²

¹Tutor, Department of Pharmacology, ANMMCH, Gaya, Bihar, India

²Associate Professor, Department of Pharmacology, ANMMCH, Gaya, Bihar, India

Received: 02-01-2024 / Revised: 10-02-2024 / Accepted: 25-03-2024

Corresponding Author: Dr. Jitendra Kumar

Conflict of interest: Nil

Abstract

Aim: The aim of the present study was to investigate the common bacteria associated with UTI cases and their antibiotic susceptibility pattern.

Methods: A retrospective study was done for one year (January 2021 to December 2021) in Department of Pharmacology, ANMMCH, Gaya, Bihar, India. The study included all in-patients of either gender above 18 years of age with a diagnosis of UTI. Data was collected from the patient's case record files, which were retrieved from the medical records department of our hospital.

Results: Out of 120 significant isolates, gram-negative organisms accounted for 81%. Fungal isolates were seen only in 2% of cases. E.coli was the most predominant uropathies isolated followed by Klebsiella spp, Enterococcus faecalis and Acinetobacter spp. Other uropathies included MRSA bacteria, Staphylococcus aureus, Candida spp, Citrobacter Freunde, Pseudomonas spp and Streptococcus spp. E.coli, showed considerable resistance to ampicillin, ceftriaxone, Co-Trimoxazole and fluoroquinolones. Low level of resistance was observed with Nitrofurantoin, piperacillin+ tazobactam and Ticarcillin. Klebsiella spp, the second most common uropathies showed high level of resistance with ampicillin, cephalexin, cefuroxime, ceftriaxone and Nitrofurantoin.

Conclusion: Most of the identified bacteria were resistant to several of the popular antibiotics used in clinical settings. Consequently, it is vital to prescribe antibiotics rationally both before and after culture reports. Pattern needs to be studied further and the outcome should be communicated to the prescribing physicians in that area.

Keywords: antibiotic resistance, Culture, UTI, uropathies

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

Introduction

Urinary tract infections (UTIs) are a major public health problem in terms of morbidity and financial cost, and incur the highest total health care cost among urological diseases, exceeding that of chronic renal failure even when renal dialysis and renal transplantation are included. [1] UTI represents one of the most common diseases encountered in medical practice today with an estimated 150 million UTIs per annum worldwide. [2] Although UTIs occur in both men and women, clinical studies suggest that the overall prevalence of UTI is higher in women. Uncomplicated UTIs in healthy women have an incidence of 50/1000/year. [3] An estimated 50% of women experience at least one episode of UTI at some point in their lifetime and between 20% and 40% of women have recurrent episodes. [4,5] Approximately 20% of all UTIs occur in men. [6]

Most episodes of UTI are caused by Escherichia coli (up to 85%) and Staphylococcus saprophyticus (up to 10%), while Klebsiella pneumoniae and Proteus

species account for most of the remaining infections. [7] The introduction of antimicrobial therapy has contributed significantly to the management of UTIs. However the main problem with current antibiotic therapies is the rapid emergence of antimicrobial resistance in hospitals and the community. [8] The resistance pattern of community acquired uropathies has not been extensively studied in the Indian subcontinent. [9-11]

UTI is the most common in patients with diabetes and in those with structural and neurological abnormalities which interfere with urinary outflow. Nosocomial UTI is also common following catheterization and cystoscopy. The manifestations of UTI may vary from mild asymptomatic cystitis to pyelonephritis and septicemia. [12] Gram-negative organisms are the most common pathogens involved in UTI, but almost all known pathogens have been incriminated as possible causative agents for UTI. [13,14] Treatment of UTI constitutes a great portion of prescription of antibiotics. Urinary pathogens

have shown a changed pattern of susceptibility to antibiotics, resulting in an increase in resistance to commonly used antibiotics. [15] The distribution of uropathogens and their susceptibility pattern to antibiotics vary regionally. [16,17]

The aim of the present study was to investigate the common bacteria associated with UTI cases and their antibiotic susceptibility pattern.

Materials and Methods

A retrospective study was done for one year (January 2021 to December 2021) in Department of Pharmacology, ANMMCH, Gaya, Bihar, India. The study included all in-patients of either gender above

18 years of age with a diagnosis of UTI. Data was collected from the patient's case record files, which were retrieved from the medical records department of our hospital.

All relevant data regarding the type of clinical presentation, demographic distribution, associated risk factors, co-morbid conditions, microbiology reports, causative uropathies and its antibiotic resistance pattern were documented into a proforma sheet prepared beforehand. The data was entered, stored and evaluated using Microsoft excel 2016. Data was calculated according to the percentages.

Results

Table 1: Antibiotic resistance pattern of gram-negative

| Antimicrobial Agents | <i>E.coli</i> | <i>Klebsiella spp</i> | <i>Acinetobacter spp</i> | <i>Citrobacter freundii</i> | <i>Pseudomonas spp</i> |
|-----------------------------|---------------|-----------------------|--------------------------|-----------------------------|------------------------|
| | R (%) | R (%) | R (%) | R (%) | R (%) |
| Ampicillin | 87 | 96 | 93 | 95 | 100 |
| Amoxicillin-Clavulanic Acid | 67.8 | 46.4 | 76 | 93 | 95 |
| Ampicillin/Salbactam | | | | | |
| Cefalexin | 84.2 | 92 | 100 | 100 | 100 |
| Cefuroxime | 78.9 | 72 | 100 | 50 | 100 |
| Ceftriaxone | 77.6 | 76 | 100 | 50 | 100 |
| Ciprofloxacin | 84.0 | 50 | 100 | 50 | 100 |
| Levofloxacin | 78.2 | 34.6 | 100 | 50 | 100 |
| Cotrimoxazole | 46.4 | 46.4 | 100 | 100 | 100 |
| Gentamicin | 51.7 | 55.6 | 100 | 50 | 50 |
| Amikacin | 7.0 | 17.8 | 100 | 50 | 100 |
| Nitrofurantoin | 6.4 | 88.2 | 100 | 0.0 | 100 |
| Piperacillin Tazobactam + | 7.9 | 30 | 50 | 50 | 0.0 |
| Meropenam | 6.6 | 9.2 | 76 | 0.0 | 0.0 |

Out of 120 significant isolates, gram-negative organisms accounted for 81%. Fungal isolates were seen only in 2% of cases. *E.coli* was the most predominant uropathies isolated followed by *Klebsiella spp*, *Enterococcus faecalis* and

Acinetobacter spp. Other uropathies included MRSA bacteria, *Staphylococcus aureus*, *Candida spp*, *Citrobacter freundii*, *Pseudomonas spp* and *Streptococcus spp*.

Table 2: Antibiotic resistance pattern of gram-Positive

| ANTIMICROBIAL AGENTS | <i>Enterococcus faecalis</i> | <i>Staphylococcus aureus</i> | MRSA | <i>Streptococcus spp</i> |
|-----------------------------|------------------------------|------------------------------|-------|--------------------------|
| | R (%) | R (%) | R (%) | R (%) |
| Ampicillin | 25 | 34.4 | 100 | 0.0 |
| Amoxicillin-clavulanic acid | 18.2 | 0.0 | 100 | 0.0 |
| Cephalexin | 100 | - | 100 | 0.0 |
| Cefuroxime | 100 | 0.0 | 100 | 0.0 |
| Ceftriaxone | 100 | 0.0 | 100 | 0.0 |
| Ciprofloxacin | 29.7 | 100 | 100 | 0.0 |
| Levofloxacin | 29.7 | 0.0 | 66.7 | 0.0 |
| Cotrimoxazole | 57.1 | 0.0 | 66.7 | 0.0 |
| Gentamicin | 72.5 | 0.0 | 0.0 | 100 |
| Amikacin | 64.6 | 0.0 | 66.7 | 100 |

| | | | | |
|--------------------------|------|-----|-----|-----|
| Nitrofurantoin | 12.5 | - | 0.0 | 0.0 |
| Piperacillin+ Tazobactam | 0.0 | - | - | - |
| Imipenem | 14.6 | - | - | - |
| Meropenem | 34.4 | - | - | - |
| Vancomycin | 0.0 | 0.0 | 0.0 | - |
| Linezolid | 0.0 | 0.0 | 0.0 | - |
| Teicoplanin | 0.0 | 0.0 | 0.0 | - |
| Azithromycin | - | - | 100 | 0.0 |

E. coli, showed considerable resistance to ampicillin, ceftriaxone, Co-Trimoxazole and fluoroquinolones. Low level of resistance was observed with Nitrofurantoin, piperacillin+ tazobactam and Ticarcillin. *Klebsiella spp*, the second most common uropathies showed high level of resistance with ampicillin, cephalexin, cefuroxime, ceftriaxone and Nitrofurantoin.

Discussion

Any infection of the urinary tract involving either the kidneys, ureters, urinary bladder, or the urethra is termed as urinary tract infection (UTI). In both community and hospital-acquired infections, urinary tract infection (UTI) is a prevalent health concern hospital-acquired infections. [18] It is one of the most frequent infections, especially among women. [19] There are many factors associated with the occurrence of urinary tract infection. Timely diagnosis and treatment of these predisposing factors is essential for preventing recurrences. [20] Treatment of UTI is usually initiated empirically with antibiotics without performing culture and sensitivity test, has led to improper usage of antibiotics. [21] Identification and practice rational prescription of antibiotics to reduce the emergence of resistant bacterial strains. [22] Culture sensitivity tests were also important in case of UTI to go for a specific antibiotic treatment rather than the empirical treatment for the patients. [23,24]

Out of 120 significant isolates, gram-negative organisms accounted for 81%. Fungal isolates were seen only in 2% of cases. *E. coli* was the most predominant uropathies isolated followed by *Klebsiella spp*, *Enterococcus fecalis* and *Acinetobacter spp*. Other uropathies included MRSA bacteria, *Staphylococcus aureus*, *Candida spp*, *Citrobacter freundii*, *Pseudomonas spp* and *Streptococcus spp*. *E. coli*, showed considerable resistance to ampicillin, ceftriaxone, Co-Trimoxazole and fluoroquinolones. Low level of resistance was observed with Nitrofurantoin, piperacillin+ tazobactam and Ticarcillin. *Klebsiella spp*, the second most common uropathogen showed high level of resistance with ampicillin, cephalexin, cefuroxime, ceftriaxone and Nitrofurantoin. *E. coli* is the commonest uropathies causing both complicated and uncomplicated UTI. [25] In this study, the most predominant uropathies isolated was *E. coli* and this

finding was similar to results of studies by Basnyat KB et al [26] and Khameneh ZR. [27]

E. coli, which was the predominant uropathies isolated showed considerable resistance to ampicillin. A resistance rate comparable to our study was observed in a study conducted by Mandal et al. in South India. [28] In contrast to our observations, in a study conducted by Das RN [29] in West Nepal, *E. coli* isolates showed high susceptibility to ampicillin and cotrimoxazole. High level of resistance was observed with ampicillin in similar with In a retrospective study conducted by Bahadin et al [30], *klebsiella* was found to be the second most prevalent isolate and a hundred percent resistance was observed with ampicillin. However, in contrary to our observations, their study noted higher sensitivity to gentamicin. Another study by Bhargavi PS et al [31] conducted in Southeast part of India showed a higher resistance to gentamicin (83.3%) and cotrimoxazole (82.4%) compared to our study results, which were 54.5% and 45.5% respectively. Our study showed resistance to Nitrofurantoin (6.4 %), piperacillin+ tazobactam (7.8%), and Ticarcillin (4.5%). *Klebsiella spp*.

Conclusion

Most of the identified bacteria were resistant to several of the popular antibiotics used in clinical settings. Consequently, it is vital to prescribe antibiotics rationally both before and after culture reports. Pattern needs to be studied further and the outcome should be communicated to the prescribing physicians in that area.

References

1. Gales AC, Sader HS, Jones RN; SENTRY Participants Group (Latin America). Urinary tract infection trends in Latin American hospitals: report from the SENTRY antimicrobial surveillance program (1997-2000). *Diagn Microbiol Infect Dis*. 2002 Nov; 44(3):289-99.
2. Karlowsky JA, Kelly LJ, Thornsberry C, Jones ME, Sahm DF. Trends in antimicrobial resistance among urinary tract infection isolates of *Escherichia coli* from female outpatients in the United States. *Antimicrob Agents Chemother*. 2002 Aug;46(8):2540-5.

3. De Backer D, Christiaens T, Heytens S, De Sutter A, Stobberingh EE, Verschraegen G. Evolution of bacterial susceptibility pattern of *Escherichia coli* in uncomplicated urinary tract infections in a country with high antibiotic consumption: a comparison of two surveys with a 10 year interval. *J Antimicrob Chemother.* 2008 Aug;62(2):364-8.
4. Rock W, Colodner R, Chazan B, Elias M, Raz R. Ten years surveillance of antimicrobial susceptibility of community-acquired *Escherichia coli* and other uropathogens in northern Israel (1995-2005). *Isr Med Assoc J.* 20 07 Nov;9(11):803-5.
5. Vasquez Y, Hand WL. Antibiotic susceptibility patterns of community-acquired urinary tract infection isolates from female patients on the US (Texas)-Mexico border. *Journal of Applied Research.* 2004 May 1;4(2).
6. Griebing TL. *Urologic Diseases In America Us Dhhs, Phs, Nih, Niddk.*
7. Dimitrov TS, Udo EE, Emara M, Awni F, Passadilla R. Etiology and antibiotic susceptibility patterns of community-acquired urinary tract infections in a Kuwait hospital. *Med Princ Pract.* 2004 Nov-Dec;13(6):334-9.
8. Habte TM, Dube S, Ismail N, Hoosen AA. Hospital and community isolates of uropathogens at a tertiary hospital in South Africa. *S Afr Med J.* 2009 Aug;99(8):584-7.
9. Akram M, Shahid M, Khan AU. Etiology and antibiotic resistance patterns of community-acquired urinary tract infections in J N M C Hospital Aligarh, India. *Ann Clin Microbiol Antimicrob.* 2007 Mar 23;6:4.
10. Kothari A, Sagar V. Antibiotic resistance in pathogens causing community-acquired urinary tract infections in India: a multicenter study. *J Infect Dev Ctries.* 2008 Oct 1;2(5):3 54-8.
11. Biswas D, Gupta P, Prasad R, Singh V, Arya M, Kumar A. Choice of antibiotic for empirical therapy of acute cystitis in a setting of high antimicrobial resistance. *Indian J Med Sci.* 2006 Feb;60(2):53-8.
12. Naveen R, Mathai E. Some virulence characteristics of uropathogenic *Escherichia coli* in different patient groups. *Indian J Med Res.* 2005 Aug;122(2):143-7.
13. Wilkie ME, Almond MK, Marsh FP. Diagnosis and management of urinary tract infection in adults. *BMJ.* 1992 Nov 7;305(68 62):1137-41.
14. Bajaj JK, Karyakarte RP, Kulkarni JD, Deshmukh AB. Changing aetiology of urinary tract infections and emergence of drug resistance as a major problem. *J Commun Dis.* 19 99 Sep;31(3):181-4.
15. Magalit SL, Gler MT, Tupasi TE. Increasing antimicrobial resistance patterns of community and nosocomial uropathogens in Makati Medical Center. *Phil J Microbiol Infect Dis.* 2004;33(4):143-8.
16. Farrell DJ, Morrissey I, De Rubeis D, Robbins M, Felmingham D. A UK multicentre study of the antimicrobial susceptibility of bacterial pathogens causing urinary tract infection. *J Infect.* 2003 Feb;46(2):94-100.
17. Mathai D, Jones RN, Pfaller MA; SENTRY Participant Group North America. Epidemiology and frequency of resistance among pathogens causing urinary tract infections in 1,510 hospitalized patients: a report from the SENTRY Antimicrobial Surveillance Program (North America). *Diagn Microbiol Infect Dis.* 2001 Jul;40(3):129-36.
18. Ronald AR, Pattullo AL. The natural history of urinary infection in adults. *Med Clin North Am* 1991; 75:299-312
19. Ahmed SS, Shariq A, Alsalloom AA, Babikir IH, Alhomoud BN. Uropathogens and their antimicrobial resistance patterns: Relationship with urinary tract infections. *Int J Health Sci.* 2019;13(2):48-55.
20. Foxman B, Gillespie B, Koopman J, Zhang L, Palin K, Tallman P, Marsh JV, Spear S, Sobel JD, Marty MJ, Marrs CF. Risk factors for second urinary tract infection among college women. *Am J Epidemiol.* 2000 Jun 15; 151(12):1194-205.
21. Spellberg B, Bartlett JG, Gilbert DN. The future of antibiotics and resistance. *N Engl J Med.* 2013;368:299-302.
22. Manges AR, Natarajan P, Solberg OD, Dietrich PS, Riley LW: The changing prevalence of drug-resistant *Escherichia coli* clonal groups in a community: evidence for community outbreaks of urinary tract infections. *Epidemiol Infect.* 2006, 134 (2): 425-31. 10.1017/S0950268805005005
23. Panayappan L, Babu AS, Davis D, Joseph N, Joshy N, Krishnakumar K. Urinary tract infection: Prescribing pattern of antibiotics at a tertiary care hospital. *Asian J Pharm Clin Res.* 2017;10(5):255-7.
24. Khatun SA, Shaha S. Prescribing Pattern of Antimicrobials in Urinary Tract Infection at Outpatient Department in a Tertiary Care Hospital in Dhaka. *J Enam Med Coll.* 2020;10 (1):23-6.
25. Beyene G, Tsegaye W. Bacterial uropathogens in urinary tract infection and antibiotic susceptibility pattern in jimma university specialized hospital, southwest ethiopia. *Ethiop J Health Sci.* 2011 Jul;21(2):141-6.
26. Khatri B, Basnyat S, Karki A, Poudel A, Shrestha B. Etiology and antimicrobial susceptibility pattern of bacterial pathogens from urinary tract infection. *Nepal Med Coll J.* 2012 Jun;14(2):129- 32

27. Khameneh ZR, Afshar AT. Antimicrobial susceptibility pattern of urinary tract pathogens. Saudi J Kidney Dis Transpl. 2009 Mar;20(2):251-3.
28. Mandal J, Acharya NS, Buddhapriya D, Parija SC. Antibiotic resistance pattern among common bacterial uropathogens with a special reference to ciprofloxacin resistant Escherichia coli. Indian J Med Res. 2012 Nov;136(5):842-9.
29. Das RN, Chandrashekhar TS, Joshi HS, Gurung M, Shrestha N, Shivananda PG. Frequency and susceptibility profile of pathogens causing urinary tract infections at a tertiary care hospital in western Nepal. Singapore Med J. 2006 Apr;47(4):281-5.
30. Bahadin J, Teo SS, Mathew S. Aetiology of community-acquired urinary tract infection and antimicrobial susceptibility patterns of uropathogens isolated. Singapore Med J. 2011 Jun;52(6):415-20.
31. Bhargavi PS, Gopala Rao TV, Mukkanti K, Dinesh Kumar B, Krishna TP. Increasing emergence of antibacterial resistance mainly in uropathogens:southeast part of India. Int J Microbiol Res. 2010;2(1):1-6.