

To Quantify the Amount of Protein Intake and to Compare Growth Parameters at Hospital Discharge in ELBW Infants: A Retrospective Study

Prashant Kumar¹, Rishika Verma², Rajaram Pd Singh³

¹Junior Resident, Upgraded Department of Pediatrics, Patna Medical College and Hospital, Patna, Bihar, India

²Junior Resident, Upgraded Department of Pediatrics, Patna Medical College and Hospital, Patna, Bihar, India

³Associate Professor (Unit Head) Upgraded Department of Pediatrics, Patna Medical College and Hospital, Patna, Bihar, India

Received: 13-01-2024 / Revised: 10-02-2024 / Accepted: 27-03-2024

Corresponding Author: Dr. Rishika Verma

Conflict of interest: Nil

Abstract

Aim: The aim of the present study was to quantify the amount of protein intake and to compare growth parameters at hospital discharge (as measured by discharge weight and head circumference percentiles) when using two different methodologies (BW versus current daily weight until BW is regained) for calculating fluid and protein intake in the first two weeks after birth in ELBW infants.

Material & Methods: The present study was conducted in the Upgraded Department Of Paediatrics, Patna Medical College and Hospital, Patna, Bihar, India from January 2014 to December 2014 and all ELBW admissions to the NICU were included except those regaining BW in ≤ 3 days, those who were transferred to another facility, or those who died before reaching discharge.

Results: We studied 150 infants with 100 and 50 infants in P1 and P2 periods, respectively. Mean gestational age was lower during Phase 1 (24.86 ± 1.34 vs. 25.45 ± 1.85 weeks). P value = 0.01). However, the birth weights were not different between the two periods. The incidence of SGA at birth was higher during Phase 2. Compared to P1, babies in P2 received 1 g/kg/day more protein till BW was regained. This average daily protein deficit would have resulted in a cumulative protein deficit of 5.1 g/kg if the infant's current weight was used for protein calculations for the duration of 1 week (P2 phase). There were no significant differences in protein and fluid intakes. However, there was a higher caloric intake in P2 compared to P1.

Conclusion: Basing nutrition calculations in ELBW on birth weight rather than current daily weight until the birth weight is regained resulted in significantly greater protein delivery, a significant decrease in the incidence of failure to thrive and smaller head circumference percentiles at discharge in ELBW infants.

Keywords: Nutrition, Prematurity, Breastfeeding, Fat mass, Fortification

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

Introduction

During the 2nd and early 3rd trimester, foetus receive about 4 ± 0.5 gr/kg/day of protein. [1] A study from NICHD showed that >90% ELBW (extremely low birth weight) infants were growth restricted by 36 weeks of post-menstrual gestational age (PMA). [2] Postnatal growth failure, which is associated with impaired neurocognitive and renal development, [3,4,5,6] was observed in nearly 100% of very low birth weight infants at the end of the 1990s and still occurs in more than half of these infants. [7] Growth impairment during early infancy can have permanent detrimental effects. [8] Impaired foetal and postnatal growth has been associated with neurodevelopmental delay, ischemic heart disease, impaired glucose tolerance, type-II

diabetes mellitus, hypertension, and metabolic syndrome. [9,10] Generally, growth failure is considered when a baby's weight is below the 10th percentile (≤ -1.28 Z-score), but with different reference charts and variable postnatal periods having been used. [11] Although length growth deficit is also very common, the vast majority of children gradually catch up between the ages of 2 and 8 years and are within normal height ranges as adults. [12,13] A deficit in fat-free mass (FFM) at discharge has been associated with suboptimal neurological outcomes, and the proportion of fat mass (FM) is known to be higher in premature infants compared to foetuses of the same gestational age (GA). [14,15] Small for gestational age (SGA)

infants had poorer weight gain than appropriate for gestational age (AGA) infants [16], and SGA infants remained at a lower body weight until the age of 3–6 years. [17]

Factors such as weight, gestational age, length of hospital stay, presence of respiratory distress syndrome (RDS), bronchopulmonary dysplasia (BPD), and sepsis have been associated with post-discharge growth and development in preterm infants. [18] According to a National Institute of Child Health and Human Development study on VLBW infants, nutritional intake deficits persist to some degree during the hospital stay, and infants with BPD, necrotizing enterocolitis (NEC), or late-onset sepsis demonstrate slower growth. Poor nutritional intake is common in VLBW in the early postnatal months. [19] This prompted the promotion of early aggressive nutrition in this vulnerable population [20]. Protein delivery increases steadily as the foetus grows and gains weight in utero.

A recent studies have shown a direct association between first-week nutritional intakes, especially higher protein intake, with improved neurodevelopment at 18-24 months. [21,22] Based on information from such studies, the use of 3-4gm/kg/day of protein in parenteral nutrition for ELBW infants within the first few days of life has become the standard goal. In the premature infant, growth and weight gain do not always occur in the first few days or weeks after birth. An ELBW infant may lose 10-20% of its birth weight (BW) and may take 2 to 3 weeks to regain its BW. Since fluid and nutrition need and delivery in ELBW infants are calculated based on their body weights, there could be a measurable difference in fluid, nutrition, and protein intake calculations based on birth weight or current weight of the infant, especially in the first two weeks of life. The issue is further complicated by the fact that fluid overload in the early newborn period is associated with negative cardiorespiratory outcomes, and most centres pay close attention to the daily fluctuations in weight when managing fluid balance in these infants. [23]

The aims of the study were to quantify the amount of protein intake and compare growth parameters at hospital discharge (as measured by discharge weight and head circumference percentiles) when using two different methodologies (birth weight versus current daily weight until birth weight is regained) for calculating fluid and protein intake in the first two weeks after birth in ELBW infant.

Materials & Methods

The present study was conducted in the Upgraded Department Of Paediatrics, Patna Medical College and Hospital, Patna, Bihar, India from January 2014 to December 2014 and all ELBW admissions to the NICU were included except those regaining BW in

≤ 3 days, those who were transferred to another facility, or those who died before reaching discharge.

Exclusion criteria

- Those regaining BW in ≤ 3 days, those who were transferred to another facility, or
- Those who died before reaching discharge.

Methodology

Birth weight, current daily weight, caloric intake (Kcal/kg/day), total fluid intake (mL/kg/day), and parenteral protein intake (gm/kg/day) were recorded daily until BW was regained and until discharge. At this centre, in P1 CW was exclusively used for calculating fluid, caloric, and protein administration till BW was regained. In P2, BW was exclusively used for all calculations.

Comorbid conditions that may influence an infant's nutritional needs and growth outcomes were studied. CLD (chronic lung disease) was defined as the requirement for oxygen at 36 weeks of corrected gestational age. [24] PDA (patent ductus arteriosus) was defined as the presence of patent ductus arteriosus on an echocardiogram. NEC (necrotizing enterocolitis) was defined as the presence of pneumatosis intestinalis or air in the portal vein on an X-ray or surgical diagnosis at the time of laparotomy. [25] IVH (intraventricular haemorrhage) was defined as the presence of any haemorrhage in the lateral ventricles of the brain on an ultrasound. [26] Gender-specific Fenton growth charts (2013) were used for growth assessment. [27] The length was not analysed due to intrinsic variability in measurement secondary to lack of standard-length board. This study received institutional review board exemption as part of the quality improvement. Feeding and nutritional practices during the two phases were as follows.

In P1, infants were gavage-fed ≤15ml/kg/day for seven days. Soon after birth, protein-based early parenteral nutrition was initiated with infants being given 2g/kg/day within 24 hours of birth. Fluids are initially started at 80-100 ml/kg/day and advanced as needed based on serum sodium levels. Enteral nutrition is calculated based on formula nutrient composition and average breast milk composition. Protein intake was calculated to provide 3-4 g/kg/day in the first week of life based on the current weight. Subsequently, daily administration of a protein of 3-4g/kg/day, calories of 120-130 kcal/kg/day, and fluids of 135-155 ml/kg/day was targeted based on CW, unless an infant's condition warranted a change. Human milk fortifier was used to fortify the breast milk to 24 cal /oz. Average targeted daily weight gain was 15-20 grams/kg/d. No probiotics were administered.

P2 infants were gavage-fed ≤ 15 ml/kg/day for five days. Soon after birth, protein-based early parenteral nutrition was initiated with infants being given 2g/kg/day within 24 hours of birth. Protein intake was calculated to provide 3-4g/kg/day after 24 hours of age. Subsequently, daily administration of a protein of 3-4g/kg/day, calories of 120- 130 kcal/kg/day, and fluids of 135-155 ml/kg/day was targeted based on BW, unless infant's condition warranted a change. Human milk fortifier was used to fortify the breast milk to 24 cal/oz. Average targeted daily weight gain was 15-20 grams/kg/d. Infants received a daily dose of probiotics (UDO's choice infant's probiotic, Flora Inc., Lynden,

Washington, USA). P1 and P2 periods were compared and analysed for differences in demographics, nutritional intake, comorbid conditions, and growth outcomes.

Statistical Analysis

Statistical analysis was completed using SPSS v.18. Descriptive statistics were summarized using frequencies and means. Wilcoxon Signed-rank test and Chi-square analyses were used to determine associations between variables.

Results

Table 1: Patient characteristics

	P1 (N=100) Count	%	P2 (N=50) Count	%	P value
Female	45	45	26	52	0.36
Gest Age-mean, std dev	24.86	1.34	25.45	1.85	0.01
Birth Wt-mean, std dev	771.89	128.92	746.84	164.76	0.34
Birth Weight <10%	15	15	40	40	<.001
Birth HC <10%	18	18	28	28	0.18
Discharge Weight <10%	76	76	43	43	<.001
Discharge HC <10%	34	34	14	28	0.42

We studied 150 infants with 100 and 50 infants in P1 and P2 periods, respectively. Mean gestational age was lower during Phase 1 (24.86 \pm 1.34 vs. 25.45 \pm 1.85 weeks). P value = 0.01).

Table 2: Daily nutritional intake before regaining birth weight

	Phase 1 (N=100)		Phase 2 (N=50)		P Value
	Mean	SD	Mean	SD	
Protein Received	2.48	0.46	3.47	0.34	0
Calories Received	64.86	12.8	67.03	13.37	0.07
Fluids Received	134.48	14.66	120.8	13.34	0

However, the birth weights were not different between the two periods. The incidence of SGA at birth was higher during Phase 2.

Table 3: Nutritional intake after regaining birth weight to the time of discharge

Parameter	Period	Mean	Std. Dev	P value
Protein g/kg/day	Phase 1	3.26	0.34	0.138
	Phase 2	3.39	0.38	
Calories kg/day	Phase 1	111.19	6.44	0.032
	Phase 2	114.6	6.04	
Fluids cc/kg/day	Phase 1	144.06	10.36	0.119
	Phase 2	138.52	8.2	

Compared to P1, babies in P2 received 1 g/kg/day more protein till BW was regained. This average daily protein deficit would have resulted in a cumulative protein deficit of 5.1 g/kg if the infant's current weight was used for protein calculations for the duration of 1 week (P2 phase).

Table 4: Morbidity before hospital discharge between the two periods of study

	Phase 1		Phase 2		p-value
	Count	%	Count	%	
CLD	36	36	48	48	0.15
PDA	63	63	45	45	0.07
IVH	18	18	26	26	0.25
NEC	10	10	5	5	0.314

There were no significant differences in protein and fluid intakes. However, there was a higher caloric intake in P2 compared to P1.

Discussion

Postnatal growth failure, which is associated with impaired neurocognitive and renal development [28-31], was observed in nearly 100% of very low birth weight infants at the end of the 1990s and still occurs in more than half of these infants. [32] The goal of nutrition for premature infants is to duplicate the in-utero growth rate and body composition of a foetus of the same gestational age. [33] However, achieving this weight gain goal is difficult and, consequently, preterm infants are frequently significantly underweight at the time of hospital discharge. [34] Recently, the Vermont Oxford Network, which evaluates the postnatal growth of preterm infants, defined postnatal growth failure (PGF) as a discharge weight that is lower than the 10th percentile for postmenstrual age. Among infants registered in this Network, the prevalence rate of PGF is 50.3%. [35] Kelleher reported that birth weight was significantly lower in low-birth-weight children with failure to thrive than in those without failure to thrive. [36] As well, small for gestational age (SGA) infants had poorer weight gain than appropriate for gestational age (AGA) infants [37] and SGA infants remained at a lower body weight until the age of 3–6 years. [38]

We studied 150 infants with 100 and 50 infants in P1 and P2 periods, respectively. Mean gestational age was lower during Phase 1 (24.86 ± 1.34 vs. 25.45 ± 1.85 weeks). P value = 0.01). However, the birth weights were not different between the two periods. The incidence of SGA at birth was higher during Phase 2. Compared to P1, babies in P2 received 1 g/kg/day more protein till BW was regained. This average daily protein deficit would have resulted in a cumulative protein deficit of 5.1 g/kg if the infant's current weight was used for protein calculations for the duration of 1 week (P2 phase). There were no significant differences in protein and fluid intakes. However, there was a higher caloric intake in P2 compared to P1. When using CW for nutritional calculations, we found that there was an average of a 2 gm/kg deficit of protein in the first week of life with some infants losing up to a 6.9 gm/kg during P1. It is possible that this protein deficit may not have been adequately compensated for, which led to significantly poorer weight and head circumference growth during P1 compared to P2. This may have an impact on future neurodevelopmental outcomes in these infants. Our findings may be corroborated by a recent study that showed that, for each gram/kg increase in protein delivery in the first week, there was an increase of 8.2 points in MDI (mental development index). [39] Another study has shown that higher protein and caloric intake reduced

postnatal head growth failure in preterm infants of <29 weeks' gestation. [40] Thus, using BW for protein calculations (till BW is regained) may be a simple, but effective, strategy to avoid significant protein deficits and consequent postnatal growth restriction. It is also important to consider when the additional protein is provided, as that may also be critical to avoid a fall in growth percentile. A study involving 560 children showed that growth during early infancy (0-4 months) predicted IQ at nine years of age. [41]

Another aspect of neonatal physiology that impacts this issue is that a fetus in utero receives increasing amounts of protein every day, which accounts for its continuously increasing growth with advancing gestation. In contrast, soon after birth, this increasing protein transfer is not accounted for in the infant's nutrition, and the infant loses weight and may become catabolic. It may take up to a week or longer to regain BW and reverse the catabolic changes. During this period, even in the best circumstances, the infant would receive significantly less protein and other nutrients than it would have in utero. Therefore, there is a need for an additional allowance of nutritional requirements for catch-up growth. [42]

Conclusion

Basing nutrition calculations in ELBW on birth weight rather than current daily weight until the birth weight is regained resulted in significantly higher protein delivery, a significant decrease in the incidence of failure to thrive, and smaller head circumference percentiles at discharge in ELBW infants. The findings from this retrospective study with historical controls need to be corroborated with a larger multicentre population of ELBW infants to confirm the above conclusion.

References

1. Hay WW, MD J, Lucas A, Heird WC, Ziegler E, Levin E, Grave GD, Catz CS, Yaffe SJ. Workshop summary: nutrition of the extremely low birth weight infant. *Pediatrics*. 1999 Dec 1;104(6):1360-8.
2. Ehrenkranz RA, Younes N, Lemons JA, Fanaroff AA, Donovan EF, Wright LL, Katsikiotis V, Tyson JE, Oh W, Shankaran S, Bauer CR. Longitudinal growth of hospitalized very low birth weight infants. *Pediatrics*. 1999 Aug 1;104(2):280-9.
3. Raghuram K, Yang J, Church PT, Cieslak Z, Synnes A, Mukerji A, Shah PS, Canadian Neonatal Network, Canadian Neonatal Follow-Up Network Investigators. Head growth trajectory and neurodevelopmental outcomes in preterm neonates. *Pediatrics*. 2017 Jul 1;140(1).

4. Vinall J, Grunau RE, Brant R, Chau V, Poskitt KJ, Synnes AR, Miller SP. Slower postnatal growth is associated with delayed cerebral cortical maturation in preterm newborns. *Science translational medicine*. 2013 Jan 16;5(168):168ra8-.
5. Bacchetta J, Harambat J, Dubourg L, Guy B, Liutkus A, Canterino I, Kassai B, Putet G, Cochat P. Both extrauterine and intrauterine growth restriction impair renal function in children born very preterm. *Kidney international*. 2009 Aug 2;76(4):445-52.
6. Ordóñez-Díaz MD, Pérez-Navero JL, Flores-Rojas K, Olza-Meneses J, Muñoz-Villanueva MC, Aguilera-García CM, Gil-Campos M. Prematurity with extrauterine growth restriction increases the risk of higher levels of glucose, low-grade of inflammation and hypertension in prepubertal children. *Frontiers in Pediatrics*. 2020 Apr 21;8:180.
7. Horbar JD, Ehrenkranz RA, Badger GJ, Edwards EM, Morrow KA, Soll RF, Buzas JS, Bertino E, Gagliardi L, Bellù R. Weight growth velocity and postnatal growth failure in infants 501 to 1500 grams: 2000–2013. *Pediatrics*. 2015 Jul 1;136(1):e84-92.
8. Embleton NE, Pang N, Cooke RJ. Postnatal malnutrition and growth retardation: an inevitable consequence of current recommendations in preterm infants?. *Pediatrics*. 2001 Feb 1;107(2):270-3.
9. Lapillonne A, Griffin IJ. Feeding preterm infants today for later metabolic and cardiovascular outcomes. *The Journal of pediatrics*. 2013 Mar 1;162(3):S7-16.
10. Barker, D. J. Te fetal and infant origins of adult disease. *BMJ* 301, 1111 (1990).
11. Fenton, T. R. et al. Preterm Infant Growth Velocity Calculations: A Systematic Review. *Pediatrics* 139 (2016).
12. Sakurai M, Itabashi K, Sato Y, Hibino S, Mizuno K. Extrauterine growth restriction in preterm infants of gestational age \leq 32 weeks. *Pediatrics International*. 2008 Feb;50(1):70-5.
13. Saigal S, Stoskopf B, Streiner D, Paneth N, Pinelli J, Boyle M. Growth trajectories of extremely low birth weight infants from birth to young adulthood: a longitudinal, population-based study. *Pediatric research*. 2006 Dec;60(6):751-8.
14. Ramel SE, Gray HL, Christiansen E, Boys C, Georgieff MK, Demerath EW. Greater early gains in fat-free mass, but not fat mass, are associated with improved neurodevelopment at 1 year corrected age for prematurity in very low birth weight preterm infants. *The Journal of pediatrics*. 2016 Jun 1;173:108-15.
15. Hamatschek C, Yousuf EI, Möllers LS, So HY, Morrison KM, Fusch C, Rochow N. Fat and fat-free mass of preterm and term infants from birth to six months: a review of current evidence. *Nutrients*. 2020 Jan 21;12(2):288.
16. Pilling, E. L., Elder, C. J. & Gibson, A. T. Growth patterns in the growth-retarded premature infant. *Best Pract Res Clin Endocrinol Metab* 22, 447–462 (2008).
17. Monset-Couchard, M., de Bethmann, O. & Relier, J. P. Long term outcome of small versus appropriate size for gestational age cotwins/triplets. *Arch Dis Child Fetal Neonatal Ed* 89, F310–314 (2004).
18. Lima PA, Carvalho MD, Costa AC, Moreira ME. Variables associated with extra uterine growth restriction in very low birth weight infants. *Jornal de pediatria*. 2014 Jan;90:22-7.
19. Ehrenkranz RA, Younes N, Lemons JA, Fanaroff AA, Donovan EF, Wright LL, Katsikiotis V, Tyson JE, Oh W, Shankaran S, Bauer CR. Longitudinal growth of hospitalized very low birth weight infants. *Pediatrics*. 1999 Aug 1;104(2):280-9.
20. Thureen PJ, Hay Jr WW. Early aggressive nutrition in preterm infants. In *Seminars in Neonatology* 2001 Oct 1 (Vol. 6, No. 5, pp. 403-415). WB Saunders.
21. Yang J, Chang SS, Poon WB. Relationship between amino acid and energy intake and long-term growth and neurodevelopmental outcomes in very low birth weight infants. *Journal of Parenteral and Enteral Nutrition*. 2016 Aug;40(6):820-6.
22. Cormack BE, Bloomfield FH, Dezoete A, Kuschel CA. Does more protein in the first week of life change outcomes for very low birthweight babies?. *Journal of paediatrics and child health*. 2011 Dec;47(12):898-903.
23. Van Marter LJ, Leviton A, Allred EN, Pagano M, Kuban KC. Hydration during the first days of life and the risk of bronchopulmonary dysplasia in low birth weight infants. *The Journal of pediatrics*. 1990 Jun 1;116(6):942-9.
24. A. T. Shennan, M. S. Dunn, A. Ohlsson, K. Lennox, and E. M. Hoskins, "Abnormal pulmonary outcomes in premature infants: Prediction from oxygen requirement in the neonatal period," *Pediatrics*, vol. 82, no. 4, pp. 527–532, 1988.
25. M. J. Bell, J. L. Ternberg, R. D. Feigin et al., "Neonatal necrotizing enterocolitis. Therapeutic decisions based upon clinical staging," *Annals of Surgery*, vol. 187, no. 1, pp. 1–7, 1978.
26. L. A. Papile, J. Burstein, R. Burstein, and H. Kofer, "Incidence and evolution of subependymal and intraventricular hemorrhage: a study of infants with birth weights less than 1,500 gm," *Journal of Pediatrics*, vol. 92, no. 4, pp. 529–534, 1978.
27. T. R. Fenton, R. Nasser, M. Eliasziw, J. H. Kim, D. Bilan, and R. Sauve, "Validating the weight gain of preterm infants between the reference

- growth curve of the fetus and the term infant,” *BMC Pediatrics*, vol. 13, no. 1, 2013.
28. Raghuram K, Yang J, Church PT, Cieslak Z, Synnes A, Mukerji A, Shah PS, Canadian Neonatal Network, Canadian Neonatal Follow-Up Network Investigators. Head growth trajectory and neurodevelopmental outcomes in preterm neonates. *Pediatrics*. 2017 Jul 1;140(1).
 29. Vinall J, Grunau RE, Brant R, Chau V, Poskitt KJ, Synnes AR, Miller SP. Slower postnatal growth is associated with delayed cerebral cortical maturation in preterm newborns. *Science translational medicine*. 2013 Jan 16;5(168):168ra8-.
 30. Bacchetta J, Harambat J, Dubourg L, Guy B, Liutkus A, Canterino I, Kassaï B, Putet G, Cochat P. Both extrauterine and intrauterine growth restriction impair renal function in children born very preterm. *Kidney international*. 2009 Aug 2;76(4):445-52.
 31. Ordóñez-Díaz MD, Pérez-Navero JL, Flores-Rojas K, Olza-Meneses J, Muñoz-Villanueva MC, Aguilera-García CM, Gil-Campos M. Prematurity with extrauterine growth restriction increases the risk of higher levels of glucose, low-grade of inflammation and hypertension in prepubertal children. *Frontiers in Pediatrics*. 2020 Apr 21;8:180.
 32. Horbar JD, Ehrenkranz RA, Badger GJ, Edwards EM, Morrow KA, Soll RF, Buzas JS, Bertino E, Gagliardi L, Bellù R. Weight growth velocity and postnatal growth failure in infants 501 to 1500 grams: 2000–2013. *Pediatrics*. 2015 Jul 1;136(1):e84-92.
 33. American Academy of Pediatrics Committee on Nutrition. Nutritional needs of low-birth-weight infants. *Pediatrics*. 1985;75(5):976-86.
 34. Embleton NE, Pang N, Cooke RJ. Postnatal malnutrition and growth retardation: an inevitable consequence of current recommendations in preterm infants?. *Pediatrics*. 2001 Feb 1;107(2):270-3.
 35. Horbar JD, Ehrenkranz RA, Badger GJ, Edwards EM, Morrow KA, Soll RF, Buzas JS, Bertino E, Gagliardi L, Bellù R. Weight growth velocity and postnatal growth failure in infants 501 to 1500 grams: 2000–2013. *Pediatrics*. 2015 Jul 1;136(1):e84-92.
 36. Kelleher KJ, Casey PH, Bradley RH, Pope SK, Whiteside L, Barrett KW, Swanson ME, Kirby RS. Risk factors and outcomes for failure to thrive in low birth weight preterm infants. *Pediatrics*. 1993 May 1;91(5):941-8.
 37. Pilling EL, Elder CJ, Gibson AT. Growth patterns in the growth-retarded premature infant. *Best practice & research Clinical endocrinology & metabolism*. 2008 Jun 1;22(3):447-62.
 38. Monset-Couchard M, De Bethmann O, Relier JP. Long term outcome of small versus appropriate size for gestational age co-twins/triplets. *Archives of Disease in Childhood-Fetal and Neonatal Edition*. 2004 Jul 1;89(4):F310-4.
 39. Stephens BE, Walden RV, Gargus RA, Tucker R, McKinley L, Mance M, Nye J, Vohr BR. First-week protein and energy intakes are associated with 18-month developmental outcomes in extremely low birth weight infants. *Pediatrics*. 2009 May 1;123(5):1337-43.
 40. Morgan C, McGowan P, Herwitker S, Hart AE, Turner MA. Postnatal head growth in preterm infants: a randomized controlled parenteral nutrition study. *Pediatrics*. 2014 Jan 1;133(1):e120-8.
 41. Pongcharoen T, Ramakrishnan U, DiGirolamo AM, Winichagoon P, Flores R, Singkhornard J, Martorell R. Influence of prenatal and postnatal growth on intellectual functioning in school-aged children. *Archives of pediatrics & adolescent medicine*. 2012 May 1;166(5):411-6.
 42. Adamkin DH. Nutrition management of the very low-birthweight infant: ii. optimizing enteral nutrition and postdischarge nutrition. *NeoReviews*. 2006 Dec 1;7(12):e608-14.