

## Assessment of Nutritional Awareness among Mothers and Iron Deficiency Anemia in Infants

Amit Kumar<sup>1</sup>, Vibhas Ranjan<sup>2</sup>

<sup>1</sup>Assistant Professor, Department of Pediatrics, DMCH, Darbhanga, Bihar, India

<sup>2</sup>Assistant Professor Department of Pediatrics, DMCH, Darbhanga, Bihar, India

Received: 10-01-2024 / Revised: 13-02-2024 / Accepted: 20-03-2024

Corresponding Author: Dr. Vibhas Ranjan

Conflict of interest: Nil

### Abstract

**Aim:** The aim of the present study was to analyze the risk factors and vital role of mothers in prevention of nutritional anemia in the early childhood.

**Methods:** This study was carried out in the Department of Pediatrics, DMCH, Darbhanga, Bihar, India and children aged 6 months–2 years admitted at the hospital during the study period were included. Any child with chronic disorders such as hemoglobinopathies, hemolytic anemia, and treated for anemia and those on any immunosuppressants such as steroids and biologics were excluded as an indirect measure to eliminate anemia of chronic disease. The study protocol was approved by the Hospital Ethics and Scientific Committee. An informed written consent was obtained from the mothers of these children. A total of 200 participants were included.

**Results:** Majority of the mothers were aware that neural development of child could be affected by anemia. Maternal knowledge regarding cow's milk was poor and had a faulty belief that it increases iron absorption, whereas 16% were unaware of the relationship between cow's milk and anemia. Many mothers believed anemia to occur only in children on vegetarian diet. Around 75% mothers recognized green leaves and 70% women recognized jaggery to be enriched with iron. Although 78% identified vitamin C containing fruits, many were not aware of its vital role in increasing the iron absorption. Mothers of children with more than one sibling had a relatively greater incidence of "poor" knowledge unlike mothers of children with single or no sibling. This implies that those with "poor" knowledge were also unaware of the importance of birth spacing.

**Conclusion:** Recognizing the causative factors plays an essential role in preventing iron deficiency anemia (IDA). Exclusive breastfeeding must be promoted, along with avoiding excessive cow's milk intake. Intervening at the right period with age-appropriate foods becomes a necessity. Maternal illiteracy has a positive correlation with anemia in infants. Improving maternal awareness by the physicians will pave the road toward a nation free from anemia.

**Keywords:** Maternal awareness, Iron deficiency anemia, Child nutrition, Weaning

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

### Introduction

Increasingly complex nutrition is required by the body during pregnancy. Therefore, difficulties in fulfilling these nutrition requirements are common. One of the most common nutritional problems in pregnant women is iron deficiency anemia, the most widespread micronutrient problem and the most difficult to overcome worldwide. [1] The incidence of anemia is still high, affecting a fairly large proportion of people worldwide. According to World Health Organization (WHO) data, approximately 29.9% of women globally in the 15–49 age group have anemia. This is equivalent to more than half a billion people. The prevalence is 29.6% in non-pregnant women and 36.5% in pregnant women. [2] The incidence of anemia in pregnant women is very high in lower-middle-income countries (LMICs), especially in Southeast

Asia and Africa. Based on the World Bank income category, the prevalence of anemia in pregnant women in LMICs is 45%. Meanwhile, in high-income countries, the prevalence of anemia in pregnant women is only around 17%. [3]

Anemia in pregnancy is potentially harmful to the mother and child. Therefore, it requires serious attention from all parties involved in health services. Anemia in pregnant women can cause complications such as bleeding, increased risk of birthweight, abortion, and increased risk of mortality for both mother and baby. [4] Low hemoglobin levels in pregnant women can impact fetal growth and development. Hence, pregnant women with a history of anemia are more likely to have infants with low birthweight. [5] Considering the impact of iron

deficiency anemia on pregnant women and the potential future health conditions, preventing anemia during pregnancy is crucial. Poor awareness and prevention restrict efforts to reduce the prevalence of iron deficiency anemia. [6] According to Rogers' theory (1962), commonly known as the AIETA theory, the body undergoes a series of sequential stages while adopting new behaviors: awareness, interest, evaluation, trial, and adoption. If the behavior adoption through this process is built on awareness, the behavior will persist.

Maternal nutritional deficiency during pregnancy affects the developmental process of the fetus, which subsequently influences the birth weight of the newborn. [7] The fetus is highly dependent on maternal nutritional intake since malnutrition during pregnancy leads to different adverse birth outcomes like LBW. During pregnancy, insufficient storage or inadequate intake of essential nutrients can cause harmful effects on both the mothers and newborn babies. [8-10] The strategies of anemia prevention are iron supplementation, regular de-worming, control, and prevention of parasitic infections in pregnancy, such as by using insecticide-treated bed net consistently, intake of iron-rich foods, nutritional counseling such as not taking coffee, tea, or milk with meals, accessing clean and adequate water, and by treating the underlying causes and complications. [11]

Hence the aim of the study was to analyze the risk factors and vital role of mothers in prevention of nutritional anemia in the early childhood.

### Material & Methods

This study was carried out in the Department of Pediatrics, DMCH, Darbhanga, Bihar, India for one year and children aged 6 months–2 years admitted at the hospital during the study period were included. Any child with chronic disorders such as hemoglobinopathies, hemolytic anemia, and treated for anemia and those on any immunosuppressants such as steroids and biologics were excluded as an indirect measure to eliminate anemia of chronic disease. An informed written consent was obtained from the mothers of these children. A total of 200 participants were included.

Later, a pre-designed pro forma was used to record the relevant information. The two pages pro forma would include six sets of questionnaire. Only the mother was allowed to answer the questionnaire. The first part comprises general details including demography, personal data, socioeconomic status,

and family background. Information pertaining to significant history of both mother and child was included in the second part. A detailed nutritional history of the child was assessed in the third part of the pro forma. Clinical presentation and investigation details were recorded in the fourth and the fifth, respectively. The final part of the questionnaire was added to analyze the awareness of the mothers on anemia and their extent of knowledge on the risk factors, clinical features, and the importance of treating IDA. Length and weight were measured uniformly using infantometer and digital weighing machine to the nearest 0.1 cm and 0.001 kg, respectively. Nourishment was graded based on the World Health Organization (WHO) guidelines and chronically malnourished children were excluded from the study. [12] Children were classified based on Modified Kuppuswamy scale of socioeconomic strata. [13] Two generations of family living in the same household were considered 'nuclear' family. Anything beyond was considered 'joint' family. Complete blood count including red blood cell indices was calculated using Coulter LH 780 Hematology analyzer. Mentzer index was calculated along to differentiate between IDA and Thalassemia. Anemia was graded as per the WHO guidelines. [14] Questionnaire was prepared in both English and vernacular language for better understanding. On designing the questionnaire, it was validated by an expert panel consisting of a Pediatrician, healthcare workers, and academic professionals including experts in vernacular language for easy comprehensibility.

### Statistical Analysis

Anemia was the primary outcome variable and knowledge on nutrition/anemia was considered as the secondary variable. Descriptive analysis was carried out by mean and standard deviation for quantitative variables, frequency, and proportion for categorical variables. The association between variables of anemia and quantitative outcome was assessed by comparing the mean values. Independent sample t-test was used to assess the statistical significance. The association between explanatory variables and categorical outcomes was assessed by cross tabulation and comparison of percentages. Chi-square test was used to test the significance of statistics.  $p < 0.05$  was considered statistically significant. IBM SPSS version 22 was used for statistical analysis.<sup>15</sup>

### Results

**Table 1: Tabular column showing the percentage of answers by the mothers**

Correct answer	No. of correct answers (%)	No. of incorrect answers (%)
Anemia is a deficiency of hemoglobin	156 (78)	44 (22)
Iron is important for carrying oxygen in the blood to various organs	52 (26)	148 (74)
Infants at weaning age are at high risk for anemia	44 (22)	156 (78)
Infants at weaning age are at high risk for anemia	50 (25)	150 (75)
Anemia affects both vegetarians and non-vegetarians equally	32 (16)	168 (84)
Large quantities of cow’s milk decreases iron absorption	32 (16)	168 (84)
Complementary feeds should be started at 6 months	36 (18)	164 (82)
Jaggery contains more iron	140 (70)	60 (30)
Green leaves are rich in iron	150 (75)	50 (25)
Meat is rich in iron	40 (20)	160 (80)
Orange is rich in vitamin C	160 (80)	40 (20)
If mother takes iron supplements, breastfed infant doesn’t get more iron	160 (80)	40 (20)
Vitamin C is necessary for absorption of iron	156 (78)	44 (22)
Anemia is often asymptomatic	156 (78)	44 (22)
Anemia can cause neurodevelopmental delay in infants	156 (78)	44 (22)
Government provides free iron supplements	24 (12)	176 (88)
Anemia can be prevented	150 (75)	50 (25)
Iron causes constipation but should not be avoided	10 (5)	190 (95)
Blood transfusion is not necessary for all children with anemia	20 (10)	180 (90)

Majority of the mothers were aware that neural development of child could be affected by anemia. Maternal knowledge regarding cow’s milk was poor and had a faulty belief that it increases iron absorption, whereas 16% were unaware of the relationship between cow’s milk and anemia. Many

mothers believed anemia to occur only in children on vegetarian diet. Around 75% mothers recognized green leaves and 70% women reconized jaggery to be enriched with iron. Although 78% identified vitamin C containing fruits, many were not aware of its vital role in increasing the iron absorption.

**Table 2: Comparison of number of siblings across knowledge about nutrition and anemia**

Number of siblings	Maternal awareness on anemia and nutrition			P Value
	Poor knowledge	Fair knowledge	Good knowledge	
Nil (20)	4	14	2	0.017
One (70)	11	49	10	
More than one (10)	4	5	1	

Mothers of children with more than one sibling had a relatively greater incidence of “poor” knowledge unlike mothers of children with single or no sibling. This implies that those with “poor” knowledge were also unaware of the importance of birth spacing.

**Discussion**

Anemia, a major global public health burden, affects 25% of the population worldwide. [16] Although anemia is multifactorial, iron deficiency anemia (IDA) is the major cause (42%). [17] Affected children have nonspecific symptoms; therefore, a greater proportion of them remains undiagnosed unless a health problem ensue. [16,18] Inappropriate weaning in the infancy and faulty feeding practices is some of the early contributors to anemia. Apart from these amenable risk factors, low birth weight and premature infants are at a higher risk of developing anemia. [19] All these factors, leading to IDA during infantile and early childhood, has a negative impact on the motor and neuro-cognitive function. [20] Iron is an essential nutrient for a

child’s proper development at every growth stage. [21] Iron is crucial for the production of new red blood and muscle cells, DNA replication, and the development of the brain, nervous and immune systems. [22,23] Iron deficiency in infants can result in poor memory and attention, a higher risk of attention-deficit hyperactivity disorder, visual and auditory system impairment and social and emotional behaviours. [24]

Majority of the mothers were aware that neural development of child could be affected by anemia. Maternal knowledge regarding cow’s milk was poor and had a faulty belief that it increases iron absorption, whereas 16% were unaware of the relationship between cow’s milk and anemia. Many mothers believed anemia to occur only in children on vegetarian diet. Around 75% mothers recognized green leaves and 70% women reconized jaggery to be enriched with iron. Although 78% identified vitamin C containing fruits, many were not aware of its vital role in increasing the iron absorption.

Mothers of children with more than one sibling had a relatively greater incidence of “poor” knowledge unlike mothers of children with single or no sibling. This implies that those with “poor” knowledge were also unaware of the importance of birth spacing. This stresses the importance of the WHO’s exclusive breast feeding in the early infancy sufficing the lesser iron requirements. Only 17% of mothers were aware of the importance of timely introduction of complementary feeds, proving the misconceptions on duration of breast feeding. Iron requirements after 6 months are 0.9–1.3 mg/kg/day [25], whereas the iron content of cow’s milk is 0.2–0.5 mg/L, of which only 10% is absorbed. [26] In addition, cow’s milk causes asymptomatic micro-hemorrhages in intestine further increasing the loss of iron. [27] This explains the reason why excess cow’s milk had a negative influence on hemoglobin level in our study.

Several mothers believed that only children on vegetarian diet would be affected by IDA. However, iron status is a common nutritional problem among both vegetarian and nonvegetarian consumers, despite few studies reporting IDA of higher incidence in the former. This relative increase in IDA among vegetarians is probably due to their dependence on non-heme iron and the presence of iron absorption inhibitors in plant foods. [28] Apt feeding practices are thus fundamentally important to ensure an appropriate nutrition in a growing child. Clinical features are non-specific in anemic children as evidenced in our study, which when untreated lead to neurodevelopment delay and cognitive deficits. Their attention span is often reduced and this reflects on the child’s academic performance. [29] On a positive note, most of the mothers had acknowledged the possibility of neural development being affected by anemia. Understanding this would stress the importance of supplementing iron in infants. The prime source of knowledge in our study was the community health workers. Training and engagement of the health workers are critical in increasing the healthcare awareness in low- to middle-income countries. [30] Parenting knowledge plays a key role in the biological, physical, socioeconomic, and cognitive needs of the child. It also has a direct influence on their everyday decisions about upbringing, developmental expectations which, in turn, determines their child’s health and well-being. [31] Overall most of the mothers had “fair” knowledge of anemia and its implications. Mothers with “poor” understanding of nutrition predominantly had anemic children. Furthermore, mothers with better educational status had better awareness in terms of questionnaire. Thus, maternal knowledge plays a key role in preventing anemia.

## Conclusion

Despite the advent of the internet available in everyone’s hand, the knowledge of anemia remains shallow. Nutrition in the latter half of infancy is vital. Information regarding the negative influences of cow’s milk on hemoglobin ought to be spread. Mothers must be enlightened on the importance and source of nourishment in a weaning child. Improving maternal educational status will address the concerns on IDA. Ultimate goal must be to target the mothers to take rational decisions rather than believing the faculty of age-old aphorisms. Role of Pediatricians in creating adequate awareness among these mothers is enormous.

## References

1. Ekwere TA, Ekanem AM, Ekwere TA. Maternal knowledge, food restriction and prevention strategies related to anaemia in pregnancy: a cross-sectional study. *International journal of community medicine and public health*. 2015 Jul;2(3):331-8.
2. WHO Prevalence of anaemia in pregnant women (aged 15-49) (%) 2021.
3. The World Bank Prevalence of anemia among pregnant women (%) -Least developed countries: UN classification | Data 2019
4. Shah T, Khaskheli MS, Ansari S, Lakhan H, Shaikh F, Zardari AA, Warsi J, Rind NA, Rind KH, Shar AH. Gestational Anemia and its effects on neonatal outcome, in the population of Hyderabad, Sindh, Pakistan. *Saudi J Biol Sci*. 2022 Jan;29(1):83-87.
5. Sari YA, Wahiduddin W, Ansariadi A, Bustan N, Russeng SS, Sirajuddin S, Wijaya E. Determinants of low birth weight occurrence in Bahteramas regional general hospital, Southeast Sulawesi Province, Indonesia. *Open Access Macedonian Journal of Medical Sciences*. 2022 Jul 3;10(E):1107-13.
6. Hermayanti D, Setiawan M, Nurdiana H. Improvement of knowledge on iron deficiency anemia and its prevention among Bustanul Athfal teachers in Malang.
7. Roland, M. C. P. et al. Maternal factors associated with fetal growth and birthweight are independent determinants of placental weight and exhibit differential effects by fetal sex. *PLoS ONE* 9(2), e87303 (2014).
8. Esmailzadeh, A., Samareh, S. & Azadbakht, L. Dietary patterns among pregnant women in the west-north of Iran. *Pak. J. Biol. Sci. PJB* 11(5), 793 (2008).
9. Abedini, Z. & Gaini, M. Dietary food intake of pregnant women based on food guide pyramid and its related factors. *Iran J. Nurs*. 24(73), 36–46 (2011).
10. Woldeamanuel, G. G., Geta, T. G., Mohammed, T. P., Shuba, M. B. & Bafa, T. A. Effect of nutritional status of pregnant women on birth weight of newborns at Butajira Referral

- Hospital, Butajira, Ethiopia. *SAGE Open Med.* 7, 2050312119827096 (2019).
11. Balcha WF, Eteffa T, Arega Tesfu A, Abeje Alemayehu B. Maternal Knowledge of Anemia and Adherence to its Prevention Strategies: A Health Facility-Based Cross-Sectional Study Design. *INQUIRY: The Journal of Health Care Organization, Provision, and Financing.* 2023 Apr;60:00469580231167731.
  12. De Onis M, Garza C, Onyango AW, Rolland-Cachera MF. WHO growth standards for infants and young children. *Arch Pediatr* 2008; 16:47-53.
  13. Saleem SM, Jan SS. Modified Kuppusswamy socioeconomic scale updated for the year 2019. *Indian J Forensic Community Med* 20 19;6:1-3.
  14. Matthews M, Nigg JT, Fair DA. Attention Deficit Hyperactivity Disorder. *Curr Top Behav Neurosci* 2014;16: 235– 66.
  15. Corp IBM. IBM SPSS Statistics for Windows. Version 22.0. Armonk, NY:IBM Corp; 2013.
  16. World Health Organization. Nutritional Anaemias: Tools for Effective Prevention and Control. Geneva: World Health Organization; 2017.
  17. Kassebaum NJ, Jasrasaria R, Naghavi M, Wulf SK, Johns N, Lozano R, Regan M, Weatherall D, Chou DP, Eisele TP, Flaxman SR. A systematic analysis of global anemia burden from 1990 to 2010. *Blood, the Journal of the American Society of Hematology.* 2014 Jan 30;123(5):615-24.
  18. World Health Organization. Iron Deficiency Anemia. Assessment, Prevention, and Control. A Guide for Programme Managers. Geneva: World Health Organization; 2001. p. 47-62.
  19. Uijterschout L, Vloemans J, Vos R, Teunisse PP, Hudig C, Bubbers S, Verbruggen S, Veldhorst M, de Leeuw T, van Goudoever JB, Brus F. Prevalence and risk factors of iron deficiency in healthy young children in the southwestern Netherlands. *Journal of pediatric gastroenterology and nutrition.* 2014 Feb 1; 58(2):193-8.
  20. Lozoff B, Jimenez E, Hagen J, Mollen E, Wolf AW. Poorer behavioral and developmental outcome more than 10 years after treatment for iron deficiency in infancy. *Pediatrics.* 2000 Apr 1;105(4):e51-.
  21. Cerami C. Iron nutriture of the fetus, neonate, infant, and child. *Annals of Nutrition and Metabolism.* 2017 Dec 21;71(Suppl. 3):8-14.
  22. Koleini N, Shapiro JS, Geier J, Ardehali H. Ironing out mechanisms of iron homeostasis and disorders of iron deficiency. *The Journal of Clinical Investigation.* 2021 Jun 1;131(11).
  23. Means RT. Iron deficiency and iron deficiency anemia: implications and impact in pregnancy, fetal development, and early childhood parameters. *Nutrients.* 2020 Feb 11;12(2):447.
  24. Kassebaum NJ, Jasrasaria R, Naghavi M, Wulf SK, Johns N, Lozano R, Regan M, Weatherall D, Chou DP, Eisele TP, Flaxman SR. A systematic analysis of global anemia burden from 1990 to 2010. *Blood, the Journal of the American Society of Hematology.* 2014 Jan 30;123(5):615-24.
  25. Domellöf M. Iron requirements in infancy. *Annals of Nutrition and Metabolism.* 2011 Nov 1;59(1):59-63.
  26. Saidalikutty FM, Sugumar R, Shanmugam K. Exclusive cow's milk intake and asymptomatic anaemia. *Indian Journal of Case Reports.* 2019 Aug 26:311-2.
  27. Wilson JF, Lahey ME, Heiner DC. Studies on iron metabolism: V. Further observations on cow's milk-induced gastrointestinal bleeding in infants with iron-deficiency anemia. *The Journal of Pediatrics.* 1974 Mar 1;84(3):335-44.
  28. Pawlak R, Bell K. Iron status of vegetarian children: a review of literature. *Annals of Nutrition and Metabolism.* 2017 Mar 21;70(2):88-99.
  29. Lukowski AF, Koss M, Burden MJ, Jonides J, Nelson CA, Kaciroti N, Jimenez E, Lozoff B. Iron deficiency in infancy and neurocognitive functioning at 19 years: evidence of long-term deficits in executive function and recognition memory. *Nutritional neuroscience.* 2010 Apr 1;13(2):54-70.
  30. Rowe SY, Kelly JM, Olewe MA, Kleinbaum DG, McGowan Jr JE, McFarland DA, Rochat R, Deming MS. Effect of multiple interventions on community health workers' adherence to clinical guidelines in Siaya district, Kenya. *Transactions of the Royal Society of Tropical Medicine and Hygiene.* 2007 Feb 1;101(2):188-202.
  31. Bornstein MH, Yu J, Putnick DL. Mothers' parenting knowledge and its sources in five societies: Specificity in and across Argentina, Belgium, Italy, South Korea, and the United States. *International Journal of Behavioral Development.* 2020 Mar;44(2):135-45.