

An Observational Study to Assess the Histopathological Spectrum of Uterine Lesions in Perimenopausal and Postmenopausal Women

Madhu Bharti¹, Shipra Singh², Richa Sharma³

¹Tutor, Department of Pathology, Darbhanga Medical College and Hospital, Darbhanga, Bihar, India

²Tutor, Department of Pathology, Darbhanga Medical College and Hospital, Darbhanga, Bihar, India

³Tutor, Department of Pathology, Darbhanga Medical College and Hospital, Darbhanga, Bihar, India

⁴Professor and HOD, Department of Pathology, Darbhanga Medical College and Hospital, Darbhanga, Bihar, India

Received: 17-02-2024 / Revised: 14-03-2024 / Accepted: 29-04-2024

Corresponding author: Dr. Shipra Singh

Conflict of interest: Nil

Abstract

Aim: The aim of the present study was to assess the histopathological spectrum of uterine lesions in perimenopausal and postmenopausal women.

Methods: The present study was conducted for a period of 1 year in the Department of Pathology, Darbhanga Medical College and Hospital, Darbhanga, Bihar, India. The study comprised of 100 cases which met the inclusion criteria were taken for this study.

Results: The maximum number of cases 24 (24%) were between the age group of 46-50 years. The most common site biopsy received was from cervix, 43 (43%). Postmenopausal bleeding was due to benign causes. Atrophic endometrium was the commonest benign.

Conclusion: PMB is a symptom which should not be taken lightly. Accurate diagnosis is usually made by histopathological examination. In our study, a wide spectrum of both neoplastic and non- neoplastic conditions of female genital tract has displayed as a cause of PMB with predominance of benign causes. Cervical cancer is still the most common cause of PMB, which point out that the effective implementation of screening program is utmost important. More awareness among people, especially elderly women should be made about the importance of pap screening.

Keywords: histopathological spectrum, uterine lesions, perimenopausal, postmenopausal women

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

Introduction

Abnormal uterine bleeding (AUB) is the most common presentation amongst the women of all age groups attending the Gynecology Out Patient Department. AUB is the term used to describe any departure from normal menstruation or from a normal menstrual cycle pattern. The key characteristics are regularity, frequency, heaviness of flow and duration of flow. [1] Menstrual bleeding is considered abnormal when the pattern is irregular of prolonged duration (i.e.>20 days) in individual cycle lengths over a period of 1 year or heavy menstrual bleeding. Recently, the International Federation of Gynecology and Obstetrics working group on menstrual disorder has developed a classification system (PALM-COEIN) for causes of the AUB in non-gravid women of reproductive age. There are nine main categories, which are arranged according to the acronym PALM-COEIN: Polyp; adenomyosis; leiomyoma; malignancy and

hyperplasia; coagulopathy; ovulatory dysfunction; endometritis; iatrogenic; and not yet classified. [2]

Two types of causes including functional causes like normal cyclical endometrium, abnormal physiological changes of endometrium (atrophic endometrium, weakly proliferative and disordered proliferative endometrium) and organic lesions like chronic endometritis, endometrial hyperplasia, endometrial polyp, carcinomas and pregnancy related complications. [2] In reproductive age group heavy menstrual bleeding accounts for 30% of total DUB cases [3], while postmenopausal bleeding accounts for 5% of all gynaecological visits. Histopathological examination of endometrial sample is used for the diagnosis of majority of lesions. Endometrial biopsy and curettage are chief sampling methods. [4]

UB is due to several factors deranging homeostasis like hormonal imbalances, infections, structural

lesions, and malignancy. Based on these possible underlying etiologies, the International Federation of Gynaecology and Obstetrics (FIGO) in 2011 devised a classification named PALM-COEIN for the etiology of AUB. PALM accounts for structural features like polyps, adenomyosis, leiomyoma, and malignancy. COEIN addresses non-structural causes like coagulation defects, ovulatory dysfunction, endometrial causes, iatrogenic causes, and non-classified ones. [5] Endometrial biopsy is used as a diagnostic aid in AUB. It is done as a first-line test in women >45 years of age presenting with AUB. Endometrial biopsy is also done in patients <45 years of age with a history of unopposed estrogen exposure, failed medical management, and persistent AUB.⁶ The prime idea is to rule out the precursor lesions like hyperplasia and aggressive endometrial carcinoma. [5]

The aim of the present study was to assess the histopathological spectrum of uterine lesions in perimenopausal and postmenopausal women.

Materials and Methods

The present study was conducted for a period of one year in the Department of Pathology, Darbhanga

Medical College and Hospital, Darbhanga, Bihar, India. The study comprised of 100 cases which met the inclusion criteria were taken for this study. Material for study was collected from endometrial, cervical, vaginal, vulval biopsies and hysterectomy specimens which were sent for histopathological examination to Department of Pathology from clinically diagnosed cases of postmenopausal bleeding. The History noted were spotting per vagina, brownish discharge, scanty flow and moderate to profuse bleeding. Premature menopause whether surgical or natural, with age <40yrs and patient on hormonal replacement therapy/on anticoagulant/having bleeding disorders were excluded from the study. After the collection of detailed data, the specimens were examined grossly. Specimens were fixed in 10% buffered formalin. Sections were processed and stained with H & E. Slides were examined under microscope and observations were done. The results were compiled, analysed using proportion and compared with other studies.

Results

Table 1: Distribution of study subjects according to age groups

AGE (years)	No of studysubjects	Percentage
41-45	8	8
46-50	24	24
51-55	20	20
56-60	18	18
61-65	13	13
66-70	10	10
>70	6	6
>80	1	1
Total	100	100

The maximum number of cases 24 (24%) were between the age group of 46-50 years.

Table 2: Type of specimens received

Type of specimen	Numbers	Percentage
Biopsy vulva and vagina	3	3
Biopsy cervix	43	43
Biopsy cervix and endometrium	9	9
Endometrial curettage	31	31
Hysterectomy	5	5
Hysterectomy with adnexa	8	8
Hysterectomy with adnexa, omentum and lymph nodes	1	1
Total	100	100

The most common site biopsy received was from cervix, 43 (43%).

Table 3: Distribution and percentage of various lesions

Histopathology diagnosis	Number
Inadequate	9
Cervicitis	4
Cervical polyp	4
Atrophic endometrium	15
proliferative endometrium	10
Endometrial hyperplasia without atypia	8
Endometrial hyperplasia with atypia	2
Endometrial polyp	3
Endometritis	1
Adenomyosis	3
Leiomyoma	5
Prolapse	1
Carcinoma cervix(total)	45
CIN	8
SCC	35
Undifferentiated	2
Malignant Uterus (Total)	12
Adenocarcinoma	7
Adenosquamous carcinoma	2
Papillary Serous Carcinoma	2
MMMT	1
Carcinoma Vulva (SCC)	3
Carcinoma Vagina (SCC and Adenocarcinoma)	2

Atrophic endometrium was the commonest benign. Most of the malignant tumors were from the cervix

Table 4: Distribution of lesions

S. No	Histopathology	Number of cases
1	Benign	70
2	Malignant	50
3	Premalignant	10

Postmenopausal bleeding was due to benign causes.

Discussion

The endometrium is a constantly changing tissue that is sensitive and responsive to hormones, particularly during active reproductive years. Among patients of all ages who are coming to gynaecology outpatient departments, excessive and irregular bleeding remains a common presenting symptom. [7] Abnormal uterine bleeding is characterized by a deviation in frequency, duration, or amount of bleeding from what is typically observed during a normal menstrual cycle or after menopause. [8]

The investigations and assessment is moving away from operation theatre, ward environment into outpatient department. However the primary assessment in all cases of PMB should be trans vaginal ultrasound scanning(TVS) as the thickening of endometrium may indicate significant pathology. [9] The present trend in investigating only lesions with PMB when endometrial thickness is >4mm as

measured by ultrasound. [10] However the authors have recommended systematic collection of biopsies from symptomatic patients [11] because there have been reports of cancer in patients presenting with ultrasound measured endometrial thickness <5mm. [12] The maximum number of cases 24 (24%) were between the age group of 46-50 years while the study done by Way sf et al, Sousa R et al, Bharani B et al and Sheikh M et al was 38-94, 43-82,52-65, 42-84 yrs respectively. [13-16] The most common site biopsy received was from cervix, 43 (43%). Postmenopausal bleeding was due to benign causes. Atrophic endometrium was the commonest benign.

Choo YC et al [17] found out that stimulation of postmenopausal endometrium can occur because of conversion of adrenal and rosteinedione by peripheral fat to estrogen which leads to proliferative endometrium and also fluctuation of low level of estrogen results in bleeding from proliferative endometrium. Menorrhagia is the commonest presentation of AUB found in studies done by Sharma et al., Sajitha et al. and

Mukhopadhyay et al. [18-20] In endometrial polyp bleeding can be as a result of injury to thin walled vein below surface epithelium or thrombosis of the vessels. The bleeding in leiomyoma can occur due to congestion or atrophy & thinning of overlying endometrium and myometrium results in ulceration and bleeding. [21]

Conclusion

PMB is a symptom which should not be taken lightly. Accurate diagnosis is usually made by histopathological examination. In our study, a wide spectrum of both neoplastic and non-neoplastic conditions of female genital tract has displayed as a cause of PMB with predominance of benign causes. Cervical cancer is still the most common cause of PMB, which point out that the effective implementation of screening program is utmost important. More awareness among people, especially elderly women should be made about the importance of pap screening. PMB indicates malignancy until proved otherwise and it demands thorough evaluation of patients with histopathological confirmation. An accurate diagnosis is immensely important as it will be helpful for the management of patient by implementing a proper treatment plan.

References

- Fraser IS, Critchley HO, Broder M, Munro MG. The FIGO recommendations on terminologies and definitions for normal and abnormal uterine bleeding. In Seminars in reproductive medicine 2011 Sep (Vol. 29, No. 05, pp.383-390).© Thieme Medical Publishers.
- Munro MG, Critchley HO, Broder MS, Fraser IS, FIGO Working Group on Menstrual Disorders. FIGO classification system (PALM-COEIN) for causes of abnormal uterine bleeding in nonpregnant women of reproductive age. International Journal of Gynecology & Obstetrics. 2011 Apr 1;113(1):3-13.
- Moodley M, Roberts C. Clinical pathway for the evaluation of postmenopausal bleeding with an emphasis on endometrial cancer detection. J Obstet Gynaecol. 2004 Oct;24(7):736-41.
- Dueholm M. Levonorgestrel-IUD should be offered before hysterectomy for abnormal uterine bleeding without uterine structural abnormalities: there are no more excuses! Acta Obstet Gynecol Scand 2009;88(12):1302-1304.
- Mishra D, Sultan S. FIGO's PALM-COEIN Classification of Abnormal Uterine Bleeding: A Clinico-histopathological Correlation in Indian Setting. J Obstet Gynaecol India. 2017 Apr;67(2):119-125.
- ACOG committee opinion no. 557: Management of acute abnormal uterine bleeding in nonpregnant reproductive-aged women. Obstet Gynecol. 2013 Apr;121(4): 891-896.
- Fa T. Pathology and Genetics. Tumours of the breast and female genital organs. Peritoneal tumours. 2003:197-202.
- Ely JW, Kennedy CM, Clark EC, Bowdler NC. Abnormal uterine bleeding: a management algorithm. The Journal of the American Board of Family Medicine. 2006 Nov 1;19(6):590-602.
- Wolman I, Amster R, Hartoov J et al. Reproducibility of transvaginal ultrasonographic measurements of endometrial thickness in patients with postmenopausal bleeding. Gynecol Obstet Invest 1998;46:191-94.
- Biriniyi L, Darago P, Torok P et al. Predictive value of hysteroscopic examination in intrauterine abnormalities. Eur J. Obstet Gynecol. Reprod Biol 2004;115(1):75-79.
- Tabor A, Watt HC, Wald NJ. Endometrial thickness as a test for endometrial cancer in women with postmenopausal bleeding. Obstet Gynecol 2002;99(4):663-70.
- Litta P, Merlin F, Saccordic C et al. Role of hysteroscopy with endometrial biopsy to rule out endometrial cancer in postmenopausal women with abnormal uterine bleeding. Marturitas 2005;50(2):117-23.
- Wong SF, Luk KL, Wong AY, Tang LC. Finding in women with postmenopausal bleeding investigated with hysteroscopy. J Obstet Gynecol 2001;21:392-95.
- Sousa R, Silvestre M, Almedia e Sousa, Falco F, Dias I, Silva T, De Oliveria C, Oliveria HM. Transvaginal ultrasonography and hysteroscopy in postmenopausal bleeding a prospective study. Acta Obstet Gynecol Scand 2001;80:856-62.
- Bharani B, Phatak SR. Feasibility and yield of endometrial biopsy using suction curette device for evaluation of abnormal pre and postmenopausal bleeding. J Obstet Gynecol India. 2008;58:322-6.
- Sheikh M, Sawhney, Khurana A, Al Yatama M. Alteration of sonographic texture of the endometrium in PMB a guide to further management. Acta Obstet Gynecol Scand. 2009;79:1006-50.
- Choo YC, Mak KC, Hsu C, Wong TS, Ma HK. Postmenopausal uterine bleeding of nonorganic cause. Obstet Gynecol 1985;66:225-28.
- Sharma R, Mishra P, Kumar N, Srivastava P. Histomorphological spectrum of endometrial lesion in women presenting with abnormal uterine bleeding: A 3-year study at a tertiary care center. Trop J Pathol Microbiol. 2018;4: 525-31.
- Sajitha K, Padma SK, Shetty JK, Kishan Prasad HL, Permi HS, Hegde P. Study of

- histopathological patterns of endometrium in abnormal uterine bleeding. CHRISMED journal of health and Research. 2014 Apr 1;1 (2):76-81.
20. Mukhopadhyay I, Rao PS, Nataraj S, Biswas M. An analysis of endometrial bleeding patterns in perimenopausal women. Int J Reprod Contracept Obstet Gynecol. 2017;6: 2776-83.
21. Tyagi R, Isaacs R, Dhar T. Postmenopausal bleeding: histopathological spectrum and association with age and clear span: case series of 328 cases. Journal of Evolution of Medical and Dental Sciences. 2014 Jun 30;3(26):7210-22.