

A Retrospective Study Assessing Role of Diabetes Mellitus in Causing Posterior Subcapsular Cataracts in Outpatients

Nandani Priyadarshini¹, Rajesh Kumar Tiwary²

¹Senior Resident, Department of Ophthalmology, Nalanda Medical College and Hospital, Patna, Bihar, India

²Professor, Department of Ophthalmology, Nalanda Medical College and Hospital, Patna, Bihar, India

Received: 10-03-2024 / Revised: 14-04-2024 / Accepted: 21-05-2024

Corresponding Author: Dr. Nandani Priyadarshini

Conflict of interest: Nil

Abstract

Aim: This study aimed to determine the role of diabetes mellitus in causing cataracts, especially posterior subcapsular cataracts.

Methods: The present study was conducted in the Department of ophthalmology, Nalanda Medical College and Hospital, Patna, Bihar, India March 2016 to February 2017. The consecutive sampling method used in this study the overall data of cataract patients treated by the Department of ophthalmology. This data is a medical record containing a history of diabetes mellitus and the types of cataracts suffered by 100 patients.

Results: In this study, 100 patients' data were used; it is known that there were 36 patients with a history of diabetes mellitus (36%) and history of non-diabetes Mellitus as many as 64 patients (64%). As for 100 patients who suffer from cataract eye disorders or diseases in this study, it was found that 25 patients (25%), had PSC type cataracts, and 75 patients (75%) had non-PSC type cataracts. From the cross-section analysis results, it was known that as many as 21 patients had a history of diabetes mellitus and had PSC cataracts, while 4 patients, did not have a history of diabetes mellitus but had PSC cataracts. In addition, the results of the analysis also showed that there were 15 patients had a history of diabetes mellitus and had non-PSC cataracts, while 60 patients did not have a history of diabetes mellitus experiencing non-PSC cataracts. Overall, there were 36 patients (36%) had a history of diabetes mellitus and had cataracts, and there were 64 patients (64% who did not have a history of diabetes mellitus but had cataracts. Thus, it can be concluded that diabetes mellitus has a relationship with cataracts, especially PSC cataracts. This condition is proven by more patients with PSC cataracts who have a history of diabetes mellitus than patients who suffer from PSC cataracts but do not have a history of diabetes.

Conclusion: As the number of people with DM is estimated to continue to increase, cataract surgery will remain important for diabetic patients. Patients with diabetes have multiple issues to be evaluated preoperatively, perioperatively, and in the postoperative period.

Keywords: Cataract; Diabetes Mellitus; PSC Cataract

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

Introduction

The eye is an essential visual organ for humans. [1] With the eye, human beings can get a variety of visual information transmitted to the brain properly to be processed as a basis for decision making in carrying out daily activities. Every organ in the eye has a vital function for humans because if one of the eye functions is problematic, it will affect the function of other organs. Eyes need lenses that are clear, transparent, and flexible or elastic. If the lens in the eye loses its translucency or clarity, the vision will become foggy and even cause a person unable to see at all [2], this is because the cloudiness that occurs in the lens makes the amount of incoming light decrease and causes a person unable to see correctly. [3] Turbidity or loss of translucency that occurs in the lens is called a cataract. Turbidity in

the lens or cataracts can occur due to hydration of the lens fluid, denaturation of the lens protein, or a result of both. [4]

According to data reported by the World Health Organization [5], it is estimated that 2.2 billion people suffer from vision problems, with 65.2 million people in the world suffering from cataracts. Cataracts consist of four types: secondary cataracts, senile cataracts, complicated cataracts, and traumatic cataracts. [6] In a population over 50 years old, most people suffered from senile cataracts consisting of nuclear cataracts, cortical cataracts, and Posterior Subcapsular Cataracts (PSC) caused by lens degeneration.⁷ PSC cataract patients may not realize they have cataracts if the visual disturbance

does not occur in the center of the lens because visual disturbances in cataract patients occur slowly and are only realized when their eyesight is getting worse, even when patients experience blindness. [8] Cataract is one of the major causes of visual impairment in diabetic patients. [9] Patients with DM are reported to be up to five times more likely to develop cataract, in particular at an early age [5-8]. Due to the increasing prevalence of DM, the incidence of diabetic cataracts has also risen. Cataract extraction is one of the most common surgical procedures among the general population, and the number of cataract surgeries each year also continues to increase. Recent technological advancements in cataract surgery have improved surgical outcomes. [10-13]

This study aimed to determine the role of diabetes mellitus in causing cataracts, especially posterior subcapsular cataracts.

Methods

The present study was conducted in the Department of ophthalmology, Nalanda Medical College and Hospital, Patna, Bihar, India from March 2016 to February 2017. The consecutive sampling method used in this study the overall data of cataract patients treated by the Department of ophthalmology. This data is a medical record containing a history of diabetes mellitus and the types of cataracts suffered by 100 patients.

The Criteria Set Are:

- 1. Cataract patients over 45 years old

- 2. Experiencing PSC cataract or Non-PSC cataract eye disease
- 3. Have a history of diabetes mellitus or non-diabetes mellitus

The data obtained in this study were then processed and analyzed with a cross-sectional design and chi-square test using IBM SPSS 26 software to determine the proportion of data distribution and the relationship between the independent and dependent variables.

In this study, the independent variable was diabetes mellitus, while the dependent variable was PSC cataract. The analysis results were recorded in a 2x2 table. The hypotheses proposed in this study are as follows.

H0 : History of diabetes mellitus does not cause PSC cataract eye disease

H1 : History of diabetes mellitus causes PSC cataract eye disease

In addition to the retrospective analysis, a regression test was also conducted to determine the value of the determinant coefficient (R²), which shows the magnitude of the influence of Diabetes Mellitus disease history (independent variable) in causing PSC cataract (dependent variable). If the R² value shows a value of 0.67 or more, then the independent variable significantly influences the dependent variable. Meanwhile, if the R² value shows a value of 0.33 or 0.19, then each independent variable has a moderate or weak influence on the dependent variable.¹⁹

Results

Table 1: Number of diabetes and cataract patients

History of Disease	N%
Diabetes Mellitus	36 (36)
Non-Diabetes Mellitus	64 (64)
Type of cataract	
PSC	25 (25)
Non PSC	75 (75)

In this study, 100 patients' data were used; it is known that there were 36 patients with a history of diabetes mellitus (36%) and history of non-diabetes Mellitus as many as 64 patients (64%). As for 100

patients who suffer from cataract eye disorders or diseases in this study, it was found that 25 patients (25%), had PSC type cataracts, and 75 patients (75%) had non-PSC type cataracts.

Table 2: Cross-Section Test Results

History of disease	Type of cataract		Total (%)
	PSC N	Non PSC N	
Diabetes Mellitus	21	15	36 (36)
Non Diabetes Mellitus	4	60	64 (64)
Total	25	75	
Chi-Square Test			
Pearson Chi Square	86.44	1.0000	0.000

From the cross-section analysis results, it was known that as many as 21 patients had a history of diabetes mellitus and had PSC cataracts, while 4 patients, did not have a history of diabetes mellitus but had PSC cataracts. In addition, the results of the analysis also showed that there were 15 patients had a history of diabetes mellitus and had non-PSC cataracts, while 60 patients did not have a history of diabetes mellitus experiencing non-PSC cataracts. Overall, there were 36 patients (36%) had a history of diabetes mellitus and had cataracts, and there were 64 patients (64% who did not have a history of diabetes mellitus but had cataracts. Thus, it can be concluded that diabetes mellitus has a relationship with cataracts, especially PSC cataracts. This condition is proven by more patients with PSC cataracts who have a history of diabetes mellitus than patients who suffer from PSC cataracts but do not have a history of diabetes. In this study, a confidence interval of 95% or α 0.05 was used. Thus, to accept hypothesis H1, the Pearson Chi-Square value on the resulting Chi-Square test must be less than the cut-off value of 0.05.

Discussion

Diabetes mellitus is reported to cause cataracts due to long-term uncontrolled hyperglycemia and significant fluctuations in blood glucose levels. [14] This condition is because most of the glucose breakdown in the lens 78% is via the anaerobic glycolysis pathway, 14% via the pentose phosphate pathway, and about 5% via the polyol pathway. In conditions of hyperglycemia, the anaerobic glycolysis pathway saturates quickly, and glucose selects the polyol pathway. In the polyol pathway, glucose is converted to sorbitol. Sorbitol is then broken down into fructose by the Polyol Dehydrogenase enzyme. However, in diabetes mellitus, the Polyol Dehydrogenase enzyme levels are low, so sorbitol accumulates in the eye lens.

In this study, 100 patients' data were used; it is known that there were 36 patients with a history of diabetes mellitus (36%) and history of non-diabetes Mellitus as many as 64 patients (64%). As for 100 patients who suffer from cataract eye disorders or diseases in this study, it was found that 25 patients (25%), had PSC type cataracts, and 75 patients (75%) had non-PSC type cataracts. From the cross-section analysis results, it was known that as many as 21 patients had a history of diabetes mellitus and had PSC cataracts, while 4 patients, did not have a history of diabetes mellitus but had PSC cataracts. In addition, the results of the analysis also showed that there were 15 patients had a history of diabetes mellitus and had non-PSC cataracts, while 60 patients did not have a history of diabetes mellitus experiencing non-PSC cataracts. Overall, there were 36 patients (36%) had a history of diabetes mellitus and had cataracts, and there were 64 patients (64% who did not have a history of diabetes mellitus but

had cataracts. Thus, it can be concluded that diabetes mellitus has a relationship with cataracts, especially PSC cataracts. This condition is proven by more patients with PSC cataracts who have a history of diabetes mellitus than patients who suffer from PSC cataracts but do not have a history of diabetes. In this study, a confidence interval of 95% or α 0.05 was used. Thus, to accept hypothesis H1, the Pearson Chi-Square value on the resulting Chi-Square test must be less than the cut-off value of 0.05. This study was in line with research conducted by Ehrlich RM and Taskapili M which states that anterior and subcapsular posterior are structural characteristics of cataracts in diabetic patients. [15,16]

In testing the hypothesis in this study, the chi-square test was carried out. As known in the previous analysis, the resulting Pearson chi-square value is 0.000 less than the cut-off value of 0.05. Thus, hypothesis H1 is accepted, which means that a history of diabetes mellitus can cause PSC cataract eye disease. Then, in the R2test, it was found that the PSC cataract eye disease was influenced by a history of diabetes mellitus by 39.1% (R2: 0.391); this means that the history of diabetes mellitus has a moderate effect in causing PSC cataracts, and as much as 60.9% is influenced by other factors that were not examined in this study. This finding supports other studies that say that the cause of PSC cataracts is multifactorial with other factors, not only by one thing, such as diabetes mellitus. [17]

The results of this study also support several studies that have shown that diabetes mellitus can cause cataracts to occur more frequently than non-diabetics. Framingham and other eye studies showing a three to fourfold increase in cataract prevalence in patients with diabetes under 65 years old and a two-fold increase in patients over 65 years old. The results of hypothesis testing in this study also in line with The Visual Impairment Project, which evaluates risk factors for cataract development in Australia. The study showed that diabetes mellitus is an independent risk factor for posterior subcapsular cataract if it occurs for more than five years. [19] In addition, eight studies in Europe, America, Africa, and Australia on 20,837 cataract patients using a meta-analysis method showed that type 2 diabetes mellitus has an increased risk of developing posterior subcapsular cataracts (OR=1.55,95% CI:1.27-1.90, P<0.001). [18] Medicare analysis from 1997 to 2001 revealed that the diagnosis rate of postoperative pseudophakic cystoid macular edema (PCME) was higher in diabetic patients than in non-diabetic patients. [19] So that PCME prophylaxis should be done immediately depending on the stage of diabetic retinopathy. However, if diabetic retinopathy is found, patients with PSC cataracts should delay surgery or cataract extraction. [20]

Conclusion

As the number of people with DM is estimated to continue to increase, cataract surgery will remain important for diabetic patients. Patients with diabetes have multiple issues to be evaluated preoperatively, perioperatively, and in the postoperative period. With the advent of modern surgical and pharmacologic therapies, these patients can, like other cataract patients without diabetes, recover excellent vision. Postoperative monitoring and management of surgical complications will also help to alleviate the risk of vision loss in these patients.

References

1. Amaliah AN, Febriza AM. The Relations Between Diabetes Mellitus Type 2 on The Incidence Cataract in Balai Kesehatan Mata Makassar in 2016. *MAGNA MEDIKA: Berkala Ilmiah Kedokteran dan Kesehatan*. 2019;6(1):99-106.
2. Ilyas S. Ilmu penyakit mata. In: Yulianti SR, editor. Jakarta: Badan Penerbit Fakultas kedokteran Universitas Indonesia; 2014.
3. Gupta VB, Rajagopala M, Ravishankar B. Etiopathogenesis of cataract: an appraisal. *Indian journal of ophthalmology*. 2014 Feb;62(2):103.
4. Nadyanti F. Hubungan durasi menderita diabetes melitus tipe 2 dengan kejadian katarak di RSUD DR. H. Abdul Moeloek Provinsi Lampung Tahun 2018.
5. Organization WH. World report on vision.
6. Girsang RR, Fahmi H. Expert System for Diagnosing Cataract Eye Disease with Web-Based Certainty Factor Method. *Matics*. 2019 Oct 10; 11(1): 27-31.
7. Hasriani RD, Syahrizal S, Misti M. Hipertensi dengan Katarak pada Peserta Skrining Gangguan Penglihatan. *HIGEIA (Journal of Public Health Research and Development)*. 2020 Oct 31;4(4):645-55.
8. Sunardi B, Iskandar E, Tamzil NS. HUBUNGAN PENGGUNAAN KORTIKOSTEROID PADA PASIEN SYSTEMIC LUPUS ERYTHEMATOSUS TERHADAP KEJADIAN KATARAK DI RUMAH SAKIT MOHAMMAD HOESIN (Doctoral dissertation, Sriwijaya University).
9. Drinkwater JJ, Davis WA, Davis TM. A systematic review of risk factors for cataract in type 2 diabetes. *Diabetes/metabolism research and reviews*. 2019 Jan;35(1):e3073.
10. Klein BE, Klein R, Wang Q, Moss SE. Older-onset diabetes and lens opacities. The Beaver Dam Eye study. *Ophthalmic epidemiology*. 1995 Jan 1;2(1):49-55.
11. Klein BE, Klein R, Lee KE. Diabetes, cardiovascular disease, selected cardiovascular disease risk factors, and the 5-year incidence of age-related cataract and progression of lens opacities: the Beaver Dam Eye Study. *American journal of ophthalmology*. 1998 Dec 1;126(6):782-90.
12. Klein BE, Klein R, Moss SE. Incidence of cataract surgery in the Wisconsin Epidemiologic Study of Diabetic Retinopathy. *American journal of ophthalmology*. 1995 Mar 1;119(3):295-300.
13. Saxena S, Mitchell P, Rochtchina E. Five-year incidence of cataract in older persons with diabetes and pre-diabetes. *Ophthalmic epidemiology*. 2004 Jan 1;11(4):271-7.
14. Wu J, Zeng H, Xuan R, Lei S, Li J, Lai X, Liu J. Bilateral cataracts as the first manifestation of type 1 diabetes mellitus: a case report. *Medicine*. 2018 Oct;97(42).
15. Ehrlich RM, Kirsch S, Daneman D. Cataracts in children with diabetes mellitus. *Diabetes Care*. 1987 Nov 1; 10(6): 798-9.
16. Taskapili M, Gulkilik G, Ozsutcu M, Kucuksahin H. Acute bilateral dense cortical cataracts as a first manifestation of juvenile diabetes mellitus in a 12-year-old girl. *J Pediatr Ophthalmol Strabismus*. 2008 May-Jun ; 45(3): S177-8.
17. Vaughan D, Asbury T. *General ophthalmology*. 17th (edtn). California: Lange Medical Publication; 2007.
18. Li L, Wan XH, Zhao GH. Meta-analysis of the risk of cataract in type 2 diabetes. *BMC ophthalmology*. 2014 Dec;14(1):1-8.
19. Pollreisz A, Schmidt-Erfurth U. Diabetic cataract—pathogenesis, epidemiology and treatment. *Journal of ophthalmology*. 2010 Oct;2010.
20. Javadi MA, Zarei-Ghanavati S. Cataracts in diabetic patients: a review article. *J Ophthalmic Vis Res*. 2008 Jan; 3(1): 52-65.