

A Prospective Assessment of the Correlation between New Born Size and Maternal First Trimester Haemoglobin Concentration

Soni Kumari¹, Kiran Kumari²

¹Senior resident, Department of Obstetrics and Gynaecology, Madhubani Medical College and Hospital, Madhubani, Bihar, India

²Assistant professor, Department of Obstetrics and Gynaecology, Madhubani Medical College and Hospital, Madhubani, Bihar, India

Received: 10-03-2024 / Revised: 15-04-2024 / Accepted: 24-05-2024

Corresponding Author: Dr. Kiran Kumari

Conflict of interest: Nil

Abstract

Aim: The aim of the present study was to assess the correlation between new born size and maternal first trimester haemoglobin concentration.

Methods: The prospective observational study, conducted in the Department of Obstetrics and Gynaecology from May 2023 to January 2024 included 200 mothers registered for antenatal care (ANC) and willing to continue till confinement were recruited after written informed consent as per routine protocol.

Results: Out of 200 women patients, 55% had normal Hb level followed by mild 32% and 13% moderate. The correlation between hemoglobin concentration and birth weight and other anthropometric parameters was measured by using chi square test and student's t-test. P-value was considered significant if it was below and highly significant in case <0.001.

Conclusion: Increased incidence of low birth weight babies is seen if the mother is anemic in her first trimester with significant association when hemoglobin is less than 8 gm%. Increased incidence of preterm deliveries is seen if the mother is anemic in her first trimester. We observed better neonatal outcome in the form of weight and anthropometry if maternal hemoglobin is in range of 10 to 13 gm%.

Keywords: new born size, maternal first trimester, haemoglobin concentration

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

Introduction

Women are more likely than men to have low iron stores because of blood loss at the time of menstruation. During pregnancy, the fetal demand for iron increases maternal daily iron requirements from ≈ 1 to 2.5 mg/d in early pregnancy and 6.5 mg/d in the third trimester. The average daily diet in the developed world contains $\approx 10-14$ mg non-heme iron¹ but not all of this can be absorbed. Evidence from stable-isotope studies suggests that the percentage of non-heme iron absorbed from food during normal pregnancy increases from 7% at 12 wk of gestation to 36% at 24 wk and 66% at 36 wk. These dramatic changes enable the healthy pregnant woman to cope with the extra demands of pregnancy without becoming anemic [1], but only if there is adequate iron in her diet. If the woman's diet is deficient in iron, as is the case in many developing countries, fetal requirements can be met only by additional contributions of iron from maternal stores. This demand by the developing fetus may cause the mother to develop iron deficiency anemia

if she had inadequate iron stores at the beginning of pregnancy.

Whether a pregnant woman is anemic cannot be assessed simply by measuring the blood hemoglobin concentration because a major factor influencing hemoglobin concentration in pregnancy is expansion of plasma volume. How this occurs is not fully established but part of the sequence might be as follows. Heat production by the feto-placental unit causes a rise in body temperature. Heat loss is increased by peripheral vasodilation, which causes a drop in blood pressure. This in turn stimulates the release of aldosterone from the adrenal gland, causing the retention of salts and water. [2] The drop in osmolality that occurs reduces blood viscosity and enhances blood flow in the low-pressure system of the intervillous space. Enhanced blood flow improves fetal growth. In women who are not given supplemental iron, the hemoglobin concentration of the maternal blood falls from an average of ≈ 133 g/L in non-pregnant women to an average of ≈ 110 g/L at 36 wk of gestation. [3] The fall is steepest up to 20

wk of gestation; the hemoglobin concentration remains fairly constant up to 30 wk and then rises slightly thereafter. [4,5] These changes in hemoglobin concentration are due mainly to changes in plasma volume; the red cell mass and total hemoglobin actually increase during pregnancy.

Maternal nutrition during pregnancy critically determines fetal nutrition and has a significant contribution to fetal and neonatal health. Maternal intakes of micro nutrients such as folic acid and iron are crucial for fetal growth and development. [6] Hemoglobin concentration is a key indicator reflecting the maternal nutrition status during pregnancy, especially the iron status. [7] Previous studies reported that maternal hemoglobin concentration was associated with neonatal birth weight as well as LBW and SGA, but there is no consistent conclusion. Rasmussen et al. found a strong independent inverse correlation between the lowest second-trimester hemoglobin and birth weight, but no relationship was observed in the first trimester. [8] Haider et al. reported that neonatal birth weight increased by 14 g for every 1 g/L increase in average hemoglobin in the third trimester. Steer et al. observed that the maximum mean birth weight was achieved with a lowest hemoglobin concentration in pregnancy of 85–95 g/L, which indicated a nonlinear relationship between maternal hemoglobin during pregnancy and birth weight. [9] A recent meta-analysis revealed a U-shaped curve association between maternal hemoglobin concentration and adverse birth outcomes, which suggested that both low and high hemoglobin concentrations might be risk factors for fetal growth. [10]

The aim of the present study was to assess the correlation between new born size and maternal first trimester haemoglobin concentration.

Materials and Methods

The prospective observational study, conducted in the Department of Obstetrics and Gynaecology, Madhubani Medical College and Hospital, Madhubani, Bihar, India from May 2023 to January 2024, included 200 mothers registered for antenatal care (ANC) and willing to continue till confinement were recruited after written informed consent as per routine protocol. In this study, cases with hemoglobin >11 gm% and <11 gm% were classified as normal and anemia respectively in first trimester of pregnancy.

Table 1: Percentage of severity of anemia in our study as per severity assessment by WHO classification

Haemoglobin level	First Trimester
Normal (>11.1 gm%)	110 (55%)
Mild (9.1-11 gm%)	64 (32%)
Moderate (7.1-9 gm%)	26 (13%)
Severe (4.1-7 gm%)	0
Total	200 (100%)

Inclusion Criteria

1. All pregnant women registered to our institute for ANC till deliveries were included in the Study.
2. Singleton pregnancy
3. Having had a USG in first trimester to accurately confirm / adjust dates and assign gestational age accordingly

Exclusion Criteria

Pregnant women with one of the following at booking were excluded:

1. Diabetes mellitus.
2. Hypertension (including pregnancy-induced hypertension).
3. Toxoplasmosis, Rubella, Cytomegalovirus, Herpes infection.
4. Diagnosed renal or cardiac illness.
5. Smoker or alcoholic.
6. Multiple gestation.
7. hemolytic anemia

The investigations done were

1. Blood-
 - a) Measurement of hemoglobin concentration was done by cyanmethemoglobin method (Analyzer–Coulter).
 - b) Complete blood picture- MCV, MCH, MCHC, RDW.
 - c) Hematocrit (Hct).
 - d) Peripheral smear for typing of anemia.
 - e) Hemoglobin electrophoresis whenever required. Birth weight was recorded in grams using a digital scale with a scale of 1 gram. As per weight of newborns categorization was done as normal if birth weight is above 2.5 kg and low birth weight if less than 2.5 kg.

Statistical Analysis

The correlation between hemoglobin concentration and birth weight and other anthropometric parameters was measured by using chi test and students-t test. P value was considered significant if it was below 0.05 and highly significant in case <0.001.

Observation and Results

Results

Out of 200 women patients, 55% had normal Hb level followed by mild 32% and 13% moderate.

Table 2: First trimester hemoglobin concentration and outcome in the form of birth weight of baby and maturity

Hb%	7.1-9.0gm%	9.1-11gm%	11.1-13gm%	>13gm%
Mean birth weight in gm	8.16±0.46 2242±124.36	11.13±0.17 2782±398.56	11.84±0.52 2864±372.54	13.57±0.74 2656±214.16
Birthwt<2500gms (n=60)	2018±106	2216±144	2282±102.56	2272±2098.52
hwt>2500g (n=140)	3144±1.04	2945±190	2965±276.84	2754.01±152.42
Preterm (n=45)	7	19	15	4
Term (n=155)	10	60	75	10

The correlation between hemoglobin concentration and birth weight and other anthropometric parameters was measured by using chi square test and student's t-test. P-value was considered significant if it was below and highly significant in case <0.001.

Discussion

Every Second women is anemic (55%). Evidence from stable-isotope studies suggest that the percentage of non-heme iron absorbed from food during normal pregnancy increases from 7% at 12 weeks of gestation to 36% at 24 weeks and 66% at 36 wks. These dramatic changes enable the healthy pregnant woman to cope with the extra demands of pregnancy without becoming anemic¹, but only if there is adequate iron in her diet. [11] Prevalence of anemia in South Asian countries is the highest in the world. WHO estimates that even among the South Asian countries, India has the highest prevalence of anemia. [12] NNMB (National Nutrition Monitoring Bureau), DLHS (District Level Household Survey) and ICMR (Indian Council of Medical Research) surveys showed that over 87% of pregnant women suffer from anemia and about 10% have severe anemia. The importance of adequate plasma volume expansion in allowing adequate fetal growth is attested by several studies that showed an increased incidence of low birth weight in association with either a high maternal hemoglobin concentration [13-15] or high hematocrit. [16]

Out of 200 women patients, 55% had normal Hb level followed by mild 32% and 13% moderate. Neonatal birth weight has been obtaining wide attention as it is a strong predictor of neonatal and perinatal mortality and disability, as well as birth weight percentiles, are used to predict the risk of growth disorders in newborns. [17] The global prevalence of low birth weight (LBW) was 14.6% in 2015 [18] and it was estimated that approximately 32.4 million infants that were small for gestational age (SGA) were born in low-income and middle-income countries in 2010. [19] Both LBW and SGA were associated with short-term and long-term adverse outcomes, such as infection, respiratory depression, jaundice, obesity, insulin resistance and type 2 diabetes. [20,21] In addition, the prevalence

of large for gestational age (LGA) and macrosomia is also increasing, especially in developing countries. [22,23] The correlation between hemoglobin concentration and birth weight and other anthropometric parameters was measured by using chi square test and student's t-test. P-value was considered significant if it was below and highly significant in case <0.001.

Maternal Hb concentration in the late-third trimester had a roughly negative association with neonatal birth weight and a faint U-shaped association with Lightweight. When maternal Hb was 100–110 g/L, the risk of Lightweight was the lowest and the risk of Heavyweight was relatively high. Similar to our results, Chen et al. showed that maternal Hb in the third trimester was inversely correlated with neonatal birth weight. [24] Relatively low maternal Hb in the third trimester usually reflects changes in plasma volume rather than poor maternal nutrition or adaptation. [25] However, a recent prospective study from Northwest China showed an inverted U-shaped association between maternal Hb in the third trimester and neonatal birth weight [26] and a significant positive correlation between maternal Hb in the third trimester and birth weight was reported in another study. [27]

In several studies, a U-shaped association was observed between maternal hemoglobin concentrations and birth weight. Abnormally high hemoglobin concentrations usually indicate poor plasma volume expansion, which is also a risk for low birth weight. Lower birth weights in anaemic women have been reported in several studies. [28,29] In our study, in first trimester clustering of normal birth weight was observed in hemoglobin (Hb) range of 9.0 to 13.0gm% and percentage of LBW was increased as hemoglobin drops below 9.0 gm% and also when hemoglobin is above 14 gm%. We also observed, high hemoglobin percentage above 14 gm% was not positively associated with proportionate increase in newborn size as noted with U shape correlation of maternal hemoglobin with newborn size and gestational age.

Conclusion

Increased incidence of low-birth-weight babies is seen if the mother is anemic in her first trimester with significant association when hemoglobin is less than 8 gm%. Increased incidence of preterm deliveries is seen if the mother is anemic in her first trimester. We observed better neonatal outcome in the form of weight and anthropometry if maternal hemoglobin is in range of 10 to 13 gm%. Supplementing iron earlier during antenatal period and maintaining optimal hemoglobin concentration between 10 to 13 gm percent has overall better outcome regarding premature deliveries and low birth weight babies. Regular antenatal care from first trimester has a vital role in assessing and managing maternal anemia timely and it directly affects the perinatal outcomes.

References

- Barrett JF, Whittaker PG, Williams JG, Lind T. Absorption of non-haem iron from food during normal pregnancy. *Bmj*. 1994 Jul 9;309(6947):79-82.
- Chapman AB, Zamudio S, Woodmansee W, Merouani A, Osorio F, Johnson A, Moore LG, Dahms T, Coffin C, Abraham WT, Schrier RW. Systemic and renal hemodynamic changes in the luteal phase of the menstrual cycle mimic early pregnancy. *American Journal of Physiology-Renal Physiology*. 1997 Nov 1;273(5):F777-82.
- Taylor DJ, Lind T. Red cell mass during and after normal pregnancy. *BJOG: An International Journal of Obstetrics & Gynaecology*. 1979 May;86(5):364-70.
- Paintin DB, Thomson AM, Hytten FE. Iron and the haemoglobin level in pregnancy.
- Lu ZM, GOLDENBERG RL, CLIVER SP, CUTTER G, BLANKSON M. The relationship between maternal hematocrit and pregnancy outcome. *Obstetrics & Gynecology*. 1991 Feb 1;77(2):190-4.
- Mousa A, Naqash A, Lim S. Macronutrient and Micronutrient Intake during Pregnancy: An Overview of Recent Evidence. *Nutrients*. 2019 Feb 20;11(2):443.
- Means RT. Iron Deficiency and Iron Deficiency Anemia: Implications and Impact in Pregnancy, Fetal Development, and Early Childhood Parameters. *Nutrients*. 2020 Feb 11;12(2):447.
- Rasmussen S, Oian P. First- and second-trimester hemoglobin levels. Relation to birth weight and gestational age. *Acta Obstet Gynecol Scand*. 1993 May;72(4):246-51.
- Steer P, Alam MA, Wadsworth J, Welch A. Relation between maternal haemoglobin concentration and birth weight in different ethnic groups. *BMJ*. 1995 Feb 25;310(6978):489-91.
- Dewey KG, Oaks BM. U-shaped curve for risk associated with maternal hemoglobin, iron status, or iron supplementation. *Am J Clin Nutr*. 2017 Dec;106(Suppl 6):1694S-1702S.
- Rasmussen KM. Is there a causal relationship between iron deficiency or iron-deficiency anemia and weight at birth, length of gestation and perinatal mortality?. *The Journal of nutrition*. 2001 Feb 1;131(2):590S-603S.
- Alwan NA, Greenwood DC, Simpson NA, McArdle HJ, Godfrey KM, Cade JE. Dietary iron intake during early pregnancy and birth outcomes in a cohort of British women. *Human Reproduction*. 2011 Apr 1;26(4):911-9
- Ahmad MO, Kalsoom U, Sughra U, Hadi U, Imran M. Effect of maternal anaemia on birth weight. *Journal of Ayub Medical College Abbottabad*. 2011 Mar 1;23(1):77-9.
- Bakhtiar UJ, Khan Y, Nasar R. Relationship between maternal hemoglobin and perinatal outcome. *Age*. 2007;25:24.
- NASIRI AF, HAJI AM, Basirat Z. Maternal hematocrit status affecting pregnancy outcome.
- Levy A, Fraser D, Katz M, Mazor M, Sheiner E. Maternal anemia during pregnancy is an independent risk factor for low birthweight and preterm delivery. *European journal of obstetrics & gynecology and reproductive biology*. 2005 Oct 1;122(2):182-6.
- Goldenberg RL, Culhane JF. Low birth weight in the United States. *Am J Clin Nutr*. 2007 Feb;85(2):584S-590S.
- Blencowe H, Krusevec J, de Onis M, Black RE, An X, Stevens GA, Borghi E, Hayashi C, Estevez D, Cegolon L, Shiekh S, Ponce Hardy V, Lawn JE, Cousens S. National, regional, and worldwide estimates of low birthweight in 2015, with trends from 2000: a systematic analysis. *Lancet Glob Health*. 2019 Jul;7(7): e849-e860.
- Lee AC, Katz J, Blencowe H, Cousens S, Kozuki N, Vogel JP, Adair L, Baqui AH, Bhutta ZA, Caulfield LE, Christian P, Clarke SE, Ezzati M, Fawzi W, Gonzalez R, Huybregts L, Kariuki S, Kolsteren P, Lusingu J, Marchant T, Merialdi M, Mongkolchat A, Mullany LC, Ndirangu J, Newell ML, Nien JK, Osrin D, Roberfroid D, Rosen HE, Sania A, Silveira MF, Tielsch J, Vaidya A, Willey BA, Lawn JE, Black RE; CHERG SGA-Preterm Birth Working Group. National and regional estimates of term and preterm babies born small for gestational age in 138 low-income and middle-income countries in 2010. *Lancet Glob Health*. 2013 Jul;1(1):e26-36.
- Lee AC, Kozuki N, Cousens S, Stevens GA, Blencowe H, Silveira MF, Sania A, Rosen HE, Schmiegelow C, Adair LS, Baqui AH, Barros FC, Bhutta ZA, Caulfield LE, Christian P, Clarke SE, Fawzi W, Gonzalez R, Humphrey J, Huybregts L, Kariuki S, Kolsteren P, Lusingu J, Manandhar D, Mongkolchat A, Mullany LC,

- Ndyomugenyi R, Nien JK, Roberfroid D, Saville N, Terlouw DJ, Tielsch JM, Victora CG, Velaphi SC, Watson-Jones D, Willey BA, Ezzati M, Lawn JE, Black RE, Katz J; CHERG Small-for-Gestational-Age-Preterm Birth Working Group. Estimates of burden and consequences of infants born small for gestational age in low and middle income countries with INTERGROWTH-21st standard : analysis of CHERG datasets. *BMJ*. 2017 Aug 17;358;j3677.
21. Barker DJ. The developmental origins of chronic adult disease. *Acta Paediatr Suppl*. 2004 Dec;93(446):26-33.
 22. Surkan PJ, Hsieh CC, Johansson AL, Dickman PW, Cnattingius S. Reasons for increasing trends in large for gestational age births. *Obstet Gynecol*. 2004 Oct;104(4):720-6.
 23. Li G, Kong L, Li Z, Zhang L, Fan L, Zou L, Chen Y, Ruan Y, Wang X, Zhang W. Prevalence of macrosomia and its risk factors in china: a multicentre survey based on birth data involving 101,723 singleton term infants. *Paediatr Perinat Epidemiol*. 2014 Jul;28(4): 34 5-50.
 24. Chen JH, Guo XF, Liu S, Long JH, Zhang GQ, Huang MC, Qiu XQ. [Impact and changes of maternal hemoglobin on birth weight in pregnant women of Zhuang Nationality, in Guangxi]. *Zhonghua Liu Xing Bing Xue Za Zhi*. 2017 Feb 10;38(2):154-157.
 25. Whittaker PG, Macphail S, Lind T. Serial hematologic changes and pregnancy outcome. *Obstet Gynecol*. 1996 Jul;88(1):33-9.
 26. Liu D, Li S, Zhang B, Kang Y, Cheng Y, Zeng L, Chen F, Mi B, Qu P, Zhao D, Zhu Z, Yan H, Wang D, Dang S. Maternal Hemoglobin Concentrations and Birth Weight, Low Birth Weight (LBW), and Small for Gestational Age (SGA): Findings from a Prospective Study in Northwest China. *Nutrients*. 2022 Feb 18;14 (4):858.
 27. Yildiz Y, Özgü E, Unlu SB, Salman B, Eyi EG. The relationship between third trimester maternal hemoglobin and birth weight/length; results from the tertiary center in Turkey. *J Matern Fetal Neonatal Med*. 2014 May;27(7): 729-32.
 28. Ezzati M, Lopez AD, Rodgers A, Vander Hoorn S, Murray CJ. Selected major risk factors and global and regional burden of disease. *the lancet*. 2002 Nov 2;360(9343): 13 47-60.
 29. Hemminki E, Rimpelä U. Iron supplementation, maternal packed cell volume, and fetal growth. *Archives of Disease in Childhood*. 1991 Apr 1;66(4 Spec No):422-5.