

A Clinical Trial to Conduct to Compare the Effectiveness and Stability of Ketamine/Fentanyl in Combination with Propofol in Patients Undergoing Dilatation and Curettage

Sujeet Kumar¹, Soniya Kumari², Muni Lal Gupta³

¹Senior Resident, Department of Anaesthesia, BMIMS Pawapuri, Nalanda, Bihar, India

²PG Student, Physiology, BMIMS Pawapuri, Nalanda, Bihar, India

³Assistant Professor and HOD, Department of Anaesthesia, BMIMS, Pawapuri, Nalanda, Bihar, India

Received 08-03-2024 / Revised: 14-04-2024 / Accepted: 27-05-2024

Corresponding Author: Dr. Soniya Kumari

Conflict of interest: Nil

Abstract

Aim: The aim of the present study was to evaluate and compare the efficacy and hemodynamic stability of ketamine / fentanyl along with propofol and also assess recovery profile, post-operative analgesia and complications in patients undergoing Dilatation and Curettage.

Methods: The present study was carried out in the Department of Anaesthesia, BMIMS Pawapuri, Nalanda, Bihar, India. the study was conducted in 50 patients, aged 18-35 years of ASA Grade 1&2 scheduled for Dilatation and Curettage procedures lasting up to 30 minutes.

Results: Pre-induction heart rate, systolic blood pressure, diastolic blood pressure, mean arterial pressure, spo2 were comparable in both the groups with a statistically no significant difference between them ($p < 0.05$).

Conclusion: It may be concluded from our present study that propofol with ketamine as an adjuvant in the dose of 0.75 mg/kg compared to propofol with fentanyl as adjuvant in the dose of 2 mcg/kg provides deep sedation. But propofol-ketamine group is more efficacious and provides better peri-operative hemodynamic stability during anaesthesia as compared to propofol-fentanyl group. Also the propofol-ketamine combination produces good analgesia with less requirement of rescue drug in post-operative period with fewer peri-operative complications than propofol-fentanyl combination. We have not encountered any psychotomimetic effects of injection Ketamine in the 0.75mg/kg dose. Hence propofol-ketamine combination is a better choice especially when hemodynamic stability is of great importance in patients undergoing dilatation and curettage.

Keywords: Propofol, Ketamine, Fentanyl, Sedation, Hemodynamic stability, Dilatation, Curettage

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

Introduction

Dilatation and Curettage (DandC), is a short lasting but painful surgical procedure, performed for the diagnosis and treatment of endometrial and intrauterine disorders. It causes significant pain due to cervical dilatation with dilators and tissue extraction. [1] Therefore, drugs that are used for this procedure, should ensure a rapid onset of action, an adequate level of sedation, good analgesia as well as rapid recovery. The most important objectives in the perioperative period is being maintenance of the hemodynamic-respiratory stability and minimizing the side effects of the drug. Propofol is a widely used sedative agent due to its rapid onset of action and fast recovery time, but it causes cardiovascular and respiratory depression in a dose dependent manner. Also it might be insufficient in painful processes, if used alone since it lacks analgesic properties. [1]

Combining propofol with other drugs such as opioids or ketamine is recommended for improving the quality of sedation, analgesia and minimizing the potential adverse effects with maintenance of a stable cardiovascular and respiratory status in the perioperative period. [2] Ketamine, a NMDA receptor agonist, in subanesthetic doses with propofol has gained attention in total intravenous anesthesia because of its powerful analgesic action without causing myocardial and respiratory depression. Ketamine also causes some degree of sympathetic stimulation, which tends to counter balance the cardiovascular effects of propofol. [3] Fentanyl on the other hand is synthetic opioid analgesic which has rapid onset and short duration of action and has been used in combination with propofol satisfactorily.

Ketofol (propofol-ketamine admixture) is a combination of ketamine and propofol that is an agent of choice for various procedures. [4] The safety and efficacy of ketofol as a sedoanalgesic agent depend on the dose and the ratio of the admixture. [5] The ratios of 1:2, 1:3 and 1:4 [sub dissociative dose] ratios were very effective for the day case procedure. [6] Dilation and curettage (D C), a brief and painful procedure, is performed for the diagnosis and treatment of endometrial and intrauterine disorders. The procedure is one of the most frequently performed gynecological surgical procedures. It causes considerable pain during cervical dilation and tissue extraction. [7]

The aim of the present study was to evaluate and compare the efficacy and hemodynamic stability of ketamine / fentanyl along with propofol and also assess recovery profile, post-operative analgesia and complications in patients undergoing Dilatation and Curettage.

Materials and Methods

The present study was carried out in the Department of Anaesthesia, BMIMS Pawapuri, Nalanda, Bihar, India for one year the study was conducted in 50 patients, aged 18-35 years of ASA Grade 1&2 scheduled for Dilatation and Curettage procedures lasting up to 30 minutes. The patients were randomly allocated using sealed envelope method in 2 groups (25 of each). Patients with respiratory, cardiovascular, neurological, liver diseases, patients on narcotic therapy were excluded from study. Pre-anaesthetic evaluation including history and a thorough general and systemic examination and all relevant investigations were done for all the patients.

Patients were kept fasting for at-least 8 hours prior to anaesthesia. Pre-operative baseline heart rate, systolic blood pressure, diastolic blood pressure, mean blood pressure, spo₂ was recorded. A peripheral iv line was established. All patients were

pre-medicated with Inj. Glycopyrrolate 0.004 mg/kg and Inj. Midazolam 0.02 mg/kg. Group A received Inj. Ketamine 0.75 mg/kg body weight IV slowly over 2 minutes, after 5 minutes Inj. Propofol given at rate of 1ml/3 seconds till loss of consciousness. Group B received Inj. Fentanyl 2 mcg/kg body weight as slow IV injection, after 5 minutes Inj. Propofol given at rate of 1ml/3 seconds till loss of consciousness.

Intra-operatively heart rate, blood pressure, oxygen saturation and Ramsay sedation score was recorded at different time intervals of 5, 10, 15, 20, 25, 30 minutes following induction of anaesthesia in both groups. Throughout the procedure patients were allowed to breathe spontaneously on room air and oxygen supplementation was given to some patients during apnoea.

Top-up dose of Inj. Propofol 0.5 mg/kg was given when patient became light during anaesthesia as indicated by rise in heart rate, blood pressure or any other movement to surgical stimulus. Total dose of propofol required for the patients was noted.

Post-operatively all vital parameters like heart rate, blood pressure, oxygen saturation and visual analogue score was recorded every 30 minutes for first 2 hours then every 2 hours till 12 hours. Any complication like nausea, vomiting, delirium, sedation, pain was noted.

Statistical Analysis

Continuous variables like age, heart rate, systolic blood pressure, diastolic blood pressure, Ramsay sedation score, spo₂, total dose of propofol etc were presented as Mean \pm SD. Continuous variables were compared at different time intervals between ketamine and fentanyl groups by performing unpaired t-test, $p < 0.05$ was considered as statistically significant.

Results

Table 1: Comparison of Heart Rate in Both Groups

Heart rate	Baseline	1 min	5 min	10 min	15 min	20 min	25 min	30 min
Group A Mean \pm SD	84.56 \pm 3.34	81.20 \pm 4.93	87.62 \pm 5.2	83.30 \pm 6.33	83.0 \pm 5.58	80.45 \pm 6.02	80.22 \pm 6.2	80.7 \pm 6.1
Group B Mean \pm SD	86.86 \pm 3.86	88.05 \pm 6.04	82.66 \pm 6.00	77.20 \pm 5.27	78.34 \pm 4.85	76.45 \pm 6.63	80.65 \pm 6.16	79.55 \pm 6.31
P- value	0.01	0.0001	0.0005	0.0003	0.0007	0.03	0.741	0.497

Table 2: Comparison of Systolic Blood Pressure (SBP) In Both Groups

SBP	Baseline	1 min	5 min	10 min	15 min	20 min	25 min	30 min
Group A Mean \pm SD	123.64 \pm 5.48	119.14 \pm 12.17	120.60 \pm 11.30	119.14 \pm 9.90	116.35 \pm 10.97	114.10 \pm 11.87	114.78 \pm 10.56	115.07 \pm 10.23
Group B Mean \pm SD	123.35 \pm 4.52	119.85 \pm 9.67	117.21 \pm 8.35	116.57 \pm 9.19	111 \pm 7.44	108.42 \pm 8.21	109.85 \pm 8.68	107 \pm 18.77
P-value	0.823	0.803	0.191	0.301	0.030	0.04	0.05	0.001

Table 3: Comparison of Diastolic Blood Pressure (DBP) In Both Groups

DBP	Base line	1 min	5 min	10 min	15 min	20 min	25 min	30 min
Group A Mean ± SD	72.47 ±4.91	69.65± 7.04	69.38± 2.95	67.42± 5.56	67.22± 5.34	67.35± 5.97	67.47± 4.39	69.25± 3.78
Group B Mean ± SD	73.37 ±4.51	69.77± 4.88	69.47± 3.51	64.75± 3.59	64.45± 3.59	65.58± 4.71	67.37± 4.39	25
P- value	0.370	0.858	0.802	0.013	0.017	0.060	0.033	0.926

Table 4: Comparison of Ramsay Sedation Score In Both Groups

RSS	Baseline	1 min	5 min	10 min	15 min	20 min	25 min	30 min
Group A Mean ± SD	1.92 ±0.27	1.92±0 .27	4.40±0 .57	4.82±0 .56	5.46±0 .57	5.86±0 .35	5.96±0 .19	5.95±0 .20
Group B Mean ± SD	1.94± 0.23	1.94±0 .23	3.94±0 .74	4.06±0 .72	5.10±0 .57	5.72±0 .49	5.9±0. 30	5.97±0 .14
P- value	0.70	0.70	0.009	<0.00 01	<0.00 01	0.20	0.35	0.65

Table 5: Comparison of VAS in Both Groups

VAS	30 min	1 hour	1 hour 30 min	2 hour	4 hour	6 hour	8 hour	10 hour	12 hour
Group A Mean ± SD	2.87 ±1.2 2	2.33± 1.12	1.57±0 .9	1.33± 0.76	1.17± 0.53	1.1±0.4	1±0	1±0	1±0
Group B Mean ± SD	4.37 ±1.5 4	3.27± 1.34	2.8±1. 13	2.3±0 .84	1.63± 0.85	1.23±0. 5	1±0	1±0	1±0
P -value	0.00 01	0.004	<0.000 1	<0.00 01	0.013	0.262	-	-	-

Pre-induction heart rate, systolic blood pressure, diastolic blood pressure, ramsay sedation score, VAS were comparable in both the groups with a statistically no significant difference between them (p<0.05).

Discussion

Day care gynecological procedures require the use of anesthetic agents which ensure rapid induction and recovery. [8] Total intravenous anesthesia (TIVA) is a combination of hypnotic agents, analgesic drugs and may be muscle relaxants, excluding simultaneous administration of any inhaled drugs. Therefore, it can be an effective alternative to inhalational anesthesia and for ambulatory surgery when the speed and completeness of recovery are important. Drugs used for TIVA should have quick onset, smooth induction, easy maintenance, quick recovery and minimal side effects. [4] Ideal drug for sedoanalgesia should have rapid onset and fast recovery time. However, there is still no consensus for best sedoanalgesic management for short-term procedures. [7]

Pre-induction heart rate, systolic blood pressure, diastolic blood pressure, ramsay sedation score, VAS were comparable in both the groups with a statistically no significant difference between them (p<0.05). The minimal fall in SBP, DBP and MBP

in group PK as compared to group PF is due to the sympathomimetic activity of ketamine which counteracts with the cardiovascular depressant action of propofol, thus maintaining a stable hemodynamic profile as compared to fentanyl. These findings are correlated well with the study done by Pawar et al [9], Phillips et al [10] and Khutia et al. [11] The opposing effect of ketamine and propofol on arterial pressure tended to cancel each other out resulting in improved cardiovascular stability. The propofol ketamine group experienced a smaller percentage decrease in SBP, which is comparable with other studies. [2,3,12]

The ketamine-propofol combination is thought to act by counteracting the cardiovascular side-effects of each other, preserving the sedative efficacy. Also the amount of propofol needed to achieve a deep sedation level was much lower in case of PK group than PF group, which contributed to the lower incidence of hypotension and apnea. Also, no patients in present study required airway manipulation. The post operative complications were also higher in group PF than in group PK. There were no incidences of psychedelic effects of ketamine like hallucination, dysphoria and psychotomimetic emergence reactions when ketamine is used in combination with a sedative-hypnotic (e.g., benzodiazepines, propofol) or a general anesthetic (e.g., halothane, nitrous oxide) is

minimal or negligible. Hypnotic doses of propofol are reported to block these hallucinations. These findings are concordance with the study done by Aarikan et al [2] Brajesh et al [3] Mahajan et al. [12]

Conclusion

It may be concluded from our present study that propofol with ketamine as an adjuvant in the dose of 0.75 mg/kg compared to propofol with fentanyl as adjuvant in the dose of 2 mcg/kg provides deep sedation. But propofol-ketamine group is more efficacious and provides better peri-operative hemodynamic stability during anaesthesia as compared to propofol-fentanyl group. Also the propofol-ketamine combination produces good analgesia with less requirement of rescue drug in post-operative period with fewer peri-operative complications than propofol-fentanyl combination. We have not encountered any psychotomimetic effects of injection Ketamine in the 0.75mg/kg dose. Hence propofol-ketamine combination is a better choice especially when hemodynamic stability is of great importance in patients undergoing dilatation and curettage.

References

1. Yuce H, Kucuk A, Altay N, Bilgic TE, Karahan M, Buyukfirat E, Havlioglu I, Yalcin S, Camuzcuoglu A, Incebiyik A, Hilali G. Propofol-ketamine combination has favorable impact on orientation times and pain scores compared to propofol in dilatation and curettage. A randomized trial. *Acta Medical Mediterranean*. 2013 Jan 1;29:539-44.
2. Aarikan M, Aslan B, Aarikan O, But A, Horasanli EY. Comparison of propofol-remifentanyl and propofol-ketamine combination for dilatation and curettage: a randomized double blind prospective trial. *European Review for Medical and Pharmaceutical Sciences*. 2015 Sep 1;19(18):3522-7.
3. Kaushal B, Dhawale Y, Gupta D. Comparative evaluation of propofol ketamine and propofol fentanyl in day care surgeries. *Journal of evolution of Medical and Dental Sciences*. 2014 Sep 4;3(41):10325-36.
4. Babita R, Dinesh K, Sangita R, Gurdeep S. A comparative evaluation of Propofol-Ketamine and Propofol-fentanyl as TIVA techniques in terms of haemodynamic variables and recovery characteristics in minor surgeries. *IOSR Journal of Dental and Medical Sciences (IOSRJDMS)*. 2015;14(4):19-26.
5. Daabiss M, Medhat E and Rashed A. (2009): Assessment of different concentration of ketofol in procedural operation. *British Journal of Medical Practitioners*, 2(1)27-31.
6. Wang Y, Jiang X, Pang L, Dong S, Feng Y, Prajapati SS, Yuan C, Ma H. A randomized double-blind controlled study of the efficacy of ketofol with propofol-fentanyl and propofol alone in termination of pregnancy. *Afr J pharm pharmacol*. 2012 Sep 15;6(34):2510-4.
7. Yuce H, Kucuk A, Altay N, Bilgic TE, Karahan M, Buyukfirat E, Havlioglu I, Yalcin S, Camuzcuoglu A, Incebiyik A, Hilali G. Propofol-ketamine combination has favorable impact on orientation times and pain scores compared to propofol in dilatation and curettage. A randomized trial. *Acta Medical Mediterranean*. 2013 Jan 1;29:539-44.
8. Ahuja H, Abraham V, Abraham J, Liddle D. Ideal anesthetic agents for day-care gynecological procedures: A clinical trial comparing thiopentone with ketamine as adjuncts to propofol. *Advanced Biomedical Research*. 2015 Jan 1;4(1):81.
9. Pawar D, Bhople P, Pandey S, Khobragade S. Comparative evaluation of propofol- ketamine and propofol- fentanyl for minor surgical procedures. *International Journal of Research in Medical Sciences*. 2015; 3(12): 3795-3801.
10. Phillips W, Anderson A, Rosengreen M, Johnson J, Halpin J. Propofol versus propofol/ketamine for brief painful procedures in the emergency department: clinical and bispectral index scale comparison. *J Pain Palliat Care Pharmacother* 2010; 24: 349-355.
11. Khutia SK, Mandal MC, Das S, Basu SR. Intravenous infusion of ketamine-propofol can be an alternative to intravenous infusion of fentanyl-propofol for deep sedation and analgesia in paediatric patients undergoing emergency short surgical procedures. *Indian J Anaesth* 2012;56:145-50.
12. Mahajan R, Swarnkar N, Ghosh A. Comparison of ketamine and fentanyl with propofol in total intravenous anesthesia: a double blind randomized clinical trial. *The Internet Journal of Anesthesiology*. 2009;23 (2):1-7.