

## Assessing Dietary Pattern and Physical Activity among Adolescents: An Observational Study

Kanchan Lata<sup>1</sup>, Chikirsha Vijay<sup>2</sup>

<sup>1</sup>Associate Professor, Department of Community Medicine, Mata Gujri Memorial Medical College, Kishanganj, Bihar, India

<sup>2</sup>Senior Resident, Department of Pediatrics, Darbangha Medical College and Hospital, Darbhanga, Bihar, India

---

Received: 25-03-2024 / Revised: 11-04-2024 / Accepted: 10-05-2024

Corresponding Author: Dr. Chikirsha Vijay

Conflict of interest: Nil

---

### Abstract

**Aim:** The current study aimed to focus specifically on objectively assessed physical activity & and dietary habits among adolescents of BIHAR.

**Methods:** The present study was conducted at department of community medicine and cross-sectional data were collected. Data were obtained from adolescents (11-16 years) standard selected recruited from three secondary schools in Bihar. Staff at participating schools selected a subset of their classes for participation. All students from nominated classes (n = 415) were eligible and received written information on the project. Consent was sought from parents prior to the study and adolescent participants provided assent before completing written surveys during class.

**Results:** In total, 393 pupils provided consent and completed the questionnaire (95.85% response rate). Of these, 150 (38.16%) provided usable accelerometer data. Comparison of these 150 adolescents with those who did not provide usable accelerometer data (n = 150) showed no significant differences in age, SES, ethnicity or fruit and vegetable consumption. However, a significantly ( $p < 0.01$ ) higher proportion of those with usable accelerometer data were girls, compared to boys (53% compared to 37%), and ate breakfast on more days per week (4.7 days compared to 3.5 days). Thus, our figures likely underestimate the prevalence of not meeting the breakfast recommendations. The final sample composition was 150 adolescents, with 70 boys, 80 girls, 68 younger adolescents and 50 older adolescents. The mean age of younger adolescents was 14.3 years and of older adolescents was 14.6 years.

**Conclusion:** Many adolescents fail to meet multiple diet and physical activity recommendations, highlighting that physical activity and dietary behaviours do not occur in isolation. Future research should investigate how best to achieve multiple health behaviour change in adolescent boys and girls.

**Keywords:** adolescence, life style, non-communicable disease, physical activity

---

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

---

### Introduction

The word “adolescence” is derived from the Latin word “adolescere,” which means “grow to maturity.” Adolescence is a grey area in the spectrum of life falling between childhood and adulthood. It is an age of transition when an individual experiences rapid growth and development, both physical and psychological and changes from being a child to an adult. [1] In India, adolescents constitute 21.4% of the population, i.e., one-fifth of the total population. [2] The development of healthy eating habits is important as the rapid physical growth in adolescence is associated with increased nutritional needs. Various studies on diet and nutrition intake of adolescents and young adults in the developed world have shown that their diets are often high in fats and

refined carbohydrate. [3] Consumption of diet high in sugar, saturated fat, salt, and calorie content in children can lead to early development of obesity, hypertension, dyslipidemia, and impaired glucose tolerance. [4]

The changing lifestyles and the easy availability of junk food near schools or even within school canteens have led children to prefer and become addicted to junk foods. There is compelling evidence that dietary habits and lifestyle during adolescence are risk factors for several nutrition-related noncommunicable diseases (NCDs) in adulthood. [5] Physical activity is defined as any bodily movement produced by skeletal muscles that result in energy expenditure above basal level and

regular engagement in physical activity is widely acknowledged as having important health benefits. [6,7]

Poor diet and physical inactivity are established risk factors for chronic disease. In young people, physical activity and healthy diets including regular breakfast consumption and adequate levels of fruit and vegetables, have important short- and long-term health protective effects. For example, physical activity in young people may benefit cardiovascular disease (CVD) risk factors, adiposity and bone health, which could influence health in adulthood. [8]

Child fruit and vegetable consumption has been associated with a lower incidence of respiratory symptoms [9] and appears to be protective against cancer in adulthood. [10] Also, cross-sectional and longitudinal research has shown that young people who regularly eat breakfast are less likely to be overweight than those who skip breakfast. [11,12] Despite such health benefits, young people are more likely to skip breakfast than any other meal [13] and data from the Health Behaviour in School-aged Children (HBSC) study shows that less than two-fifths of young people eat fruit daily, and only about a third eat vegetables daily. [14]

The current study aimed to focus specifically on objectively assessed physical activity & and dietary habits among adolescents of BIHAR.

### Materials & Methods

The present study was conducted at department of community medicine, Mata Gujri Medical College, Kishanganj, Bihar, India for 12 months and cross-sectional data were collected. Data were obtained from adolescents (12-16 years) recruited from three secondary schools in Bihar. Staff at participating schools selected a subset of their classes for participation. All students from nominated classes (n = 415) were eligible and received written information on the project. Consent was sought from parents prior to the study and adolescent participants provided assent before completing written surveys during class. Participants completed questionnaires during Physical Education or Personal, Social and Health Education (PHSE) lessons, under the supervision of trained researchers and class teachers.

### Measures

Demographic information, including date of birth and ethnicity, was provided by the school. Questionnaires collected information on demographic characteristics of adolescents including gender and home postcodes. Socioeconomic status (SES) was determined using the Index of Multiple Deprivation (IMD), a measure of compound social and material deprivation, calculated from a variety of data including income,

employment, health, education, and housing. It is based on the postcode of the participant's home, and thus represents an area level approximation of SES.

### Adolescent Dietary Behaviour

Adolescent food intake was assessed using a 30-item food frequency questionnaire (FFQ), based on the previously validated Youth/Adolescent Food Frequency questionnaire (YAQ). [15] Adolescents were asked how often they ate ten different fruits and twelve different vegetables in the past month. Responses to questions on the frequency of consumption of specific fruits and vegetables were summed to compute total frequency of fruit and vegetable consumption/day, respectively. For the purpose of this study each item was summed to calculate the frequency of consumption of 'fruits and vegetables' (FV) per day. Guided by the current recommendations for fruit and vegetable consumption (5 portions of FV/day), the total daily frequency of FV consumption was dichotomised into <5 times/day or > 5 times/day.

Breakfast consumption was assessed with a single-item asking adolescents how often they ate breakfast in the past seven days. While there are no current national recommendations for frequency of breakfast consumption, evidence suggests that young people who eat breakfast on most days of the week see health benefits (e.g. lower Body Mass Index) compared to those who skip breakfast. [16,17] A recent government initiative in the Bihar recommends that parents encourage their children to eat breakfast regularly. Given such evidence, frequency of breakfast consumption/week was dichotomised into <5 days/week or > 5 days/week.

### Physical Activity

Physical activity was assessed by Actigraph GT1M accelerometers (ActiGraph, Fort Walton Beach, FL) using a 5 second measurement interval (epoch). Participants were instructed to wear the accelerometer over their right hip for one-week. Exceptions included time spent sleeping, showering and during water-based activities. Duration of MVPA was computed only for adolescents who wore the accelerometer for a minimum of three days [18], defined as days on which accelerometer counts were between 10,000 and 20 million. [20] To estimate the time spent per day in moderate intensity physical activity (3.0-5.9 metabolic equivalent of rest [METs]) and vigorous intensity physical activity (6.0+ METs), age specific movement count thresholds were applied. Time-per-day in MVPA was derived by summing these values across valid days. The proportion of adolescents meeting the physical activity recommendations for young people was calculated according to whether they performed an average of > 60 mins/day of MVPA. Time spent in MVPA was dichotomised as <60 mins/day or > 60 mins/day

### Statistical Analyses

All analyses were conducted using SPSS statistical software version 16.0. Descriptive statistics were used to summarise sociodemographic, physical activity and dietary characteristics of the sample. Adolescents were categorised as younger or older adolescents by dichotomising at the mean (14.4 years). Mann-Whitney tests were performed to examine gender and age-group differences in mean minutes/day spent in MVPA, fruit and vegetable consumption per day, and breakfast consumption per week. The proportion of adolescents achieving > 60 mins/day of MVPA, and the proportion of adolescents consuming fruit and vegetables > 5 times/day, and breakfast > 5 times/ week were

compared by gender and age-group using Pearson's chi-square ( $\chi^2$ ) tests of significance. A total risk behaviour score was calculated for each participant based on the total number of unmet health recommendations (range from zero to three). Pearson's chi-square tests were used to examine gender and age-group differences in the number of risk behaviours. The proportion of adolescents in each multiple risk behaviour combination was determined to examine behaviour clustering patterns, and Pearson's chi-square tests were used to examine gender and age- group differences in the clustering patterns.

### Results

**Table 1: Distribution fruit and vegetables consumption per day, breakfast consumption per week and minutes per day spent in moderate-to-vigorous intensity physical activity (MVPA)**

	Total (n = 150)	Boys (n = 70)	Girls (n = 80)	Younger adolescents(n = 68)	Older adolescents (n = 50)
<b>MVPA minutes/day</b>					
25th percentile	18.6	23.7	22.5	38.2	24.6
50th percentile	33.6	38.2	38.4	48.2	38.3
75 th percentile	69.0	54.4	49.2	65.2	49.6
<b>Frequency of FV consumption/day</b>					
25th percentile	2.8	1.7	3.6	2.6	3.7
50th percentile	4.4	3.8	4.3	3.9	4.9
75th percentile	7.3	6.7	5.7	6.2	8.2

In total, 393 pupils provided consent and completed the questionnaire (95.85% response rate). Of these, 150 (38.16%) provided usable accelerometer data. Comparison of these 150 adolescents with those who did not provide usable accelerometer data (n = 150) showed no significant differences in age, SES, ethnicity or fruit and vegetable consumption. However, a significantly ( $p < 0.01$ ) higher proportion of those with usable accelerometer data were girls, compared to boys (53% compared to

37%), and ate breakfast on more days per week (4.7 days compared to 3.5 days). Thus, our figures likely underestimate the prevalence of not meeting the breakfast recommendations. The final sample composition was 150 adolescents, with 70 boys, 80 girls, 68 younger adolescents and 50 older adolescents. The mean age of younger adolescents was 14.3 years and of older adolescents was 14.6 years.

**Table 2: Gender and age distribution of meeting health recommendations**

	Total (n = 150)	Boys (n = 70)	Girls (n = 80)	Younger adolescents (n= 68)	Older adolescents N = 50)
<b>Meet &gt;60 minutes MVPA per day, n (%)</b>					
No	115 (76.66)	49 (70)	66 (82.5)	42 (69.4)	44 (88)
Yes	35 (23.34)	21 (30)	14 (17.5)	26 (30.6)	6 (12)
<b>Meet &gt;5 portions fruits/vegetables, n (%)</b>					
No	80 (53.34)	37 (52.85)	43 (53.75)	44(64.3)	22(44)
Yes	70 (46.66)	33 (47.15)	37 (46.25)	24 (35.7)	28 (56)
<b>Meet &gt;5 days a week eating breakfast, n (%)</b>					
No	36 (24)	12 (17.14)	24 (30)	16 (23.5)	12 (24)
Yes	114(76)	58 (82.85)	56 (70)	52 (76.5)	38 (76)
<b>Number of risk behaviours, n</b>					

(%)					
0	10 (6.6)	7 (10)	3 (3.75)	5 (7.1)	3 (6)
1	60 (40)	32 (45.71)	28 (35)	25 (36.7)	22 (44)
2	64 (42.66)	26 (37.14)	38 (47.5)	33 (48.0)	19 (38)
3	16 (10.66)	5 (7.14)	11 (13.75)	5 (8.2)	6 (12)

### Prevalence of meeting health recommendations

A higher proportion of boys, compared to girls, ate breakfast on more than 5 days/week ( $p < 0.01$ ). A higher proportion of younger adolescents, compared to older adolescents, met the physical activity recommendations and did not meet the

recommendations for fruit and vegetable consumption ( $p < 0.01$ ). Only 6% of the adolescent sample met the recommendations for all three of the health behaviours and almost 10% had three health risk behaviours. A higher proportion of girls, compared to boys, had three health risk behaviours ( $p < 0.01$ ).

**Table 3: Descriptive cluster pattern of multiple risk behaviours**

Number of risk behaviours	Percent of sample	Boys (n = 70)	Girls (n = 80)	Younger Adolescents (n = 68)	Older adolescents (n = 50)
All three risk behaviours	9.8	5.8	13.7	8.2	11.6
2: MVP A < 60 minutes/day and < 5 fruit/vegetables per day	34.0	32.0	34.8	36.4	28.4
2: MVP A < 60 minutes/day and < 5 days a week eating breakfast	6.4	1.2	11.4	5.1	7.8
2: < 5 fruit/ vegetables per day and < 5 days a week eating breakfast	5.7	5.7	5.6	8.2	2.8
1: MVPA < 60 minutes/day	28.0	34.5	24.6	19.4	42.1
1: < 5 fruit/ vegetables per day	7.4	9.3	5.6	11.2	2.8
1: < 5 days a week eating breakfast	2.8	2.3	3.4	4.1	1.4
0: No risk behaviours	6.4	10.4	2.3	7.1	5.2

For adolescents with two risk behaviours, the most prevalent cluster was formed by not meeting the physical activity and fruit and vegetables recommendations. A higher proportion of girls, compared to boys, had the cluster pattern of not meeting the recommendations for physical activity and breakfast consumption. ( $p < 0.01$ ). A higher proportion of boys, compared to girls, and older adolescents, compared to younger adolescents, had the most prevalent single risk factor of not meeting the recommendations for physical activity ( $p < 0.01$ ). A higher proportion of younger adolescents, compared to older adolescents, had the single risk factor of not meeting the recommendations for fruit and vegetable consumption ( $p < 0.01$ ).

### Discussion

This study described the prevalence and clustering patterns of three health behaviours (physical

activity, fruit and vegetable consumption, breakfast consumption) in a sample of adolescents from the Bihar. Almost 54% of adolescents had multiple diet and physical activity risk behaviours, and only 6% achieved the recommendations for all three of the health behaviours. Poor diet and physical inactivity are established risk factors for chronic disease. In young people, physical activity and healthy diets including regular breakfast consumption and adequate levels of fruit and vegetables, have important short- and long-term health protective effects. For example, physical activity in young people may benefit cardiovascular disease (CVD) risk factors, adiposity and bone health, which could influence health in adulthood. [20]

In total, 393 pupils provided consent and completed the questionnaire (95.85% response rate). Of these, 150 (38.16%) provided usable accelerometer data. Comparison of these 150 adolescents with those

who did not provide usable accelerometer data ( $n = 150$ ) showed no significant differences in age, SES, ethnicity or fruit and vegetable consumption. However, a significantly ( $p < 0.01$ ) higher proportion of those with usable accelerometer data were girls, compared to boys (53% compared to 37%), and ate breakfast on more days per week (4.7 days compared to 3.5 days). Thus, our figures likely underestimate the prevalence of not meeting the breakfast recommendations. The final sample composition was 150 adolescents, with 70 boys, 80 girls, 68 younger adolescents and 50 older adolescents. The mean age of younger adolescents was 14.3 years and of older adolescents was 14.6 years. A higher proportion of boys, compared to girls, ate breakfast on more than 5 days/week ( $p < 0.01$ ). A higher proportion of younger adolescents, compared to older adolescents, met the physical activity recommendations and did not meet the recommendations for fruit and vegetable consumption ( $p < 0.01$ ). Only 6% of the adolescent sample met the recommendations for all three of the health behaviours and almost 10% had three health risk behaviours. A higher proportion of girls, compared to boys, had three health risk behaviours ( $p < 0.01$ ).

Higher levels of MVPA were found among boys, compared to girls, and in younger adolescents, compared to older adolescents. Such findings support previous research [21,22] highlighting gender differences and age related declines in physical activity, using objective measures of physical activity among adolescents. Recent research using accelerometry has also shown that age and gender differences are evident when comparing children as young as six and eleven years of age. [23,24] This suggests that the primary school years may be critical for the development of disparities in physical activity behaviours. [25] Efforts to promote physical activity should begin in these critical years given that active children are more likely to become active adults. [26] Consistent with previous research, adolescent girls ate breakfast on fewer days per week compared to boys. [27] Despite the evidence that adolescents who skip breakfast are more likely to be overweight than those who regularly eat breakfast [28], skipping breakfast may be a chosen method of weight control for girls, and in some individuals may be associated with dieting, or disordered eating. [29]

Older adolescents ate more fruit and vegetables per day, and were more likely to meet the recommendations for fruit and vegetable consumption, compared to younger adolescents. In contrast, review level evidence has shown a negative association between age and fruit and vegetable consumption. [30] Contrasting findings may reflect a difference in the methodologies employed to

assess fruit and vegetable consumption. Several studies have identified overestimation of fruit and vegetable intake when using food frequency questionnaires. [31,32] Girls were more likely to have the two behaviour risk cluster that paired physical activity and breakfast consumption. A possible explanation for such clustering is that adolescents who frequently skip breakfast have lower daily energy intakes, with higher daily energy intake being associated with more time spent being physically active. [33] Girls who skip breakfast as part of a diet or method of weight control, may have less energy to be physically active. Efforts to promote both physical activity and regular breakfast consumption to adolescent girls, in particular, are needed. Consistent with previous research, girls had a higher number of risk factors related to physical activity and dietary behaviours compared to boys. [34] Such findings provide additional evidence for support of gender specific interventions promoting physical activity and dietary behaviours. Although health promotion programmes frequently target multiple behaviours, little is known about the best approaches to stimulating multiple behaviour change in adolescents. [34] Reviews of multiple behaviour interventions with young people have revealed changes in some but not all behaviours, with significant effects more likely for dietary as opposed to physical activity outcomes. [35] There is little evidence of covariation among diet and physical activity behaviours in adolescents over time. [36]

For adolescents with two risk behaviours, the most prevalent cluster was formed by not meeting the physical activity and fruit and vegetables recommendations. A higher proportion of girls, compared to boys, had the cluster pattern of not meeting the recommendations for physical activity and breakfast consumption. ( $p < 0.01$ ). A higher proportion of boys, compared to girls, and older adolescents, compared to younger adolescents, had the most prevalent single risk factor of not meeting the recommendations for physical activity ( $p < 0.01$ ). A higher proportion of younger adolescents, compared to older adolescents, had the single risk factor of not meeting the recommendations for fruit and vegetable consumption ( $p < 0.01$ ).

### Conclusion

Many adolescents fail to meet multiple diet and physical activity recommendations, supporting previous evidence that physical activity and dietary behaviours do not occur in isolation. Differences in dietary and physical activity behaviours between adolescent boys and girls, as well as between older and younger adolescents, should be taken into consideration when assessing the efficacy of strategies promoting multiple health behaviour change.

**References**

1. Kotecha PV, Patel SV, Baxi RK, Mazumdar VS, Shobha M, Mehta KG, Mansi D, Ekta M. Dietary pattern of schoolgoing adolescents in urban Baroda, India. *J Health Popul Nutr.* 2013 Dec ;31 (4):490-6.
2. Vir S. Adolescent growth in girls--the Indian perspective. *Indian Pediatr.* 1990 Dec;27(12): 12 49-55.
3. Bull NL. Studies of the dietary habits, food consumption and nutrient intakes of adolescents and young adults. *World Rev Nutr Diet.* 1988;5 7:24-74.
4. Bull NL. Studies of the dietary habits, food consumption and nutrient intakes of adolescents and young adults. *World Rev Nutr Diet.* 1988;5 7:24-74.
5. Caspersen CJ, Powell KE, Christenson GM. Physical activity, exercise, and physical fitness: definitions and distinctions for health-related research. *Public Health Rep.* 1985 Mar- Apr;100 (2): 126-31.
6. Lynda A. Centres for disease control and prevention. Surgeon general's report on physical activity and health. *JAMA Netw Open* 1996; 276:22-9.
7. Parthasarathy A, Menon PS, Agarwal RK, Choudhury P, Thacker NC, Ugra D, et al. *IAP Text Book of Paediatrics* 4th ed New Delhi Jaypee Brothers Medical Publishers 2010 153-5.
8. Biddle SJ, Gorely T, Stensel DJ. Health-enhancing physical activity and sedentary behaviour in children and adolescents. *Journal of sports sciences.* 2004 Aug 1;22(8):679-701.
9. Antova T, Pattenden S, Nikiforov B, Leonardi GS, Boeva B, Fletcher T, Rudnai P, Slachtova H, Tabak C, Zlotkowska R, Houthuijs D. Nutrition and respiratory health in children in six Central and Eastern European countries. *Thorax.* 2003 Mar 1;58(3):231-6.
10. Maynard M, Gunnell D, Emmett P, Frankel S, Smith GD. Fruit, vegetables, and antioxidants in childhood and risk of adult cancer: the Boyd Orr cohort. *Journal of Epidemiology & Community Health.* 2003 Mar 1;57(3):218-25.
11. Barton BA, Eldridge AL, Thompson D, Affenito SG, Striegel-Moore RH, Franko DL, Albertson AM, Crockett SJ. The relationship of breakfast and cereal consumption to nutrient intake and body mass index: the National Heart, Lung, and Blood Institute Growth and Health Study. *Journal of the American Dietetic Association.* 2005 Sep 1; 105(9): 1383-9.
12. Albertson AM, Franko DL, Thompson D, Eldridge AL, Holschuh N, Affenito SG, Bauserman R, Striegel-Moore RH. Longitudinal patterns of breakfast eating in black and white adolescent girls. *Obesity.* 2007 Sep;15(9):2282-92.
13. T DWYER JO, Evans M, Stone EJ, Feldman HA, Lytle L, Hoelscher D, Johnson C, Zive M, Yang M. Adolescents' eating patterns influence their nutrient intakes. *Journal of the American Dietetic Association.* 2001 Jul 1;101(7):798-802.
14. World Health Organization: Young people's health in context: selected key findings from the Health Behaviour in School-aged Children study.
15. Rockett HR, Breitenbach M, Frazier AL, Witschi J, Wolf AM, Field AE, Colditz GA: Validation of a youth/adolescent food frequency questionnaire. *Prev Med* 1997, 26:808-816.
16. Barton BA, Eldridge AL, Thompson D, Affenito SG, Striegel-Moore RH, Franko DL, Albertson AM, Crockett SJ: The relationship of breakfast and cereal consumption to nutrient intake and body mass index: the National Heart, Lung, and Blood Institute Growth and Health Study. *J Am Diet Assoc* 2005, 105:1383-1389.
17. Delva J, Johnston LD, O'Malley PM: The Epidemiology of Over- weight and Related Lifestyle Behaviors. Racial/Ethnic and Socioeconomic Status Differences Among American Youth. *Am JPrevMed*2007, 33:s178-s186.
18. Mattocks C, Ness A, Leary S, Tilling K, Blair SN, Shield J, Deere K, Saunders J, Kirkby J, Smith GD, et al.: Use of accelerometers in a large field-based study of children: Protocols, design issues, and effects on precision. *JPhys Act Health* 2008, 5:S98-S 111.
19. Telford A, Salmon J, Jolley D, Crawford D: Reliability and validity of a self-report and proxy-report physical activity questionnaire: the Children's Leisure Activities Study Survey. *Pediatr Exerc Sci* 2004, 16:64-78.
20. Biddle SJ, Gorely T, Stensel DJ. Health-enhancing physical activity and sedentary behaviour in children and adolescents. *Journal of sports sciences.* 2004 Aug 1;22(8):679- 701.
21. Trost SG, Pate RR, Sallis JF, Freedson PS, Taylor WC, Dowda M, Sirard J. Age and gender differences in objectively measured physical activity in youth. *Medicine & Science in Sports & Exercise.* 2002 Feb 1;34(2):350-5.
22. KIMM SY, GLYNN NW, KRISKA AM, FITZGERALD SL, AARON DJ, SIMILO SL, McMAHON RP, BARTON BA. Longitudinal changes in physical activity in a biracial cohort during adolescence. *Medicine & Science in Sports & Exercise.* 2000 Aug 1;32(8):1445-54.
23. Hesketh K, Crawford D, Salmon J. Children's television viewing and objectively measured physical activity: associations with family circumstance. *International Journal of Behavioral Nutrition and Physical Activity.* 2006 Dec;3(1):1-0.
24. Bagley S, Salmon JO, Crawford D. Family structure and children's television viewing and physical activity. *Medicine and science in sports and exercise.* 2006 May 1;38(5):910-8.
25. Brodersen NH, Steptoe A, Boniface DR, Wardle J.

- Trends in physical activity and sedentary behaviour in adolescence: ethnic and socioeconomic differences. *British journal of sports medicine*. 2007 Mar 1;41(3): 140-4.
26. Telama R, Yang X, Viikari J, Valimaki I, Wanne O, Raitakari O. Physical activity from childhood to adulthood: a 21-year tracking study. *American journal of preventive medicine*. 2005 Apr 1;28(3):267-73.
  27. Pearson N, MacFarlane A, Crawford D, Biddle SJ. Family circumstance and adolescent dietary behaviours. *Appetite*. 2009 Jun 1;52(3):668-74.
  28. Rashidi A, Mohammadpour-Ahranjani B, Karandish M, Vafa MR, Hajifaraji M, Ansari F, Sadeghi S, Maddah M, Kalantari N, Akhavi-Rad MB. Obese and female adolescents skip breakfast more than their non-obese and male peers. *Open Medicine*. 2007 Dec 1;2(4):481-7.
  29. Albertson AM, Franko DL, Thompson D, Eldridge AL, Holschuh N, Affenito SG, Bauserman R, Striegel-Moore RH. Longitudinal patterns of breakfast eating in black and white adolescent girls. *Obesity*. 2007 Sep;15(9):2282-92.
  30. Rasmussen M, Krolner R, Klepp KI, Lytle L, Brug J, Bere E, Due P. Determinants of fruit and vegetable consumption among children and adolescents: a review of the literature. Part I: quantitative studies. *International journal of behavioral nutrition and physical activity*. 2006 Dec;3(1):1-9.
  31. Kremers SP, Brug J, De Vries H, Engels RC. Parenting style and adolescent fruit consumption. *Appetite*. 2003 Aug 1;41(1):43-50.
  32. Van Assema P, Brug J, Ronda G, Steenhuis I, Oenema A. A short dutch questionnaire to measure fruit and vegetable intake: relative validity among adults and adolescents. *Nutrition and Health*. 2002 Apr;16(2):85-106.
  33. Berkey CS, Rockett HR, Gillman MW, Field AE, Colditz GA. Longitudinal study of skipping breakfast and weight change in adolescents. *International journal of obesity*. 2003 Oct;27(10):1258-66.
  34. Sanchez A, Norman GJ, Sallis JF, Calfas KJ, Cella J, Patrick K. Patterns and correlates of physical activity and nutrition behaviors in adolescents. *American journal of preventive medicine*. 2007 Feb 1;32(2):124-30.
  35. Resnicow K, Robinson TN. School-based cardiovascular disease prevention studies: review and synthesis. *Annals of Epidemiology*. 1997 Oct 1;7(7):S14-31.
  36. Rosenberg DE, Norman GJ, Sallis JF, Calfas KJ, Patrick K. Covariation of adolescent physical activity and dietary behaviors over 12 months. *Journal of Adolescent Health*. 2007 Nov 1;41(5):472-8.