

## Spectrum of Supraclavicular Lymph Node on Fine Needle Aspiration Cytology: A One–Year Study in a Tertiary Care Hospital

Satyajit Behera<sup>1</sup>, Abinasha Mohapatra<sup>2</sup>, Himansu Shekhar Mishra<sup>3</sup>

<sup>1</sup>Assistant Professor, Department of General Surgery, D.D. Medical College and Hospital, Keonjhar, Odisha, India, 758001.

<sup>2</sup>Associate Professor, Department of General Surgery, F.M. Medical College and Hospital, Balasore, Odisha, India, 756019.

<sup>3</sup>Assistant Professor, Department of General Surgery, S.C.B. Medical College and Hospital, Cuttack, Odisha, India, 753001.

Received: 01-05-2025 / Revised: 15-06-2025 / Accepted: 21-07-2025

Corresponding author: Dr. Himansu Shekhar Mishra

Conflict of interest: Nil

### Abstract

**Introduction:** Supraclavicular lymphadenopathy is frequently associated with significant underlying pathology. Fine Needle Aspiration Cytology (FNAC) serves as a cost-effective, rapid, and minimally invasive diagnostic method that plays an essential role in the initial evaluation of lymph node lesions. This study analyses the spectrum of supraclavicular lymph node lesions diagnosed via FNAC in a tertiary care centre over a one-year period.

**Methods:** This retrospective observational study was conducted in the Department of General Surgery, F.M. Medical College and Hospital, Balasore, Odisha, included 50 patients presenting with supraclavicular lymphadenopathy who underwent FNAC from January to December 2024. FNAC was performed using a 23-gauge needle. Smears were stained with May-Grünwald-Giemsa (MGG) and special stains like Ziehl-Neelsen were applied when indicated. Cytological diagnoses were categorized into reactive lymphadenitis, acute suppurative lymphadenitis, granulomatous lymphadenitis, metastatic malignancy, lymphoma, and miscellaneous cases.

**Results:** Out of 50 cases, 66% were male and 34% female. Metastatic malignancy was the most common diagnosis (50%), followed by granulomatous lymphadenitis (24%) and reactive lymphadenitis (18%). Left-sided lymphadenopathy was more frequent (60%). Most metastatic cases occurred in the age group of 40–80 years. FNAC is a dependable and cost-effective diagnostic tool in evaluating supraclavicular lymphadenopathy.

**Conclusion:** In this study, metastatic malignancies were the leading cause, especially in older adults. The technique is invaluable for early diagnosis and directing further investigations and treatment.

**Keywords:** Supraclavicular lymph node, FNAC, metastasis, granulomatous lymphadenitis, squamous cell carcinoma, adenocarcinoma.

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

### Introduction

Supraclavicular lymphadenopathy is a common clinical presentation, often prompting investigation for serious underlying conditions. In both young adults and the elderly, such lymphadenopathy may be due to reactive changes, infections such as tuberculosis, or a first sign of malignancy [1].

FNAC has gained wide acceptance as an initial diagnostic procedure due to its affordability, simplicity, and rapid turnaround.

In resource- limited settings like eastern Odisha, where both infectious diseases and malignancies are prevalent, FNAC of supraclavicular nodes becomes especially significant. In selected cases, radiologically guided FNAC further improves

accuracy [2]. This study was undertaken to assess the cytological spectrum of supraclavicular lymphadenopathy using FNAC over a one-year period in Department of General Surgery, F.M. Medical College and Hospital, Balasore, Odisha.

### Aim

To assess the cytological spectrum of supraclavicular lymphadenopathy through FNAC in a tertiary care hospital in Eastern Odisha.

### Materials and Methods

This retrospective study was conducted in the Department of General Surgery, F.M. Medical College and Hospital, Balasore, Odisha. Data were

collected for all patients presenting with supraclavicular lymphadenopathy who underwent FNAC from January 2024 to December 2024. A total of 50 cases were included. Aspirations were done using a 23- gauge needle and 10 mL disposable syringe. Smears were stained with May-Grünwald-Giemsa (MGG) and Papanicolaou stain. Ziehl- Neelsen staining was performed when tuberculosis was suspected. Inadequate aspirates were excluded.

Based on cytomorphology, the cases were categorized into six groups:

- (A) Reactive lymphadenitis
- (B) Acute suppurative lymphadenitis
- (C) Granulomatous lymphadenitis
- (D) Metastatic malignancy
- (E) Lymphoma
- (F) Miscellaneous/non-specific

### Results

**Table 1: Distribution of FNAC Diagnosis of 50 Cases of Lymphadenopathy**

Cause	Male	Female	No.of Cases	Percentage (%)
Metastatic malignancy	17	08	25	50
Granulomatous lymphadenitis	08	04	12	24
Reactive lymphadenitis	06	03	09	18
Lymphoma	01	01	02	04
Acute suppurative lymphadenitis	01	00	01	02
Miscellaneous/non-specific	00	01	01	02

Out of 50 patients, 33 (66%) were male and 17 (34%) were female. The most common diagnosis was metastatic malignancy, seen in 25 cases (50%), followed by granulomatous lymphadenitis in 12 cases (24%) and reactive lymphadenitis in 9 cases (18%). The most frequent site of involvement was the left supraclavicular lymph node (60%).

**Table 2: Age Distribution of Lymph Node Lesions**

Cause	Age in years					Total
	<20	21-40	41-60	61-80	>81	
Metastatic malignancy	0	01	10	11	03	25
Granulomatous lymphadenitis	01	06	04	01	00	12
Reactive lymphadenitis	01	04	03	01	00	09
Lymphoma	00	00	00	02	00	02
Acute suppurative lymphadenitis	00	01	00	00	00	01
Miscellaneous/non-specific	00	00	01	00	00	01

The ages of the patient ranged from 11 years to 88 years. Metastatic malignancy was most common in the age group of 40–80 years.

### Discussion

#### Metastatic:

Present study - 50% (25/50) of cases.

Literature: 65.1 % (Radhakrishnan et al. [3]) and 79.7% (Mitra et al. [4])

Interpretation: In the present study, the most common cause of supraclavicular lymphadenopathy was metastatic malignancy. The majority of metastatic cases were seen in the 40–80 years age group.

#### Granulomatous Inflammation:

Present study: 24% (12/50) of cases

Range: 8.8 % (Radhakrishnan et al. [3]) and 14.2% (Mitra et al. [4])

Interpretation: Among these, 7 cases were confirmed as tuberculous on the basis of cytological features and Ziehl-Neelsen staining. The relatively high incidence in our study reflects

the endemic nature of tuberculosis in Eastern Odisha.

#### Reactive Lymphadenopathy:

Present study: 18 % of cases

Literature: 22 % (Radhakrishnan et al. [3])

Interpretation: This percentage is lower than that reported in other studies, which may be due to the fact that supraclavicular nodes are less commonly involved in localized inflammatory responses.

#### Lymphoma:

Present study: 4 %

Range: 5 % (Mitra et al. [4])

Interpretation: Both cases were provisionally diagnosed as non- Hodgkin lymphoma on cytology and were referred for further evaluation including biopsy and immunophenotyping.

### Conclusion

The present study was conducted over a one-year period and included 50 cases of supraclavicular lymphadenopathy evaluated by fine needle aspiration cytology (FNAC). The most common

cytological diagnosis was metastatic malignancy, observed in 50% (25/50) of cases, followed by granulomatous lymphadenitis in 24% (12/50) and reactive lymphadenitis in 18% (9/50). Lymphoma, suppurative lymphadenitis, and nonspecific categories were seen in 4%, 2%, and 2% of cases respectively.

The findings highlight the predominance of malignancy among supraclavicular lymph node enlargements, especially in the elderly population. The significant proportion of granulomatous lesions, including confirmed cases of tuberculosis, reflects the persistent endemic nature of the disease in Eastern Odisha.

FNAC proved to be a simple, rapid, and cost-effective diagnostic tool that enabled early detection and categorization of lymphadenopathy. Its role remains indispensable in guiding clinical decisions and avoiding unnecessary surgical interventions, particularly in resource-limited settings.

## References

1. (PDF) Pattern of disease in palpable supraclavicular lymph node: A cytopathological perspective [Internet]. [Cited 2025 Jul 23]. Available from: [https://www.researchgate.net/publication/239526239\\_Pattern\\_of\\_disease\\_in\\_palpable\\_supraclavicular\\_lymph\\_node\\_A\\_cytopathological\\_perspective](https://www.researchgate.net/publication/239526239_Pattern_of_disease_in_palpable_supraclavicular_lymph_node_A_cytopathological_perspective)
2. Department Of Pathology, NKP Salve Institute Of Medical Sciences and Research Centre, Digdoh Hills, Hingna,, Nagpur MAHARASHTRA 440019, INDIA, Mohanty R. Utility of Fine Needle Aspiration Cytology of Lymph nodes. IOSR J Dent Med Sci. 2013; 8(5):13–8.
3. Radhakrishnan D, Nair RG, P.P S, K RV. Fine Needle Aspiration Cytology of Supraclavicular Lymph Node- Three Year Retrospective Study. J Evid Based Med Healthc. 2018 Feb 2; 5(6):522–6.
4. Mitra S, Ray S, Mitra PK. Fine needle aspiration cytology of supraclavicular lymph nodes: Our experience over a three-year period. J Cytol Indian Acad Cytol.