

Retrospective Study of Abruptio Placenta and Its Maternal and Fetal Outcome in a Tertiary Care Hospital in the West Coast of India**Priyanka Kushwaha¹, Nalini I. Anand², Inderpal J. Anand³, Mona D. Gandhi⁴, Nita Rada⁵, Trupti C. Nayak⁶**¹2nd Year Resident Doctor, Department of Obstetrics & Gynaecology, Shri M.P. Shah Govt. Medical College, Jamnagar.²Professor and Head, Department of Obstetrics & Gynaecology, Shri M.P. Shah Govt. Medical College, Jamnagar.³Professor, Department of Paediatrics, Shri M.P. Shah Govt. Medical College, Jamnagar.⁴Associate Professor and HOU, Department of OBGY, Shri M.P. Shah Govt. Medical College, Jamnagar.⁵Associate Professor and HOU, Department of OBGY, Shri M.P. Shah Govt. Medical College, Jamnagar.⁶Associate Professor and HOU, Department of OBGY, Shri M.P. Shah Govt. Medical College, Jamnagar.

Received: 01-05-2025 / Revised: 15-06-2025 / Accepted: 21-07-2025

Corresponding author: Dr. Priyanka Kushwaha

Conflict of interest: Nil

Abstract**Background:** Abruptio placenta is one of the common causes of antepartum haemorrhage and is defined as premature separation of normally implanted placenta. It is more common in second half of pregnancy. Abruptio placenta is serious complication of pregnancy and causes high maternal and neonatal morbidity and mortality.**Methods:** This retrospective study of abruptio and its maternal and perinatal outcome was carried out between October 2023 and March 2025 at Shree MP Shah Government Medical College Jamnagar Gujarat.**Results:** There were 12640 deliveries during the study period out of which 50 had abruptio placenta, which had incidence of 0.44%. Majority of patients were unbooked (80%). Abruptio was commonest among 25-30 years of age had lower educational status, more often had 2-3 deliveries. 70% of cases were associated with severe preeclampsia. Live birth was 54% where is still birth were 46%. PPH occurred in 20% of patients and DIC accounts for 12% of complication, acute renal failure accounts for 10%, shock for 10%, 8% patients had infections and 10% with other complications and remaining 30% patients did not have much complications and were stable.**Conclusions:** Abruptio placenta is still a life threatening complication of pregnancy and it is associated with poor maternal and fetal outcome if not managed appropriately. Hence early diagnosis and prompt resuscitative measures would prevent both perinatal and maternal mortality and morbidity. The availability of advanced emergency obstetric care across greater number of referral hospitals has been responsible for decreasing the mortality and morbidity associated with many obstetric conditions. As of now, early referral to tertiary care hospital, better availability of blood and blood products and early interventions have the potential to limit adverse maternal and perinatal outcome.**Keywords:** Abruptio placenta, Maternal, Mortality, Morbidity, Neonatal, Pre-eclampsia.This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.**Introduction**

Placental abruption is the most common cause of antepartum haemorrhage and is defined as premature separation of normally implanted placenta. [1] Placental abruption occurs when there is a compromise of the vascular structures supporting the placenta. In other words, the vascular networks connecting the uterine lining and the maternal side of the placenta are torn away. These vascular structures deliver oxygen and nutrients to the fetus. Disruption of the vascular network may

occur when the vascular structures are compromised because of hypertension or substance use or by conditions that causes stretching the uterus. The uterus is a muscle and is elastic whereas the placenta is less elastic than the uterus. Therefore, when the uterine tissue stretches suddenly, the placenta remains stable and the vascular structure connecting the uterine wall to the placenta tear away. [2,3] Placental abruption is one of the causes of vaginal bleeding in the second half of pregnancy. A focused

history and physical examination is critical to differentiate placental abruption from other causes of vaginal bleeding. Placental abruption is a potentially life-threatening situation. Therefore, accurate assessment of the patient and accordingly management is very important to prevent a potentially poor outcome. The exact etiology of placental abruption is unknown.

However, a number of factors are associated with its occurrence. Risk factors can be thought of in four groups: health history, past obstetrical events, current pregnancy, and unexpected trauma. Factors that increase the risk of placental abruption includes smoking, cocaine use during pregnancy, maternal age over 35 years, hypertension, and placental abruption in a prior pregnancy. Conditions specific to the current pregnancy which may precipitate placental abruption are multiple gestation pregnancies, polyhydramnios, preeclampsia, diabetes mellitus, sudden uterine decompression, and short umbilical cord.

Finally, trauma to the abdomen such as a motor vehicle accident, fall or violence resulting in a blow to the abdomen may lead to placental abruption. Abruption is a significant cause of maternal and perinatal morbidity and mortality. Placental abruption may be total or partial, causing pain and vaginal bleeding-which are the hallmarks of placental abruption.

Abruption placenta is the major cause of haemorrhagic shock, DIC, renal failure and neonatal complications include hypoxia, anemia, growth restriction, prematurity, neurodevelopmental problems and premature death. Hypertensive disorder of pregnancy is associated with placental separation. [4] Maternal and fetal survival depends on early diagnosis and intervention.

The history begins with a review of the prenatal course, especially placental location on prior sonograms and if there is a history of placental abruption in previous pregnancies. History of smoking and potential trauma especially in the abdominal area is important.

Assessment of the patient can provide very important clue for diagnosing onset of abruptio placenta. The physical examination includes palpation of the uterus for tenderness, consistency, and frequency and duration of uterine contractions.

The vaginal area is inspected for the presence of bleeding. Digital examination of the cervix should be delayed until a sonogram is obtained for placental localization and to rule out a placenta previa. If bleeding is present, the quantity and characteristic of the blood, as well as the presence of clots is evaluated. However, the absence of vaginal bleeding does not eliminate the diagnosis of placental abruption.

Methods

This is a retrospective study conducted at Shree MP Shah Government Medical College Jamnagar Gujarat, India from October 2023 to March 2025. Total number of deliveries from October 2023 to March 2025 were 12640.

Total number of abruptio placenta cases from October 2023 to March 2025 were 50. From case sheets, details of the patient like age, parity and maternal high-risk factors were collected. Patients with other causes of APH like placenta previa and other extra placental causes were excluded.

All study patients underwent a complete obstetrical examination and clinical workup including history, general physical examination and abdominal and pelvic examination. Detailed obstetric history was obtained and maternal high-risk factors like PIH, GDM, and polyhydramnios was noted. As 95% patients were admitted as emergencies, placental abruption was suspected depending on clinical features like vaginal bleeding, uterine tenderness, hypertonic uterus and diagnosis was confirmed by retroplacental clots. After initial resuscitation, mode of delivery was decided depending upon state of mother and fetus. Relevant blood investigations were done.

Ultrasound for fetal well-being was done. Diagnosis was confirmed by the presence of retroplacental clots which was used to estimate the amount of bleeding and severity of abruption. Patients were managed according to the fetal and maternal conditions. Accordingly, results were analysed. Maternal complications studied were PPH, DIC, ARF, shock, pulmonary edema and infections. Fetal outcome were studied in the form of perinatal mortality (still births and neonatal deaths), prematurity and admission to the neonatal care unit.

Results

Maximum no. of cases (40%) of abruptio placenta were between 25 to 30 years of age. Increase maternal age is risk factor for abruptio placenta (Table 1). Maximum number of cases was Gravida 2 (35%) (Table 2). Chances of abruptio placenta increase with increase parity. Patients who had severe preeclampsia have maximum rate of occurrence of abruption (70%) (Table 3). Most of them were associated with anemia and PIH. 6 % patients with eclampsia had abruptio placenta. Even normotensive groups had abruption which was about 24 %. 54 % had live birth, 46% had still born baby (Table 4).

Among them 5 died in early neonatal period due to prematurity. Fetal complications included hypoxia, anemia, growth restriction, and prematurity, and neurodevelopmental problems, prematurity and fetal death. Maternal complications associated with

abruption were Postpartum Haemorrhage (PPH) (20%), Disseminated Intravascular Coagulation (DIC) (12%), Acute renal failure (ARF) (10%),

Shock (10%). 8% patients had infections and 10% suffered from other complication (Table 5).

Table 1: Association with age.

Age	Percentage
20-25 years	10%
25-30 years	40%
30-35years	30%
>35 years	20%

Maximum number of cases (40%) of abruptio placenta were between 25 to 30 years. Next most common age group were between 30 to 35 years (30%). Least incidence was seen among the age group 20-25 years (10%).

Table 2: Association with parity.

Parity	Percentage
G1	15
G2	35
G3	30
G4,G5	20

Maximum numbers of cases were gravida 2 (35%). 15 % patients were gravida 1 and 30% were third gravida and 20% were gravida 4th and 5th.

Table 3: Association with PIH.

Type	Percentage
Severe preeclampsia	70%
eclampsia	6%
Normal BP	24%

Patients who had severe preeclampsia have maximum rate of abruptio (70%). Most of them were associated with anemia and PIH. 6 % patients with eclampsia had abruptio placenta. Even normotensive groups had abruption which was about 24%.

Table 4: Fetal outcome.

Birth	Percentage
Still birth	46%
Live birth	54%

54% had live birth, 46% had still born baby. Among live born 5 died in early neonatal period due to prematurity. Fetal complications included hypoxia, anemia, growth restriction, and prematurity, and neurodevelopmental problems, prematurity and fetal death.

Table 5: Maternal complications.

Complication	Percentage
PPH	20
DIC	12
ARF	10
Shock	10
Infections	08
Others	10

Maternal complications associated with abruption were postpartum haemorrhage (pph) (20%), disseminated intravascular coagulation (dic) (12%), acute renal failure (arf) (10%), shock (10%). 8% patients had infections and 10% suffered from other complication and rest others (30%) did not had any complications.

Discussion

Placental abruption is one of the serious complications of pregnancy, as it causes both poor maternal and fetal morbidity and mortality. The incidence of abruptio placenta was 0.44 in our study, which is somewhat near to study by Wasnik SK. [4]

The signs and symptoms of abruptio placenta vary depending upon the severity of bleeding and the degree of separation of the placenta. Abruptio can occur at any trimester in pregnancy but mostly it occurs at 28-32 weeks of pregnancy. [5] It was found that 70% of patients with severe preeclampsia, 6% of patients with eclampsia, 24 % of patients with normal BP developed abruptio in the study. Among the maternal complications, postpartum hemorrhage (PPH) was commonest followed by disseminated intravascular coagulation (DIC). PPH occurred in 20% of patients in the study, were as study by Talpur NN reported PPH in 28% of patients. [6] DIC was associated with 12% of the patients in the study. Sher G observed DIC in 10-20% of his study patients with severe abruptio and fetal demise which is comparable to the study. [7] It was found that ARF is reported in 10% of the cases and shock in 10% of patients whereas study from Shrivatsava V reported 24.6% shock cases. [9] Infection was found to be in 17.5% of patients in the study by Choudhary V, in our study it is reported in 8% of the patients. [10] Regarding fetal outcome, 54% were born alive and 46% were still births. A premature delivery can increase the fetal morbidity in cases of abruptio. [11-15] Routine antenatal check-up, treatment of anemia, timely referral to advanced centres and proper management with timely caesarean section, blood transfusion and good neonatal intensive care unit will further lower the perinatal and maternal morbidity and mortality.

Conclusion

Increasing age has been implicated as a predisposing factor in Abruptio placenta. Most of the patients were unbooked and incidence of abruptio was high in multiparous. The study reveals that severe preeclampsia increases incidence of abruptio. Mainly abruptio was seen in term pregnancy. Majority of patients had associated anemia and PIH, and the mode of delivery varied according to maternal and fetal factors. Major complication on maternal side was PPH and fetal complications included hypoxia, anemia, growth restriction, and prematurity, and neurodevelopmental problems, prematurity and fetal death. Thus this study suggests that severe pre-eclampsia, eclampsia, high parity is independent risk factors for abruptio placenta. Antenatal care which identifies the risk factors like PIH plays an important role in decreasing the incidence of abruptio placenta and improving the maternal and fetal outcome. Regular antenatal checkup, anemia correction, early diagnosis and identification of gestational hypertension would

prevent the maternal and perinatal morbidity and mortality. It should be managed in centres with advanced maternal and neonatal facilities. Though maternal morbidity can be reduced with modern management of abruptio placenta but timely diagnosis and intervention is necessary. Early detection and active management will reduce morbidity and mortality.

References

1. Workalemahu T, Enquobahrie DA, Gelaye B, Thornton TA, Tekola F, et al. Abruptio placentae risk and genetic variations in mitochondrial biogenesis and oxidative phosphorylation: replication of a candidate gene association study. *Am J Obstet Gynecol*. 2018; 219(6):617.
2. Sylvester HC, Stringer M. Placental abruptio leading to hysterectomy. *BMJ*. 2017; 11:211-5.
3. Miller C, Grynspan D, Gaudet L, Ferretti E, Lawrence S, et al. Maternal and neonatal characteristics of a Canadian urban cohort receiving treatment for opioid use disorder during pregnancy. *J Dev Orig Health Dis*. 2018; 16:1-6.
4. Talpur NN, Memon SR, Jamro B, Korejo R. Maternal and fetal morbidity with abruptio placentae. *Rawal Med J*. 2011; 36(4):297-300.
5. Sher G. Pathogenesis and management of uterine inertia complicating abruptio placentae with consumption coagulopathy. *Am J Obstet Gynecol*. 1977; 129:164-70.
6. Campbell S, Lee C. Disorders of placentation. *Obstetrics*. 17th ed. Arnold London. 2002:171-3.
7. Shrivastava V, Kotur P, Jauhari A. Maternal and fetal outcome among abruptio placentae at a rural tertiary hospital in Karnataka, India: A Retrospective analysis. *Int J Res Med Sci*. 2014; 2(4):1655-8.
8. Choudhary V, Rathi S, Somani S. Evaluation of risk factors and obstetric and perinatal outcome in abruptio placentae. 2015;14(5):36-9
9. Sheiner E, Shoham-Vardi I, Hadar, Hallak M. Incidence, obstetric risk factors and pregnancy outcome of preterm placental abruptio: a retrospective analysis. *J Matern Fetal Neonatal Med*. 2002; 11(1):34-9.
10. Allred LS, Batton D. The effect of placental abruptio on the short-term outcome of premature infants. *Am J Perinatol*. 2004;21(3): 157-62.