REVIEW ARTICLE

Novel Approaches and Emerging Trends in Hydrotropic Solubilization Technology

Aliunui Aime, Harpritkaur Bagga, Gunjan Chaudhari, Ujwal Vyas*

Datta Meghe College of Pharmacy, Datta Meghe Institute of Higher Education & Research (DMIHER), Deemed to be University, Wardha, Maharashtra, India.

Received: 27th July, 2024; Revised: 12th August, 2024; Accepted: 31st August, 2024; Available Online: 25th September, 2024

ABSTRACT

Hydrotropic solubilization technology enhances the solubility and bioavailability of poorly water-soluble drugs, a major challenge in pharmaceutical research. This review discusses hydrotropic agents' structures, the solubilization process, and factors affecting drug solubility. It explores optimizing agent concentrations, the role of computer modeling and QSAR studies, and hydrotropes' impact on overcoming biological barriers and improving pharmacokinetics and pharmacodynamics. The review highlights hydrotropic compounds' potential in advancing pharmaceutical formulations and clinical applications.

Keywords: Hydrotropic solubilization, Bioavailability, Poorly water-soluble drugs, Pharmacokinetics, Quantitative structure-activity relationship (QSAR).

International Journal of Drug Delivery Technology (2024); DOI: 10.25258/ijddt.14.3.69

How to cite this article: Aime A, Bagga H, Chaudhari G, Vyas U. Novel Approaches and Emerging Trends in Hydrotropic Solubilization Technology. International Journal of Drug Delivery Technology. 2024;14(3):1736-1744.

Source of support: Nil.

Conflict of interest: None

INTRODUCTION

Numerous factors, including the amount of drug, pH, gelation temperature (Tsol–Tgel), mucoadhesive strength, viscosity, statistical analysis, and study on histology, rapid stability analysis, *in-vitro* dispersion analysis, and ex vivo penetration analysis to improve our formulation.[1] The drug component's nature, physical characteristics, and qualities were examined, and the melting point was determined using a capillary and the melting point device.[2] In drug development, insufficient solubility remains a prevalent and worsening issue, underscoring the need to enhance drug solubility through various techniques.[3] Factors such as temperature, molecular weight, pH, and the physical and chemical properties of solutes and solvents influence solubility. Thermodynamic equilibrium solubility is reached when the system's energy is minimized.[4] BCS class II drugs, like phenytoin and nifedipine, and BCS class IV drugs, such as hydrochlorothiazide and taxol, illustrate poorly soluble medications.[5,6] Hydrotropy, first reported by Neuberg in 1916, increases the solubility of one solute in water by adding a significant quantity of another solute, a hydrotrope. Hydrotropic solutions generally show weak interactions between hydrotropes and solutes and lack colloidal properties.[7] Hydrotropes, salts of organic acids, greatly enhance the solubility of sparingly soluble solutes.[8] The rise of genomics, combinatorial chemistry, and high-throughput screening has led to an increase in potential drug candidates, many of which are poorly soluble and hydrophobic. Solubility is a critical property for drug development, affecting bioavailability and the feasibility of various dosage forms. Experimental formulations often require high concentrations for preclinical studies, and while organic co-solvents like DMSO are used, they pose toxicity risks and are unsuitable for therapeutic applications.[9] Hydrotropes offer a promising solution to these challenges, providing a robust solubilizing system across a range of poorly soluble drugs. Despite the extensive study of hydrotropes like nicotinamide, which offers limited solubility enhancement, there is a need for stronger hydrotropes to create more effective solubilization vehicles.[10,11] The development of a versatile hydrotropic system faces challenges, including accommodating diverse pharmacological compounds and optimizing hydrotrope doses. This study explores the solubilizing characteristics of hydrotropic systems, with ongoing research addressing these challenges.[11-15]

Steps of Hydrotropic Solubilization

Hydrotropic solubilization enhances the solubility of poorly water-soluble drugs using hydrotropic agents

- Selection of Hydrotropic Agents: Choose agents like urea, sodium benzoate, sodium salicylate, nicotinamide, or sodium citrate based on their solubilization efficacy for the target drug.

*Author for Correspondence: ujval.pharmacy@dmiher.edu.in



Figure 1: Steps of hydrotropic solubilization



Figure 2: Diagram of hydrophilic part

- Preparation of Hydrotropic Solution: Create a water-based solution with the selected agent, adjusting concentration based on drug solubility and desired final formulation concentration.
- Drug-Hydrotrope Interaction: Study interactions to optimize solubilization and formulation parameters.
- Solubilization Technique: Use methods like mixing, heating, sonication, or stirring to improve drug solubility.
- Formulation Characterization: Assess the solubilized drug-hydrotrope complex using UV-visible spectroscopy, HPLC, or DSC.
- Optimization of Formulation: Adjust hydrotrope concentration, temperature, and pH for maximum solubilization. The steps of hydrotropic solubilization are shown in Figure 1.

The basic structure of hydrotrope and its history:

Neuberg introduced the term "hydrotropy" to describe increased solubility of a solute with high concentrations of organic acid alkali metal salts.[7] Initially applied to non-micelle-forming substances, hydrotropy involves cooperative intermolecular interactions rather than complexation or cosolvency. Hydrotropic agents, like sodium benzoate, typically have a hydrophobic aromatic ring and an anionic group that enhances aqueous solubility. Studies[16,17] found that amphiphilic drugs, such as sodium ibuprofen, can act as effective hydrotropes. El-Khordagui[16] proposed that some cationic salts and neutral molecules, including procaine HCl and pyrogallol, may also exhibit hydrotropy. The basic structure of a hydrotrope is shown in Figure 2.

Significance of hydrotropic solubilization technique[17]

- Because hydrotropy's solvent character is very selective, pH-independent, and emulsification-free, it is considered a superior solubilization technique compared to other methods such as salting in, cosolvency, micellar solubilization, and miscibility.
- All that needs to be done is combine the medication and hydrotrope in water.
- The medication and hydrotrope only need to be mixed with water.

Mechanism of solubilization

Numerous studies have investigated hydrotropic solubilization mechanisms. One proposed mechanism is complexation, where hydrotropic agents form complexes with poorly soluble drugs, enhancing solubility, exemplified by the riboflavin-nicotinamide system. Another mechanism involves the self-association of hydrotropes in aqueous solutions, such as nicotinamide, which improves solubility.[18] Molecular dynamics simulations have shed light on interactions during dissolution and self-assembly of hydrotropes like nicotinamide.[19] Research also explores other agents, like N,N-diethyl nicotinamide, to understand solubilization mechanisms.[20]

Other Mechanisms Proposed Are

Salt-induced hydrotropy

Salt-induced hydrotropy enhances the solubility of poorly soluble substances by adding salts like sodium salicylate, sodium benzoate, and sodium citrate to aqueous solutions. These salts dissociate into ions, which interact with solute molecules through ion-dipole interactions, breaking hydrophobic bonds and facilitating the solute's dispersion in water.[21]

Micellar hydrotropy

Amphiphilic compounds, like surfactants, form micelles in aqueous solutions, similar to micellar hydrotropy, which involves a critical micelle concentration (CMC). Hydrotropes also have a solubilization threshold called the "minimum hydrotrope concentration" (MHC). Micelles feature a hydrophobic core with hydrophilic heads outward and hydrophobic tails inward, allowing hydrophobic solutes to dissolve in the aqueous solution.[22] Lower alkanoates, alkyl sulfates, and alkylbenzenesulfonates exhibit hydrotropic behavior, while longer alkyl chains show micellar behavior.[23]

Co-solvent hydrotropy

Co-solvent hydrotropy is a technique that entails adding organic solvents that are miscible with water to aqueous

solutions in order to enhance the solubility of hydrophobic substances. The organic solvent molecules engage in interactions with both water molecules and solute molecules, creating a more advantageous environment for solubilization.[24] Typical solvents used include ethanol, propylene glycol, and polyethylene glycol.

Polymeric hydrotropy

Water-soluble polymers are used in polymeric hydrotropy to increase the solubility of hydrophobic compounds in aqueous solutions. By interacting or forming complexes with the solute molecules, these polymers may efficiently increase the solute molecules' dispersibility in water.[25] Among the polymers used in polymeric hydrotropy include cyclodextrins, hydroxypropyl methylcellulose (HPMC), and polyvinylpyrrolidone (PVP).

Types of Hydrotropic Agent

Aromatic hydrotropes

Toluene, xylene, and cumene, benzene derivatives, act as aromatic hydrotropes, dissolving hydrophobic chemicals through π-π interactions and hydrophobic forces. Their effectiveness depends on alkyl chain length and aromatic ring substitution. The minimum hydrotropic concentration (MHC) is measured using an aggregate-to-monomer emission ratio method, avoiding interference from probes or additives.[26] Nicotinamide, a safe vitamin B3, is a common hydrotropic agent, with complexation as the primary solubilization method.[14] Hydrotropes include anionic (e.g., sodium benzoate) and cationic (e.g., para-aminobenzoic acid hydrochloride) types.[27]

Aliphatic hydrotropes

Alcohols with a straight or branched chain, like ethanol and isopropanol, are classified as aliphatic hydrotropes. These substances cause hydrogen bonding to break down the structure of water, which promotes solute solubility. Examples of aliphatic hydrotropes are sodium alkanoate, urea and *N, N*-dimethyl urea.[27]

Polymeric hydrotropes

Because they are amphiphilic, polymers like polyethylene glycol (PEG) and polyvinylpyrrolidone (PVP) display hydrotropic behavior. These polymers can solubilize a wide variety of substances because they include segments that are both hydrophilic and hydrophobic. So far, many polymeric solubilizing systems, such as polymeric micelles and micelle-like aggregates from polymeric amphiphiles, have been used to enhance the solubility of pharmaceuticals that are not easily soluble in water.[25]

Sugar-based hydrotropes

Sugars such as glucose and sucrose have hydrotropic qualities because of the hydrophilic hydroxyl groups that they contain. As a result of their formation of hydrogen bonds with water molecules, hydrophobic substances become more easy to dissolve.[26]

Factors Affecting Solubility of Drugs[14,26-29]

Nature of solvent

The solubilization process is tremendously impacted by the nature of the solvent that is used. Using hydrogen bonding, polar solvents like water are able to effectively dissolve polar solutes. On the other hand, nonpolar solvents, which are characterized by van der Waals forces, are more successful in dissolving nonpolar solutes. In order to achieve effective solubilization, it is essential that the solute be compatible with the polarity composition of the solvent.

Temperature

The kinetics of solubilization are significantly impacted by temperature, which is an important factor. In endothermic dissolving processes, the solubility of the substance often rises as the temperature becomes higher. The solubility of substances may, however, decrease with increasing temperature in the case of exothermic activities. When it comes to optimizing solubilization settings, having a solid understanding of the correlations between solubility and temperature is very necessary.

Pressure

Pressure may affect solubility, especially in systems where gases are present. For example, Henry's law says that the solubility of a gas in a liquid is exactly proportional to the partial pressure of the gas above the liquid. High pressure may promote gas solubility, altering solubilization processes in systems like carbonation.

Particle size and surface area

Finely split solutes with greater surface area display higher solubility owing to improved interaction with the solvent. Particle size reduction using processes like milling or micronization may considerably enhance solubilization efficiency, especially for weakly soluble substances.

Polarity

According to the principle of "like dissolves like," polar chemicals dissolve readily in polar solvents and nonpolar substances dissolve readily in nonpolar solvents.

Commonly Used Hydrotropes[30-34]

Hydrotropes are known to self-assemble in solutions. Due to the wide range of compounds exhibiting hydrotropic behavior, including ethanol, salicylates, caffeine, and SDS, classifying them by chemical structure is complex. The present investigation focuses on the enhancement of the solubility and dissolution rate of poorly water-soluble drug rosuvastatin calcium by hydrotropic and mixed hydrotropic agents in solid dispersion techniques. The techniques of rotary evaporation and lyophilization prepared solid dispersion by using urea, sodium salicylate, and sodium benzoate. The study will evaluate the compatibility of pharmaceutical excipients and compare the solubility and dissolution profiles of the rosuvastatin calcium solid dispersion.

MATERIALS AND METHODS

This review article examines hydrotropic solubilization technology through a comprehensive analysis of literature on hydrotrope composition, history, and types. Key sources include recent studies, computational modeling, and QSAR analyses related to hydrotropic systems. The review synthesizes findings on hydrotropic agent optimization and their impact on drug solubility, bioavailability, and pharmacokinetics. Data were collected from peer-reviewed journals and scientific databases, focusing on advancements in hydrotropic technology and its implications for pharmaceutical formulations and patient care.

Evaluation of Rosuvastatin Calcium Solid Dispersions

- To determine the percent drug content, 100 mL of 0.05 M citrate buffer pH 6.6 was added to 10 mg rosuvastatin calcium in SDRE, SDFD and physical mixtures with hydrotropic agents (sodium benzoate, sodium salicylate, urea) in a 100 mL volumetric flask. Then, the solution was filtered through a 0.45 µm syringe filter after 30 minutes of shaking and appropriately diluted with the same solvent. Read the UV-visible spectrophotometer at 241 nm.[35] The analysis was repeated three times for accuracy.
- Percentage determination of drug yield the weight of the dried solid dispersion was determined by the following procedure, which then multiplied by the weight of the starting components yielded the percentage yield.[36]
- The solubility of pure rosuvastatin calcium, solid dispersions, and physical mixtures was determined by doing the solubility study in distilled water. Saturation solubility was measured in triplicate by the Higuchi and Connors method. Samples were shaken for 24 hours at 37 ± 0.5°C in a mechanical isothermal bath shaker and then allowed to equilibrate for a further 24 hours. The aliquots were filtered through a 0.45 µm syringe filter, diluted, and analyzed using a UV-visible double-beam spectrophotometer.[37]
- Dissolution rate studies for rosuvastatin calcium solid dispersions, physical mixtures, and pure drug were conducted using USP apparatus I with a basket stirrer. The dissolution medium consisted of 900 mL of 0.05 M citrate buffer (pH 6.6), recommended by the FDA for rosuvastatin calcium.[32] The system was maintained at 37 ± 0.5°C with a rotation speed of 50 rpm. Ten mg of rosuvastatin calcium from each solid dispersion was used, and 5 ml samples were taken at 5, 10, 15, 20, 30, and 45 minutes and replaced with fresh medium. Samples were filtered through a 0.45 µm filter, diluted, and analyzed spectrophotometrically at 241 nm. Results were plotted against time and compared with the pure drug. Each formulation was tested in triplicate.[36,37]
- Fourier Transform Infrared Spectroscopy (FTIR) was conducted using an FT/IR-6700 JASCO spectrophotometer. FTIR spectra were recorded for samples (2 mg in 200 mg KBr) across a 400 to 4000 cm^{-1} range at 4 cm^{-1} resolution. Analyzed samples included pure rosuvastatin calcium, sodium benzoate, sodium salicylate, and urea.[38]
- Differential Scanning Calorimetry (DSC) was performed using a DSC 3 Mettler Toledo. Thermograms were recorded for hydrotropic agents, solid dispersions (SDs), drug physical mixtures (PM), and pure rosuvastatin (ROS). Four milligrams of each sample were weighed, sealed in an aluminum pan, and heated from 30 to 300°C at 10°C per minute in a nitrogen environment with a 20 mL/min flow rate.[38]

Computational Modelling and Simulation

In many branches of science and engineering, computational modeling and simulation are effective methods for comprehending, forecasting, and improving complicated systems. In the context of developing new medications and employing hydrotropic agents to solubilize poorly water-soluble medications.[39]

Molecular Dynamics (MD) Simulations

Purpose

Molecular dynamics (MD) simulations provide detailed insights into molecular interactions, making them essential for studying the solubilization of poorly water-soluble drugs with hydrotropic agents. Acting as an *in-silico* laboratory, MD simulations visualize structural changes and molecular behaviors, revealing the formation and mechanisms of drug-hydrotrope complexes. These simulations also predict the dynamic stability and thermodynamic properties of drug-hydrotrope systems, enabling the optimization of formulations by modeling varying concentrations, solvents, and temperatures. MD simulations thus support experimental studies and guide the rational design of effective drug delivery systems by offering crucial molecular-level insights.[40,41]

Applications

Comprehending molecular interactions

The MD simulations can thus provide valuable insights into the formation process and stability of molecular complexes, that is, with dynamic interactions between hydrotropic agents and weakly water-soluble medicines.

Virtual experimentation

They permit *in-silico* tests in a manner tantamount to a 'virtual laboratory.' Hence, MD simulations allow for testing under very different parameters, like temperature or solvent type and concentration—all relevant when attempting to optimize drug delivery systems.[41]

Atomic-level visualization

In addition, MD simulation is able to offer a unique view into atomic-level interactions and the structure of molecules. This kind of graphic representation helps in the best communication of research findings and explanations of complicated chemical processes.

Optimization of formulation conditions

MD simulations allow for virtual experiments in which several formulation conditions can be changed—for instance,

concentration, temperature, and choice of solvent. It helps in optimizing conditions for maximum solubilization efficiency.

Correlation with experimental data

Such MD simulations deliver results correlating with the experimental data to validate computational predictions. This iterative process shall guarantee a thorough understanding of the solubilization process and enhance the reliability of the findings.[42]

Quantitative Structure-Activity Relationship (QSAR)

Computational techniques like quantitative structure-activity relationship (QSAR) investigations help establish the link between a molecule's chemical structure and its solubility efficiency. QSAR connects molecular characteristics, such as lipophilicity and molecular weight, to the solubilization efficiency of hydrotropic agents with poorly water-soluble drugs. After pre-processing the dataset, statistical methods or machine learning algorithms develop a predictive QSAR model, which is validated using statistical metrics. This approach aids in optimizing hydrotropic agents for effective drug delivery systems and their logical design.[43] The QSAR study for solubilization of poorly water-soluble drugs is shown in Table 1.

Optimization Of Hydrotropic Agent Concentrations

Selecting optimal hydrotropic agents for poorly water-soluble drugs requires careful consideration. The process involves experimental studies and computer modeling to determine the best concentration for maximum solubilization efficiency. Agents are chosen based on chemical structures, with quantitative structure-activity relationship (QSAR) analyses predicting their solubilizing potential. Experimental validation, including *in-vitro* assays and dissolution studies, fine-tunes these predictions. Systematic optimization varies agent concentrations to improve drug solubility, bioavailability, and therapeutic efficacy, merging computational insights with empirical data for effective drug delivery systems.[44]

Table 1: QSAR Study for solubilization of poorly water-soluble drugs

Step	*Description*
Specify the goal	Outline the purpose and targets of the research, with an emphasis on the solubilization of medications with hydrotropic agents.
Data collection	Compile a dataset that includes the chemical structures, solubilization efficiencies, and hydrotropic agents.
Molecular descriptors	Determine the pertinent molecular descriptors (such as lipophilicity and molecular weight) in order to quantify chemical structures.
Data pre-processing	Treat outliers, handle missing values, and normalize the dataset to make it clean and ready for pre-processing.
Model development	Utilizing statistical methods (such as machine learning and linear regression) to establish a relationship between molecular descriptors and solubilization efficiency, create the QSAR model.
Validation	To guarantee dependability and prediction precision, validate the QSAR model with statistical measures.

Virtual Screening of Hydrotropic Agents

Virtual screening is a key computational method in drug development for identifying potential hydrotropic solubilizers for poorly water-soluble drugs. This approach involves screening chemical databases using algorithms like pharmacophore-based methods or molecular docking. Scoring functions assess binding affinity, while molecular descriptors and filters refine candidate selection. Experimental validation confirms predictions, streamlining lead optimization. This method accelerates the development of high-bioavailability drug delivery systems, making drug development faster and more cost-effective, especially for low-solubility pharmaceuticals.

Database Selection

In the *in-silico* screening of hydrotropic compounds for poorly water-soluble pharmaceuticals, a well-curated and diversified chemical database is crucial. Researchers should select a database including custom libraries, natural products, and commercial substances, considering solubilization mechanisms, structural complexity, and chemical diversity. A well-chosen database ensures comprehensive chemical space coverage and improves the likelihood of identifying effective solubilizers. It should align with study objectives, focusing on agents known to enhance medication solubility.[45]

Lead Optimization

In virtual screening, lead optimization involves refining hydrotropic agents with the best interactions through structural changes or computational techniques to enhance solubilization. This iterative process improves chemical structure for better drug solubility. Lead optimization bridges computational predictions with real applications, guiding the selection of agents with high potential for experimental validation and advancement in drug delivery systems.[45]

Synergistic Effects in Solubilization

Synergistic solubilization involves combining hydrotropic agents with co-solvents or surfactants to enhance the solubility of poorly water-soluble drugs. This approach leverages the strengths of each component, achieving a more effective outcome than individual contributions. By integrating hydrotropic agents with other agents, the rate of drug solubilization increases, overcoming solubility challenges and improving bioavailability and pharmacological efficiency. This method also reduces individual component concentrations, minimizing potential toxicity. The synergy between solubilization techniques supports the development of advanced drug delivery systems with better stability, controlled release, and improved patient outcomes.[46]

Different Techniques for Solubility Enhancement

Physical modifications

Particle size reduction by micronization, nanosuspension, dispersion of drug in carriers like eutectic mixes, solid dispersions, solid solutions, and cryogenic procedures; crystal habit modification by polymorphs, amorphous form, and co-crystallization.

Chemical modifications

Change of pH, use of buffer, derivatization, complexation, and salt formation.

Miscellaneous methods

Adjuvants such as surfactants, solubilizers, co-solvency, hydrotrophy, and innovative excipients are used in the supercritical fluid process.[47]

Biological Barrier and Transport Mechanism

Biological barriers

Poorly water-soluble drugs struggle with effective delivery due to biological barriers in cellular membranes, which hinder their passive diffusion and impact absorption and bioavailability. Hydrophobic properties limit their passage through these membranes, which are primarily composed of proteins and lipids. Hydrotropic agents enhance solubility, aiding in overcoming these barriers and improving drug absorption. Advances in nanotechnology further refine drug delivery by targeting specific cellular uptake mechanisms. Addressing these biological limitations is crucial for enhancing the therapeutic efficacy of hydrophobic medications and optimizing formulations.[48]

Passive diffusion

Molecules travel across biological membranes by a basic process called passive diffusion, which transfers them from a region of higher concentration to one of lower concentration. It runs on the gradient of concentration and doesn't need any energy.

Characteristics of Passive Diffusion

Concentration gradient

This is a type of passive diffusion based on differences in concentration for a given substance inside and outside the cell. Movement of substances takes place in this direction of gradient till it reaches equilibrium.

Lipid solubility

The main passive diffusion barrier is the lipid bilayer of cellular membranes. Thus, substances that are lipophilic or hydrophobic cross the membrane rich in lipids more easily.

Size and charge

The smaller the molecule and more neutral the charge, the more easily the molecule will diffuse. Charged molecules may be completely unable to move through the plasma membrane because the lipids are hydrophobic.[49]

Active transport

Active transport is a key biological process by which cells transfer molecules against their concentration gradient, from low to high areas of concentration. Unlike passive diffusion, active transport requires an energy supply that drives the translocation of materials through the cell membrane; this is often provided by adenosine triphosphate, ATP. This process makes sure of the selective intake of certain molecules and thus sustains cellular homeostasis.

Characteristics of Active Transport

Energy utilization

Active transport is an application process in which energy is used to move the molecules against their concentration gradient. The energy is usually derived from the hydrolysis of ATP.

Carrier proteins

Proteins in the membrane-associated with active transport are referred to as carrier proteins or pumps. In this case, the protein undergoes a conformational change in an energetically dependent process for the active movement of substances across the membrane.

Against concentration gradient

Active transport moves molecules against their natural tendency to diffuse down the concentration gradient, allowing cells to accumulate substances even when their extracellular concentration is low.[50]

Facilitated diffusion

Facilitated diffusion allows compounds to move from higher to lower concentration across cell membranes via specific carrier or channel proteins, aiding those that struggle to cross the lipid bilayer due to their hydrophilic nature, size, or charge.

Features of Diffusion with Assistance

Protein facilitation

Substances can pass across cell membranes more easily when they are transported by carrier or channel proteins. Certain chemicals can be transported through the channels made by these proteins.

Selective specificity

Carrier proteins that display selectivity for specific compounds enable extremely selective diffusion. These proteins' binding sites are designed to identify and transfer specific compounds.

Even though proteins are involved in assisted diffusion, it is still a passive process because it depends on the concentration gradient and doesn't use any energy.[51]

Endocytosis and Exocytosis

Materials are moved into and out of cells by the biological processes of exocytosis and endocytosis, respectively. These processes all significantly impact the regulation of cellular communication, waste elimination, and nutrition intake.

Endocytosis

Endocytosis is a cellular process by which cells engulf external materials, forming vesicles derived from the cell membrane. There are several types of endocytosis, including phagocytosis, pinocytosis, and receptor-mediated endocytosis.

- Phagocytosis: In phagocytosis, large particles such as bacteria or cellular debris are engulfed by the cell. Specialized cells, like macrophages, are often involved in this process.
- Pinocytosis: Pinocytosis, or "cell drinking," involves the non-selective uptake of fluid and solutes. The cell membrane invaginates, trapping extracellular fluid and solutes in small vesicles.

Exocytosis

Exocytosis is the reverse process of endocytosis. It involves the release of substances from the cell by fusion of vesicles with the cell membrane. This results in the extracellular release of the vesicle's contents.

- *Vesicle fusion*

Secretory vesicles, formed within the cell, move towards the cell membrane and fuse with it. This fusion releases the contents of the vesicle into the extracellular space.

- *Role*

Exocytosis is crucial for the secretion of various substances, including hormones, neurotransmitters, and enzymes. It is involved in processes such as cell communication, neurotransmission, and the release of digestive enzymes.[52]

In-Vivo Pharmacokinetics and Pharmacodynamics

Solubilization of poorly water-soluble drugs

Many drugs, especially those with low aqueous solubility, face challenges in absorption and bioavailability. Hydrotropic agents act by increasing the solubility of such drugs in aqueous solutions through the formation of complexes or micelles.

Pharmacokinetics and pharmacodynamics

Enhanced solubility from hydrotropic agents improves drug absorption and bioavailability, leading to better dissolution in the gastrointestinal tract and across biological membranes. This increased bioavailability can result in more effective therapeutic concentrations and improved pharmacological effects.

In-vivo studies

In-vivo pharmacokinetic studies involve the assessment of drug absorption, distribution, metabolism, and elimination in living organisms. In the presence of hydrotropes, such studies could reveal changed pharmacokinetic profiles like increased bioavailability and faster absorption.

In-vivo pharmacodynamic studies

Pharmacodynamic studies assess the *in-vivo* relationship between drug concentrations and therapeutic effects. The application of hydrotropic agents can result in more predictable and reproducible pharmacodynamic responses due to improved solubility and absorption.[53]

Clinical significance

Hydrotropic agents have an enormous application in the development of pharmaceutical formulations and, more particularly, for pharmaceuticals with poor solubility in water. It will help in developing formulations that will maximize drug delivery, improve patient compliance, and enhance therapeutic outcomes.[54]

Clinical Translation and Market Preservative

One of the major problems in pharmaceutical formulation is to increase the solubility of weakly water-soluble medicaments; hydrotropic agents play a key role in this regard. The term "hydrotropy" designates the phenomenon where some compounds, called hydrotropic agents, enhance the solubility of poorly water-soluble drugs in aqueous solutions. This process is based on the complex structural formation between the hydrotropic agent and drug molecules that gives rise to enhanced solubility. Applications of hydrotropic agents affect markets and therapeutic settings.

Clinical Translation

Improved bioavailability

Poorly water-soluble drugs generally have low bioavailability due to inadequate absorption. Hydrotropic agents enhance the solubility, which aids in better absorption and hence improves the bioavailability of the drug.

Enhanced therapeutic efficacy

In general, higher solubility tends to result in more consistent and reliable drug concentrations, thereby probably increasing the overall therapeutic efficiency by improving its bioavailability at an appropriate site of action.

Dose reduction

Enhanced solubility can have the clinical benefit of reduced dosing while maintaining therapeutic efficacy, which minimizes side effects and improves patient compliance.

Formulation flexibility

These hydrotropic agents can give the formulator more flexibility in designing a formulation. This is especially important in oral formulations, wherein the ability to solubilize quite often influences absorption.[55]

Market Perspective

Expanded market potential

This enhances the solubility of poorly water-soluble drugs, hence opening opportunities for the development and marketing of drugs that were otherwise considered tough because of solubility issues.

Life cycle management

To sum up, the re-formulation of existing drugs, which are facing patent expiry, using hydrotropic agents can be considered a cycle time extension strategy for such a product, either by means of increased efficacy or through the introduction of new dosage forms.

Competitive advantage

Companies applying novel solubilization technologies, of which hydrotropic agents form one, are better placed to command product differentiation in the market—important in developing competitive advantage. This can become particularly relevant for the generic drug markets.

Market share and patient preference

Products with increased solubility and bioavailability will get more preference in the market, leading to an increased market share. Patients are also more attracted to formulations that give better efficacy and potentially fewer side effects.[56]

CONCLUSION

The development of hydrotropic solubilization technology has dramatically improved the problem of the low water solubility of medications. It offers a valid route to the improvement of therapeutic formulations by increasing drug solubility and bioavailability with the aid of hydrotropic agents. This paper is able to outline basic processes and variables impacting medication solubility by doing an in-depth review of the basic composition, history, and types of hydrotropes. One has to optimize the concentrations of hydrotropic agents and synergistic effects on solubilization in order to derive maximum advantages through the technique. Computational modeling and QSAR studies on the solubilizing characteristics of hydrotropic systems have created a better understanding and underlined its potential use in the development of drugs. The clinical significance of hydrotropes is further put forward by the capability to improve *in-vivo* pharmacokinetics and pharmacodynamics while letting go of the biological obstacles.

REFERENCES

1. Fatouh AM, Elshafeey AH, Abdelbary A. Intranasal agomelatine solid lipid nanoparticles to enhance brain delivery: formulation, optimization and *in-vivo* pharmacokinetics. Drug Design, Development and Therapy. 2017;11:1815-1825.DOI: 10.2147/DDDT.S102500
2. Ahmed EM. Hydrogel: Preparation, characterization, and applications: A review. Journal of Advanced Research. 2015;6(2):105-121. DOI: 10.1016/j.jare.2013.07.006
3. Kuchekar AB, Gawade A, Boldhane S. Hydrotropic solubilization: an emerging approach. Journal of Drug Delivery and Therapeutics. 2021;11(1-s):200-206. DOI: 10.22270/jddt.v11i1-s.4724
4. Badwan AA, El-Khordagui LK, Saleh AM, Khalil SA. The solubility of benzodiazepines in sodium salicylate solution and a proposed mechanism for hydrotropic solubilization. International Journal of Pharmaceutics. 1982;13(1):67-74. DOI: 10.1016/0378-5173(82)90143-0
5. Khadka P, Ro J, Kim H, Kim I, Kim JT, Kim H, Cho JM, Yun G, Lee J. Pharmaceutical particle technologies: An approach to improve drug solubility, dissolution and bioavailability. Asian Journal of Pharmaceutical Sciences. 2014;9(6):304-316. DOI: 10.1016/j.ajps.2014.05.005
6. Ghadi R, Dand N. BCS class IV drugs: Highly notorious candidates for formulation development. Journal of Controlled Release. 2017;248:71-95. DOI: 10.1016/j.jconrel.2017.01.014
7. Mehringer J, Kunz W. Carl Neuberg's hydrotropic appearances (1916). Advances in Colloid and Interface Science. 2021;294:102476. DOI: 10.1016/j.cis.2021.102476
8. Kumar VS, Raja C, Jayakumar C. A review on solubility enhancement using hydrotropic phenomena. International Journal of Pharmacy and Pharmaceutical Sciences. 2014;6(6):1-7. ISSN- 0975-1491
9. Li P, Zhao L. Developing early formulations: practice and perspective. International Journal of Pharmaceutics. 2007;341(1-2):1-9. DOI: 10.1016/j.ijpharm.2007.05.049
10. Saleh AM, El-Khordagui LK. Hydrotropic agents: a new definition. International Journal of Pharmaceutics. 1985;24(2-3):231-238. DOI: 10.1016/0378-5173(85)90023-7
11. Lee J, Lee SC, Acharya G, Chang CJ, Park K. Hydrotropic solubilization of paclitaxel: analysis of chemical structures for hydrotropic property. Pharmaceutical Research. 2003;20:1022-1030. DOI: 10.1023/A:1024458206032
12. Sanghvi R, Evans D, Yalkowsky SH. Stacking complexation by nicotinamide: A useful way of enhancing drug solubility. International Journal of Pharmaceutics. 2007;336(1):35-41. DOI: 10.1016/j.ijpharm.2006.11.025
13. Agrawal S, Pancholi SS, Jain NK, Agrawal GP. Hydrotropic solubilization of nimesulide for parenteral administration. International Journal of Pharmaceutics. 2004;274(1-2):149-155. DOI: 10.1016/j.ijpharm.2004.01.012
14. Suzuki H, Sunada H. Mechanistic studies on hydrotropic solubilization of nifedipine in nicotinamide solution. Chemical and Pharmaceutical Bulletin. 1998;46(1):125-130. DOI: 10.1248/cpb.46.125
15. Rasool AA, Hussain AA, Dittert LW. Solubility enhancement of some water-insoluble drugs in the presence of nicotinamide and related compounds. Journal of Pharmaceutical Sciences. 1991;80(4):387-393. DOI: 10.1002/jps.2600800422
16. Badwan AA, El-Khordagui LK, Saleh AM, Khalil SA. The solubility of benzodiazepines in sodium salicylate solution and a proposed mechanism for hydrotropic solubilization. Journal of Pharmacy and Pharmacology. 1980;32(Supplement_1):74P-74P. DOI: 10.1111/j.2042-7158.1980.tb10877.x
17. Poochikian GK, Cradock JC. Enhanced chartreusin solubility by hydroxybenzoate hydrotropy. Journal of Pharmaceutical Sciences. 1979;68(6):728-732. DOI: 10.1002/jps.2600680620
18. Coffman RE, Kildsig DO. Hydrotropic solubilization—mechanistic studies. Pharmaceutical Research. 1996;13:1460-1463. DOI:10.1023/A:1016011125302
19. Beig A, Lindley D, Miller JM, Agbaria R, Dahan A. Hydrotropic solubilization of lipophilic drugs for oral delivery: the effects of urea and nicotinamide on carbamazepine solubility–permeability interplay. Frontiers in Pharmacology. 2016;7:379. DOI: 10.3389/fphar.2016.00379
20. Kim JY, Kim S, Papp M, Park K, Pinal R. Hydrotropic solubilization of poorly water-soluble drugs. Journal of Pharmaceutical Sciences. 2010;99(9):3953-3965. DOI: 10.1002/jps.22241
21. Martins AC, Benfica J, Peréz-Sanchéz G, Shimizu S, Sintra TE, Schaeffer N, Coutinho JA. Assessing the hydrotropic effect in the presence of electrolytes: competition between solute salting-out and salt-induced hydrotrope aggregation. Physical Chemistry Chemical Physics. 2022;24(36):21645-21654. DOI:10.1039/D2CP00749E
22. Shimizu S, Matubayasi N. Hydrotropy: Monomer–micelle equilibrium and minimum hydrotrope concentration. The Journal of Physical Chemistry B. 2014;118(35):10515-10524. DOI: 10.1021/jp505869m
23. Srinivas V, Balasubramanian D. When does the switch from hydrotropy to micellar behavior occur? Langmuir.

1998;14(23):6658-6661. DOI: 10.1021/la980598c
24. Soares B, Silvestre AJ, Rodrigues Pinto PC, Freire CS, Coutinho JA. Hydrotropy and cosolvency in lignin solubilization with deep eutectic solvents. ACS Sustainable Chemistry & Engineering. 2019;7(14):12485-12493 DOI: 10.1021/acssuschemeng.9b02109
25. Lee SC, Acharya G, Lee J, Park K. Hydrotropic polymers: Synthesis and characterization of polymers containing picolylnicotinamide moieties. Macromolecules. 2003;36(7):2248-2255. DOI: 10.1021/ma021629k
26. Neumann MG, Schmitt CC, Prieto KR, Goi BE. The photophysical determination of the minimum hydrotrope concentration of aromatic hydrotropes. Journal of Colloid and Interface Science. 2007;315(2):810-813. DOI: 10.1016/j.jcis.2007.07.020
27. Dhapte V, Mehta P. Advances in hydrotropic solutions: An updated review. St. Petersburg Polytechnical University Journal: Physics and Mathematics. 2015;1(4):424-435. DOI: 10.1016/j.spjpm.2015.12.006
28. Molinier V, Aubry JM. Sugar-based hydrotropes: preparation, properties and applications. DOI: 10.1039/9781849739986-00051
29. Lieberman H, Rieger M, Banker GS, editors. Pharmaceutical Dosage Forms: Disperse Systems. CRC Press; 2020.
30. Paul BK, Moulik SP. Microemulsions: An overview. Journal of Dispersion Science and Technology. 1997;18(4):301-367. DOI: 10.1080/01932699708943740
31. Peracchia MT. Colloidal drug delivery systems. In: Kreuter J, editor. Marcel Dekker; 1994.
32. Tenjarla S. Microemulsions: An overview and pharmaceutical applications. Critical Reviews™ in Therapeutic Drug Carrier Systems. 1999;16(5). DOI: 10.1615/CritRevTherDrugCarrierSyst.v16.i5.20
33. Rahman A, Haider MD, Naseem N, Rahman N. Solubility of Drugs, Their Enhancement, Factors Affecting and Their Limitations: A Review. International Journal of Pharmaceutical Sciences Review and Research. 2023;79:78-94. DOI: 10.47583/ijpsrr.2023.v79i02.014
34. Maheshwari R, Singh M. Quantitative Determination of Ketoprofen Bulk Drug Using Sodium Salt of Aspirin as Hydrotropic Solubilizing Agent. ajc [Internet]. 2010 Mar. 30;20(6):4922-4.Available from: https://asianpubs.org/index.php/ajchem/article/view/19118.
35. Patil SS, More AS. A Review on Hydrotropy: The Approach Towards Green Chemistry. DOI: 10. 20959/wjpr202212-25303
36. Sharma P, Kaur T, Singh AP. Solubility and dissolution enhancement of water insoluble drug by using different hydrophilic carriers and formulated into tablet dosage form. Journal of Drug Delivery and Therapeutics. 2017;7(5):86-93. DOI: 10.22270/jddt.v7i5.149
37. Maheshwari RK, Sharma S, Rai N, Rajput M. Simple titrimetric method to estimate ketoprofen in bulk using mixed hydrotropy. Journal of Pharmacy Research. 2010;3(3):442-443. ISSN: 0974-6943
38. Rani TN, Muzib YI. Formulation and *in-vivo* evaluation of rosuvastatin rapid melts by direct compression and sublimation methods. International Journal of Drug Delivery Technology. 2018;8:212-221. DOI: 10.25258/ijddt.8.4.8
39. Nainwal P, Sinha P, Singh A, Nanda D, Jain DA. A comparative solubility enhancement study of rosuvastatin using solubilization techniques. ISSN 0976-4550
40. Butt S, Hasan SM, Hassan MM, Alkharfy KM, Neau SH. Directly compressed rosuvastatin calcium tablets that offer hydrotropic and micellar solubilization for improved dissolution rate and extent of drug release. Saudi Pharmaceutical Journal. 2019;27(5):619-628. DOI: 10.1016/j.jsps.2019.03.002
41. Ramu A, Vidyadhara S, Devanna N, Anusha C, Keerthi J. Formulation and evaluation of rosuvastatin fast dissolving tablets. Asian Journal of Chemistry. 2013;25(10):5340. DOI: 10.14233/ajchem.2013.14180
42. González R, Peña MÁ, Torres NS, Torrado G. Design, development, and characterization of amorphous rosuvastatin calcium tablets. PLoS ONE. 2022;17(3):e0265263. DOI: 10.1371/journal.pone.0265263
43. Gupta A. Computational intelligence in time series forecasting: theory and engineering applications. In: Palit AK, Popovic D, editors. Springer; 2005. DOI: 10.1002/rnc.1153
44. Adcock SA, McCammon JA. Molecular dynamics: survey of methods for simulating the activity of proteins. Chemical Reviews. 2006;106(5):1589-1615. DOI: 10.1021/cr040426m
45. Paul R, Chattaraj KG, Paul S. Role of hydrotropes in sparingly soluble drug solubilization: insight from a molecular dynamics simulation and experimental perspectives. Langmuir. 2021;37(16):4745-4762. DOI: 10.1021/acs.langmuir.1c00169
46. Lounnas V, Pettitt BM. A connected-cluster of hydration around myoglobin: Correlation between molecular dynamics simulations and experiment. Proteins: Structure, Function, and Bioinformatics. 1994;18(2):133-147. DOI: 10.1002/prot.340180206
47. Isarankura-Na-Ayudhya C, Naenna T, Nantasenamat C, Prachayasittikul V. A practical overview of quantitative structure-activity relationship. DOI:10.17877/DE290R-690
48. Khan AD, Tabish M, Kaushik R, Saxena V, Kesharwani P, Gupta S, Alam MN, Sharma V. Hydrotropy: Recent advancements in enhancement of drug solubility and formulation development. International Journal of Drug Delivery Technology. 2021;11. DOI: 10.25258/ijddt.11.3.47
49. Zakharova LY, Vasilieva EA, Mirgorodskaya AB, Zakharov SV, Pavlov RV, Kashapova NE, Gaynanova GA. Hydrotropes: Solubilization of nonpolar compounds and modification of surfactant solutions. Journal of Molecular Liquids. 2023;370:120923. DOI: 10.1016/j.molliq.2022.120923
50. Ahire E, Thakkar S, Darshanwad M, Misra M. Parenteral nanosuspensions: a brief review from solubility enhancement to more novel and specific applications. Acta Pharmaceutica Sinica B. 2018;8(5):733-755. DOI: 10.1016/j.apsb.2018.07.011
51. Wanat K. Biological barriers, and the influence of protein binding on the passage of drugs across them. Molecular Biology Reports. 2020;47(4):3221-3231. DOI: 10.1007/s11033-020-05361-2
52. Smith D, Artursson P, Avdeef A, Di L, Ecker GF, Faller B, Houston JB, Kansy M, Kerns EH, Kramer SD, Lennernäs H. Passive lipoidal diffusion and carrier-mediated cell uptake are both important mechanisms of membrane permeation in drug disposition. Molecular Pharmaceutics. 2014;11(6):1727-1738. DOI: 10.1021/mp400713v
53. Forrest LR, Krämer R, Ziegler C. The structural basis of secondary active transport mechanisms. Biochimica et Biophysica Acta (BBA) - Bioenergetics. 2011;1807(2):167-188. DOI: 10.1016/j.bbabio.2010.10.014
54. Stupp SI, Son S, Lin HC, Li LS. Synthesis of two-dimensional polymers. Science. 1993 Jan 1;259(5091):59-63. doi: 10.1126/science.259.5091.59. PMID: 17757473.
55. Battey NH, James NC, Greenland AJ, Brownlee C. Exocytosis and endocytosis. The Plant Cell. 1999;11(4):643-659. DOI: 10.1105/tpc.11.4.643
56. Bhalani DV, Nutan B, Kumar A, Singh Chandel AK. Bioavailability enhancement techniques for poorly aqueous soluble drugs and therapeutics. Biomedicines. 2022;10(9):2055. DOI: 10.3390/biomedicines10092055