

The Relationship Between Blood Group Types and Haemoglobin Levels in Bihar's Rural Population

Bharat Kumar¹, Sunil Kumar², Richa Kumari³, Malti Kumari⁴

¹Tutor, Department of Physiology, Jan Nayak Karpoori Thakur Medical College and Hospital, Madhepura, Bihar, India

²Tutor, Department of Physiology, Jan Nayak Karpoori Thakur Medical College and Hospital, Madhepura, Bihar, India

³Senior Resident, Department of Physiology, AIIMS, Rishikesh, Uttarakhand, India.

⁴Professor and HOD, Department of Physiology, Jan Nayak Karpoori Thakur Medical College and Hospital, Madhepura, Bihar, India

Received: 06-05-2021 / Revised: 02-06-2021 / Accepted: 18-07-2021

Corresponding author: Dr. Sunil Kumar

Conflict of interest: Nil

Abstract

Aim: Study of association between blood groups and blood hemoglobin levels in rural Population of Bihar region. **Methods:** The present Prospective study was conducted in the Department of Physiology, Jan Nayak Karpoori Thakur Medical College and Hospital, Madhepura, Bihar, India from Jan 2020 to December 2020. 100 adults (males and females) who are in the age group of 18-30 years were included in this study. Blood sample is taken using finger prick, under aseptic conditions and blood group is determined using glass slide method using antisera A, B and D. Hemoglobin concentration is estimated using Sahli's method. **Results:** In this study, adults with hemoglobin concentration less than 10g/dl were taken as anemic. Total 30 adults are anemic, while remaining 70 adults are non-anaemic. 41 adults are females out of which 59 are anemic whereas 59 adults are males where only 10 males were found to be anemic. 29 adults with Hb less than 10g/dl were Rh positive and 1 adult Rh negative. 67 adults and 3 adults were Rh positive and Rh negative respectively with Hb more than 10g/dl. Also 27 adults are blood group A, 5 with blood group AB, 35 with blood group B and O blood group adults are 33. Among the A blood group, 5 adults were anemic, among blood group B, 14 were anemic, among blood group AB, 2 were anemic and 9 adults with O blood group were anemic. **Conclusion:** Individuals with blood group B are more prone to anaemia followed by blood group O, AB and least is with blood group A. Based on their blood groups, we can advise regular intake of diet rich in iron and vitamins or also their supplements to the individual who are more susceptible to anaemia.

Keywords: Blood Groups, Anaemia, Hb.

This is an Open Access article that uses a fund-ing model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

Introduction

Blood is a specialized connective tissue with complete and unchangeable identity. It

provides one of the means of connection between the cells of different parts of the

body and external environment[1]. In modern medicine blood transfusion is an important measure for replacing blood loss[2]. At least 30 commonly occurring antigens and hundreds of other rare antigens have been found in human blood cells, especially on the surfaces of the cell membranes. Most of the antigens are weak and therefore are of importance principally for studying the inheritance of genes to establish parentage. Two particular types of antigens are much more likely than the others to cause blood transfusion reactions. They are the A B O system of antigens and the Rh system[3]. ABO and Rh are recognized as the major clinically significant blood group antigens. The rhesus blood group system was the fourth system to be discovered[1]. The ABO blood group individuals are divided into four major blood groups, namely, A, B, AB and O, according to the presence of antigens and agglutinins. Blood group A blood has type A antigens, blood group B has type B antigens and blood group O has neither A nor B antigens. The Rhesus antigen (name because a related antigen was first discovered in Rhesus monkeys), is found in the red blood cells of approximately 85%-90% of the people. This is the second most important blood group system due to its immunogenicity in Rh negative individuals in blood transfusion or pregnancy. Rh positive individuals have RhD antigen on the surface of their red blood cells and Rh-negative individuals do not have this antigen. Blood group antigens are not only important in relation to blood transfusion and organ transplantation, but also have some association with diseases like duodenal ulcer, diabetes mellitus, urinary tract infection and Rh incompatibility of the newborn[1]. As per the World Health Organization (WHO) database on anemia globally, anemia affects 1.62 billion people (95%), which corresponds to 24.8% of the population. WHO also estimates that anemia contributes to about 20% of maternal and perinatal death in developing countries[4,5]. Anaemia can be defined by a condition in which the total haemoglobin

(Hb) level or number of red blood cells (RBCs) is poorly lowered. The World Health Organisation (WHO) defines anaemia as Hb < 130 g/L in men older than 15 years, 110 g/L in pregnant women, and < 120 g/L in non-pregnant women older than age 15 years[6,7]. Anemia is defined as a clinical condition which is characterized by reduction in hemoglobin concentration of the blood below the normal for the age.² It is a global problem, mainly affecting poor people in developing countries[2,3]. Anemia during adolescence severely impairs the physical and mental development; weakens behavioral & cognitive development; reduces physical fitness; decreases the work performance and even contributes to the adverse pregnancy outcome[1]. Though, the oral iron supplements are given commonly to correct anemia but should be used only when dietary measures have failed. Moreover, iron supplement can correct anemia which is due to iron deficiency[4]. Besides this, oral iron supplements can cause gastrointestinal problems like nausea, vomiting, diarrhea and constipation and can even aggravate pre-existing ulcers and ulcerative colitis. Long term iron supplements also can cause heart disease. Anaemia even can have a negative impact on physical performance, due to reduced oxygen transport the reduced cellular oxidative capacity[6,8].

Material and methods

The present Prospective study was conducted in the Department of Physiology, Jan Nayak Karpooori Thakur Medical College and Hospital, Madhepura, Bihar, India from January 2020 to December 2020.

Methodology

100 adults (males and females) who are in the age group of 18-30 years were included in this study. Blood sample is taken using finger prick, under aseptic conditions and blood group is determined using glass slide method using antisera A, B and D. Hemoglobin concentration is estimated

using Sahli's method. The brief of the study and procedure is given to the subjects and informed consent taken.

Results

In our study, adults with hemoglobin concentration less than 10g/dl were taken as anemic. Total 30 adults are anemic, while remaining 70 adults are non anaemic. 41 adults are females out of which 59 are anemic whereas 59 adults are males where only 10 males were found to be anemic. 29 adults with Hb less than 10g/dl were Rh positive and 1 adult Rh negative. 67 adults and 3 adults were Rh positive and Rh

negative respectively with Hb more than 10g/dl. Also 27 adults are blood group A, 5 with blood group AB, 35 with blood group B and O blood group adults are 33. Among the A blood group, 5 adults were anemic, among blood group B, 14 were anemic, among blood group AB, 2 were anemic and 9 adults with O blood group were anemic.

In our study we found that adults with blood group B (14%) are more prone for anemia, followed by O (9%), AB (2%) and blood group A (5%). Blood group A almost resistant to anemia.

Table No. 1: Distribution of anaemic and non-anaemic adults

Hb	Frequency	Percentage
<10	30	30
>=10	70	70
Total	100	100

Table No. 2: Anaemic status in males and females

Sex	Hb		Total
	<10	>=10	
Female	20	21	41
Male	10	49	59
Total	30	70	100

$$\chi^2 = 51.70, p < 0.0001$$

Table No. 3: Different blood group status in anaemic and non-anaemic adults

Blood group	Hb		Total
	<10	>=10	
A +ve	5	20	25
A -ve	0	2	2
AB +ve	1	3	4
AB -ve	1	0	1
B -ve	0	1	1
B +ve	14	20	34
O +ve	9	24	33
Total	30	70	100

$$\chi^2 = 7.16, p = 0.17$$

Discussion

In our study, the distribution of blood group B was the highest with a percentage frequency of 35%, followed by blood group O and A with a percentage frequency of 33 and 27% respectively and the least

percentage frequency is that of blood group AB which is 5%. The prevalence of anaemia in our study was found in blood group B, followed by O, AB and then A. The same trend of prevalence of blood groups (B>O>A>AB) was observed and reported by study conducted by Kaur M[1].

In many other studies, blood group O has been found to be the most common blood group. The frequency of Rh+ve was about 96%, while 4% were Rh-ve in our study. Similar patterns of distribution were also observed in other studies. Thus, the frequencies of ABO and rhesus blood groups vary from one population to another. On further evaluating the distribution of subjects according to blood groups and their relationship to anemia it was found that anemia was more frequent in blood group B (40%) followed by blood group AB (40%), A (18.51%) and least in the blood group O (27.27%) but it was not statistically significant ($p=0.136$). Similar findings were seen in the study conducted by Basak Asim Kumar[4], in which individuals having blood group B, A or AB were prone to anemia compared to blood group O but was statistically significant. On the other hand, there was no such relationship between Rh factor and the occurrence of anemia between the above-mentioned populations. Anemia during adolescence severely impairs the physical and mental development; weakens behavioral and cognitive development; reduces physical fitness; decreases the work performance and even contributes to the adverse pregnancy outcome Mild anemia can adversely affect the productivity and is also known to reduce the immune competence. The average prevalence of anemia is 25-80% seen in several studies across the Indian subcontinent in a study conducted by ICMR in sixteen districts in eleven different states reporting a prevalence rate of anemia 90.1% among the adolescent girls of 11-18 years age groups. In their study, 35% anemia was seen in the 300 adults studied. In further evaluating gender wise it was found that anemia was significantly more in the girls (66.67%) compared to the boys (33.33%) with a 'p' value of 0.0001. The similar higher prevalence of anemia was found in the rural girls (98%) and boys (56%) of Punjab. This study reveals that there is a relationship between blood group and anemia, though not statistically significant which could be

due to small sample size. Similar finding was found in our study where girls are more prone to anaemia. The individuals with blood group antigen alpha and beta are comparatively more prone to be anemia, whereas the individuals devoid of these antigens are resistant to anemia. So, the regular intake of iron and vitamin rich diet in individuals having blood groups A, B, and AB can prevent the occurrence of anemia.

Conclusion

We conclude that individuals with blood group B are more prone to anaemia followed by blood group O, AB and least is with blood group A. Based on their blood groups, we can advice regular intake of diet rich in iron and vitamins or also their supplements to the individual who are more susceptible to anaemia.

Reference

1. Kaur M. Relationship and Distribution of ABO, Rh Blood Groups and Hemoglobin Concentration among the Adolescents J Phys Pharm Adv 2015, 5 (9): 703-712.
2. Reshmarani, Shilpa N., Subhash Chimkode A Study of Correlation Between Blood Groups and Anemia in Young Adults, International Journal of Physiology, 2019; 7(4): 199-202.
3. Guyton and Hall, Textbook of Medical Physiology; Second South Asia Edition July 2016;201-202.
4. BasakAsim Kumar and Maji Kaushik: Blood group and anemia: Exploring a new relationship, Journal of Public Health and Epidemiology 2013;5(1):43-45.
5. Harvey P (2004). A strategic approach to anemia control, MOST, The USAID micronutrients program, Arlington, Virginia. <http://health.nytimes.com/health/guides/disease/anemia/treatment.html> retrieved on 12.02.12.
6. Salma AlDallal, Iron Deficiency Anemia: A Short Review, Journal of Cancer Research and ImmunoOncology, 2016; 2(1):3-6.

7. Goddard AF, James MW, McIntyre AS, Scott BB. Guidelines for the management of iron deficiency anaemia. *Gut* 2011;60: 1309-1316.
8. Zhang AS, Enns CA. Molecular mechanisms of normal iron homeostasis. *Hematology. Am Soc Hematol Educ Program* 2009;1: 207-214.