

## Questionnaire-based Research to Determine the Level of Knowledge of Diabetic Retinopathy among People with type 2 Diabetics

Ganesh Kumar

Assistant Professor, Department of Ophthalmology, Lord Buddha Koshi Medical College & Hospital, Saharsa, Bihar, India

---

Received: 03-07-2021 / Revised: 18-08-2021 / Accepted: 28-10-2021

Corresponding author: Dr. Ganesh Kumar

Conflict of interest: Nil

---

### Abstract

**Aim:** Awareness of diabetic retinopathy among Type 2 diabetes mellitus patients in Bihar region.

**Methods:** This was a descriptive, cross sectional, non-randomized, questionnaire-based study conducted in the Department of Ophthalmology, Lord Buddha Koshi Medical College & Hospital, Saharsa, Bihar, India, for 15 months, Participants were asked to answer questions from a structured questionnaire developed in English and Hindi, which included questions about awareness of DR due to DM and compliance with DM and DR management. All interviewed patients were with Type 2 DM and were randomly selected using multistage random cluster sampling from the general population in and around Bihar. 200 study subjects having Type 2 diabetes mellitus were selected. 200 Patients with Type 2 diabetes mellitus who had normal cognitive ability and could speak Hindi and resided in and around Bihar was selected.

**Result:** Of 200 randomly selected Type 2 diabetes mellitus patients, 78% of these people had awareness about diabetes mellitus causing a condition called diabetic retinopathy. But only 36% knew that Diabetic Retinopathy can cause blindness. 86% of them are aware that Diabetic retinopathy can be prevented. Only 15% are aware that laser treatment for DR does not improve vision but reduces further deterioration in vision. 78% feel regular eye check-ups are required in DM patients and nearly 78% patients are fairly frequent with their eye check-up. 80% of people are aware that maintenance of blood sugar levels can reduce the risk of diabetic retinopathy and 72% check their blood sugar levels regularly while 28% are irregular with blood glucose checking. 94% of the patients have a source of information about diabetes mellitus and diabetic retinopathy with only 6% answered they do not receive any information. Only 33% of people are aware that a diabetic patient should first see an eye doctor at the time of diagnosis.

**Conclusion:** There is a need to implement strategies to increase the awareness of DR and the importance of early retinal screening among affected patients, in order to reduce the risk of visual complications.

**Keywords:** diabetic retinopathy, diabetes mellitus, awareness

---

This is an Open Access article that uses a fund-ing model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

---

## Introduction

It has been estimated that there are about 69.2 million Indians suffering from diabetes[1]The increasing prevalence of diabetes and the increased life expectancy; raises the time period for which patients have been suffering from diabetes; this increases the risk of long term complications of diabetes, a very important one being diabetic retinopathy. The overall global prevalence of diabetic retinopathy is 34.6% and it accounts for 4.8% of blindness in the world[2,3] The prevalence of diabetic retinopathy in India from various studies range from 7.3% to 20%[4,7] Wisconsin Epidemiological study has proved that micro vascular complications such as diabetic retinopathy are linked to duration of diabetes[8].Routine dilated fundus examination is recommended at the time of diagnosis of diabetes and then yearly review is required in all patients having type 2 diabetes mellitus. Awareness of importance of routine check-up for the screening of diabetic retinopathy is poor even in developed countries and the situation is much worse in a developing country like India. Previous studies have shown that 63% of the rural diabetic population has not had an eye examination[9]Magnitude of blindness caused by diabetic retinopathy is increasing. Lack of awareness about diabetic retinopathy and the preventable complications associated with it worsens the situation .Early detection of diabetic retinopathy and its appropriate management is very important to prevent irreversible visual loss. This can only be achieved with better knowledge and awareness among patients. There have been many studies done in other states in India to assess the knowledge and awareness on diabetes and diabetic retinopathy among patients with diabetic retinopathy, however very few studies assessed knowledge, awareness and practices amongst patients with diabetic retinopathy[10,13]The objective of the study was to assess the

knowledge, attitudes and practices about diabetes and diabetic retinopathy among patients diagnosed with diabetic retinopathy at a tertiary care hospital in bihar.

## Material and Methods

This was a descriptive, cross sectional, non-randomized, questionnaire-based study conducted in the Department of Ophthalmology, Lord Buddha Koshi Medical College & Hospital, Saharsa, Bihar, India, for 15 months,

## Methodology

Participants were asked to answer questions from a structured questionnaire developed in English and hindi, which included questions about awareness of DR due to DM and compliance with DM and DR management.

All interviewed patients were with Type 2 DM and were randomly selected using multistage random cluster sampling from the general population in and around Bihar. 100 study subjects having Type 2 diabetes mellitus were selected.

For the assessment of the reliability of the questionnaire, a random sample of 20 patients with diabetes was recruited to complete the questionnaire and the obtained Cronbach's alpha value was 0.8. Based on the preliminary results which showed a redundancy in some questions, the questionnaire was amended, and these results were disregarded in the final data analysis. 200 Patients with Type 2 diabetes mellitus who had normal cognitive ability and could speak hindi and resided in and around Bihar were selected. All non-diabetic patients and any patient with other type of Diabetes were excluded from the study.

## Data collection

The Questionnaire had two sections:

One was the basic sociodemographic information that included age, gender, educational level, whether they wear glasses or not and address. The second section was to assess the knowledge about diabetes, DR and source(s) of knowledge; patient's compliance with DM control, treatment and routine eye check-up visits. A sample of the questions related to this section is given below:

1. Are you aware that Diabetes Mellitus can affect the retina of your eyes and cause a condition called Diabetic Retinopathy?
2. Are you aware that diabetic retinopathy can lead to blindness?
3. Do you think regular eye check-ups are required in diabetic patients?
4. Do you think maintenance of blood sugar levels can reduce the risk of diabetic retinopathy?
5. How frequently do you think you should undergo eye check-ups?
6. How frequently do you check your blood glucose levels?
7. What are your sources of information about diabetes mellitus and diabetic retinopathy?

8. Do you know that laser treatment for diabetic retinopathy does not improve vision but reduces further deterioration in vision?

9. Do you think diabetic retinopathy can be prevented?

10. When do you feel that a diabetic should first see an eye doctor?

Informed consent was obtained from all individual participants included in the study prior to their participation in the study.

#### Data analysis

Numbers and percentages were calculated to summarize categorical and nominal data. According to assigned key, awareness present or absent is calculated.

#### Result

Of 200 randomly selected Type 2 diabetes mellitus patients, 111(55.5%) were women and 89 (44.5%) were males. 25.5% of them were illiterates, 27% had only primary education and 33% had education above middle school. 66% of the selected patients wore glasses for refractive error and they were aged from 18 yrs to 75 yrs

**Table 1: Demographic characteristics of the study population**

Variables	Number	Percentage
<b>Age</b>		
Below 30	21	10.5
30-40	45	22.5
40-50	70	35
50-60	40	20
Above 60	24	12
<b>Gender</b>		
Male	89	44.5
Female	111	55.5
<b>Educational status</b>		
Uneducated	51	25.5
class 1-12	120	60
college level	29	14.5
<b>Socio economic status</b>		
Lower	110	55
Middle	57	28.5
Upper	33	16.5

78% of these people had awareness about diabetes mellitus causing a condition called diabetic retinopathy. But only 36% knew that Diabetic Retinopathy can cause blindness. 86% of them are aware that Diabetic retinopathy can be prevented. Only 15% are aware that laser treatment for DR does not improve vision but reduces further deterioration in vision.

78% feel regular eye check-ups are required in DM patients and nearly 78% patients are fairly frequent with their eye check-up.

**Table 2:**

Question	Yes (Aware)	No (Not aware)
1. Are you aware that diabetes mellitus can affect the retina of your eyes and cause a condition called diabetic retinopathy?	78%	22%
2. Are you aware that diabetic retinopathy can lead to blindness?	36%	64%
3. Do you think regular eye check-ups are required in Diabetic patients?	78%	22%
4. Do you think maintenance of blood sugar levels can reduce the risk of diabetic retinopathy?	80%	20%
5. How frequently do you think you should undergo eye check-ups?	a. 34%	
	b. 34%	22%
	c. 10%	
6. How frequently do you check your blood glucose levels?	a. 4%	
	b. 4%	28%
	c. 64%	
7. What are your sources of information about diabetes mellitus and diabetic retinopathy?	a. 0%	
	b. 0%	6%
	c. 82%	
	d. 12%	
8. Do you know that laser treatment for diabetic retinopathy does not improve vision but reduces further deterioration in vision?	15%	85%
9. Do you think diabetic retinopathy can be prevented?	86%	4%
		10%
10. When do you feel that a diabetic should first see an eye doctor?	33%	15%
		42%
		10%

Question 6: 1) Weekly 2) Monthly 3) Not Regularly

Question 7: 1) Internet 2) Magazines 3) Doctor 4) Friends and Relatives.

Question 10: 1) Not necessary 2) When they need glasses 3) Only if they are referred 4) When their vision goes bad 5) At the time of diagnosis

80% of people are aware that maintenance of blood sugar levels can reduce the risk of

diabetic retinopathy and 72% check their blood sugar levels regularly while 28% are irregular with blood glucose checking.

94% of the patients have a source of information about diabetes mellitus and diabetic retinopathy with only 6% answered they do not receive any information.

Only 33% of people are aware that a diabetic patient should first see an eye doctor at the time of diagnosis.

Based on this data, while nearly 3/4<sup>th</sup> of this urban population are aware about Diabetic Retinopathy and feel blood glucose control and regular eye check-ups are necessary in reducing the risk of Diabetic Retinopathy, less than 1/4<sup>th</sup> know about blindness caused by DR, that laser treatment does not improve the vision but only reduces further deterioration and when a Diabetic patient should first visit the eye doctor- therein suggestive that most are aware of the disease but are not well informed about the complications and treatment of the disease.

### Discussion

It is well known that awareness is a vitally important step in the creation of a successful program to battle against any disease in the community. This is especially true of the growing problem of diabetic retinopathy. Many studies have revealed that Diabetic Retinopathy, despite its status as one of the greatest causes of blindness in both developed and developing countries, is virtually unknown to a large majority of the population. The lack of awareness about DR is considered a major health problem that could interfere with proper management and prevention of possible visual impairment.

Of 200 randomly selected Type 2 diabetes mellitus patients, 111(55.5%) were women and 89 (44.5%) were males. 25.5% of them were illiterates, 27% had only primary education and 33% had education above middle school. 66% of the selected patients wore glasses for refractive error and they were aged from 18 yrs to 75 yrs. similar to another study.[14]

From the study we conducted, it can be concluded that a high level of awareness among more than 3/4ths of the population about diabetic retinopathy is present but only 1/4<sup>th</sup> is aware about its complications and treatment and management of DM to prevent DR. Despite the high level of awareness about DR in the study, the level of patients' compliance with check-ups and efforts to reduce risk of DR is low[15] This discrepancy between the levels of

awareness and compliance in terms of routine eye examination seems to be common among patients with diabetes in the world, coinciding with reports stating that only half of the patients in Myanmar[16] and two-thirds of Japanese patients attended a routine eye examination.

Poorly controlled diabetes raises the likelihood of complications like DR. This higher proportion of diabetics with poorly controlled diabetes may be attributed to lack of awareness and limited eye health care facilities. Information given to diabetic patients should not just be on the nature of ocular complications of diabetes, but also on the risk factors for these complications and how to prevent them. Proper education also plays a major role.

The main source of information about DR in our sample was doctors, followed by friends and relatives and almost negligible from mass media which suggests awareness through mass media such as newspapers or advertisements should be encouraged so that even people who don't have access to Internet get notified and become more aware.

The reason given by patients for not getting an early DR screening was mainly a lack of information about DR, lack of time and will to take the examination and a fear of discovering any eye diseases which may burden the financial status of their families. This finding warrants further investigation into how to encourage patients with diabetes to routinely comply with vision examinations and retinal assessments every 12 months, as recommended by the international guidelines.

DR complications become more severe with prolonged duration of diabetes; it is imperative that spreading awareness on DR should be adopted by the practitioners at early stages of DM. Furthermore, public health awareness campaigns need to follow well-planned strategies. As part of the community may not be able to pursue higher education, school programs need to be revised to increase awareness of non-

communicable diseases such as DM and also awareness programmes about DM and DR and eye camps have to be conducted even in remote places by medical personnel and students.

### Conclusion

There is a need to implement strategies to increase the awareness of DR and the importance of early retinal screening among affected patients, in order to reduce the risk of visual complications.

### Reference

1. International Diabetes Federation. IDF Diabetes Atlas. 7th edn. Brussels, Belgium: International Diabetes Federation; 2015
2. Yau JWY, Rogers SL, Kawasaki R, Lamoureux EL, Kowalski JW, Bek T, et al. Global prevalence and major risk factors of diabetic retinopathy. *Diabetes Care*. 2012;35(3):556-64.
3. Resnikoff S, Pascolini D, Etya'ale D, Kocur I, Pararajasegaram R, Pokharel GP, et al. Global data on visual impairment in the year 2002. *Bull World Health Organ*. 2004;82(11):844-51.
4. Rema M, Deepa R, Mohan V. Prevalence of retinopathy at diagnosis among type 2 diabetic patients attending a diabetic centre in South India. *Br J Ophthalmol*. 2000; 84:1058-60.
5. Rema M, Premkumar S, Anitha B, Deepa R, Pradeepa R, Mohan V. Prevalence of diabetic retinopathy in urban India: The Chennai Urban Rural Epidemiology Study (CURES) eye study, I. *Invest Ophthalmol Vis Sci*. 2005;46:2328-33.
6. Raman R, Rani PK, Reddi Racheppalle S, Gnanamoorthy P, Uthra S, Kumaramanickavel G, et al. Prevalence of diabetic retinopathy in India: Sankara Nethralaya diabetic retinopathy epidemiology and molecular genetics study report 2. *Ophthalmology*. 2009; 116:311-18.
7. Jonas JB, Nangia V, Khare A, Matin A, Bhojwani K, Kulkarni M, et al. Prevalence and associated factors of diabetic retinopathy in rural central India. *Diabetes Care*. 2013;36: e69.
8. Klein R, Klein B, Moss S. The Wisconsin Epidemiological Study of Diabetic Retinopathy: a review. *Diabetes Meta Review*. 1989; 5:5559-5570.
9. Padmaja KR, Rajiv R, Pradeep GP, Swati A, Kumaramanickavel G, Sharma T. Use of eye care services by people with diabetes-South India experience. *British J Ophthalmology*. 2005
10. Srinivasan NK, John D, Rebekah G, Kujur ES, Paul P, John SS. Diabetes and diabetic retinopathy: KAP among diabetes patients in a tertiary care centre. *J Clin Diag Res*. 2017; 11(7):1-7.
11. Hussain R, Rajesh B, Giridhar A, Gopalakrishnan M, Sadasivan S, James J, et al. Knowledge and attitudes about diabetes and diabetic retinopathy in suburban population of a South Indian state and its practice among patients with diabetes mellitus: A population-based study. *Indian J Ophthalmol*. 2016;64(4):272-6.
12. Rani PK, Raman R, Subramani S, Perumal G, Kumaramanickavel G, Sharma T. Knowledge of diabetes and diabetic retinopathy among rural population in India and its influence on knowledge of diabetic retinopathy on attitude and practice. *Rural and remote health*. 2008;8(3):838.
13. Koshy J, Varghese DL, Mathew T, Kaur G, Thomas S, Bhatti SM. Study on KAP of ocular complications due to diabetes among type 2 diabetes visiting a tertiary teaching hospital. *Indian J Community health*. 2012; 24:27-31.
14. Raman R, Rani PK, Racheppalle SR, Gnanamoorthy P, Uthra S, Kumaramanickavel G, et al. Prevalence of diabetic retinopathy in India. *Ophthalmol*. 2009; 116:311-8.

15. Rani PK, Raman R, Subramani S, Perumal G, Kumaramanickavel G, Sharma T. Knowledge of diabetes and diabetic retinopathy among rural populations in India, and the influence of knowledge of diabetic retinopathy on attitude and practice. *Rural Remote Health*. 2008;8(3):838.
16. Muecke JS, Newland HS, Ryan P, Ramsay E, Aung M, Myint S. Awareness of diabetic eye disease among general practitioners and diabetic patients in Yangon, Myanmar. *Clin Exp Ophthalmol*. 2008;36(3):265–73