

A Hospital Based Descriptive Assessment of Clinical and Hematological Profile in Dengue Fever

Shatrughan Kumar¹, MD. Tarique Anwar²

¹Department of Medicine, Specialist Medical Officer, Sadar Hospital, Motihari, Bihar, India

²Department of Medicine, Specialist Medical Officer, Sadar Hospital, Motihari, Bihar, India

Received: 25-09-2022 / Revised: 20-10-2022 / Accepted: 02-11-2022

Corresponding author: Dr. MD.Tarique Anwar

Conflict of interest: Nil

Abstract

Aim: This study is an attempt to elucidate the positive laboratory profile of serologically diagnosed dengue patients so as to facilitate early diagnosis, treatment, management and vector control measures, to reduce the morbidity and mortality because of this disease.

Methods: This was a descriptive study with analysis of patients who were admitted for dengue fever in the Department of Medicine, Sadar Hospital, Motihari, Bihar, India for duration of 6 months. This study was conducted on 100 indoor patients. Patients presenting to the emergency department, outpatient department (OPD) or pediatric OPD with complaints of fever and clinical features of dengue with positive NS1 antigen test or dengue antibody serology IgM or IgG or both were included in the study.

Results: Most of the cases were seen in the 20-30 years age group followed by 31-40 years age group. Majority of the patients were males compared to females and the male to female ratio was 2:1. Fever was the most common presentation and was seen in 40 cases (40%) cases. Present study showed hemoglobin range of 6 gm% to 17 gm% (Table 3). Raised hematocrit (>47%) was noted in 15 (15%) of patients at presentation and the hematocrit ranged from 20-51%. The total leukocyte count ranged from 1500 cells/cumm to >11000 cells/cumm. Leucopenia with less than 4000 cells/cumm was present in 25% cases. In the present study out of 100 cases of dengue fever, 90% cases had thrombocytopenia and 10% cases had severe thrombocytopenia (<20,000/cumm) with bleeding manifestations.

Conclusion: Hemoconcentration, leucopenia, thrombocytopenia, and raised liver enzymes SGOT and SGPT along with reactive/ plasmacytoid lymphocytes on peripheral smear gives enough clues to test for dengue serology so that dengue cases can be diagnosed in their initial stages. This facilitates early treatment and aggressive fluid replacement therapy with good nursing care so that fatality rates can be reduced.

Keywords: Dengue, Hemogram, Hematocrit, Leucopenia

This is an Open Access article that uses a fund-ing model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

Introduction

Dengue is an acute, rapidly spreading, self-limited disease (typically lasting 5–7 days), characterized by fever, prostration, headache, myalgia, rash,

lymphadenopathy, and leukopenia, and caused by four antigenically related, but distinct types of dengue viruses transmitted by the bite of infected

mosquitoes of the genus *Aedes*, especially *Aedes aegypti*, *Aedes albopictus*, and *Aedes polynesiensis*. It occurs epidemically and sporadically in India, Japan, West Africa, and Southeast Asia. [1]

An estimated 50 million dengue infections occur worldwide annually and approximately 2.5 billion people, i.e., two-fifth of world's population in tropical and subtropical countries are at risk. The reported case fatality rate in India is 3–5%. [2,3] Most developing countries have epidemics of febrile illnesses including typhoid, measles, leptospirosis, and severe acute respiratory distress syndrome that can be confused with dengue due to similar clinical features. The non-specific presentation underscores the importance of laboratory testing and a high index of suspicion to reduce the morbidity and mortality due to this disease. [2]

Dengue fever with warning signs (DFWS) and severe dengue (SD) with severe plasma leakage, severe bleeding or severe organ involvement have emerged as important public health threat in urban areas. This is attributable to population migration to cities resulting in urban overcrowding and infrastructure construction in these areas providing unhindered opportunities for breeding of the vector. [4]

The clinical presentation of DF is triphasic with the febrile phase typically characterized by high fever, headache, myalgia, body ache, vomiting, joint pain, transient rash and mild bleeding manifestations such as petichiae, ecchymosis at pressure sites and bleeding from venipunctures. [5] In the next critical phase there is a heightened risk of progression of the patient to severe dengue which is defined by presence of plasma leakage that may lead to shock and/or fluid accumulation such as ascites or pleural effusion with or without respiratory distress, severe bleeding, and/or severe organ impairment. [6] The risk of severe

bleeding in dengue is much higher with a secondary infection and is seen in about 2-4% of cases having secondary infection. [7-10] Atypical presentations are also encountered with acute liver failure, encephalopathy with seizures, renal dysfunction, lower gastrointestinal bleeding. [11]

Leukopenia is the most prominent hematological change, sometimes with counts of less than $2 \times 10^3/\mu\text{L}$. However, there are reports of mild leukocytosis at the onset of the disease, with neutrophilia. Lymphocytosis is a common finding, with the presence of atypical lymphocytes. The hematocrit concentration should be monitored according to the days of illness, remembering that, with the progression to DHF, there will be a 20% increase in hematocrit from the patient's baseline, associated with thrombocytopenia ($< 100 \times 10^9/\text{L}$). [12,13]

This study is an attempt to elucidate the positive laboratory profile of serologically diagnosed dengue patients so as to facilitate early diagnosis, treatment, management and vector control measures, to reduce the morbidity and mortality because of this disease.

Methods

This was a descriptive study with analysis of patients who were admitted for dengue fever in the Department of Medicine, Sadar Hospital, Motihari, Bihar, India for duration of 6 months. This study was conducted on 100 indoor patients. Patients presenting to the emergency department, outpatient department (OPD) or pediatric OPD with complaints of fever and clinical features of dengue with positive NS1 antigen test or dengue antibody serology IgM or IgG or both were included in the study. Age, gender, clinical presentation, duration of fever, dehydration, hemodynamic status, urine output, hepatomegaly, ascites, pleural effusion, presence of petechiae, positive tourniquet test, other bleeding manifestations,

hematocrit and platelet count were recorded at presentation.

Inclusion criteria

Inclusion criteria were febrile patients with positive NS1 antigen or IgM or both on rapid card tests. IgG may be positive or negative.

Exclusion criteria

Exclusion criteria were patients with only IgG positive on rapid card tests were excluded from the study. Patients with other identified illnesses like typhoid, malaria which were coexisted with dengue positive serology were excluded from the study.

Hemogram was done on automated cell counter analyzer (Sysmex XP 100) which included hemoglobin, hematocrit, total leucocyte count (TLC), differential leucocyte count (DLC) and platelets count.

Platelets counts were cross checked on stained smears. Hematocrit raised >20% of normal was considered as hemoconcentration. Leukopenia was taken as total leucocyte count <4,000/mm³. Thrombocytopenia was taken as platelets count <1,00,000/mm³.

Biochemical parameters included serum Aspartate aminotransferase (AST), Alanine aminotransferase (ALT), total bilirubin (T. Bil.) and alkaline phosphatase (ALP) were done on Cobas c 311 from Roche (Hitachi) biochemistry machine.

Statistical analysis

The recorded data was compiled and entered in a spreadsheet computer program (microsoft excel 2007) and then exported to data editor page of SPSS version 15 (SPSS Inc., Chicago, Illinois, USA).

Results

Table 1: Age wise distribution of study participants

Age in years	n	%
20-30	45	45
31-40	25	25
41-50	15	15
51-60	8	8
>61	7	7
Total	100	100

Most of the cases were seen in the 20-30 years age group followed by 31-40 years age group.

Table 2: Distribution of clinical features

Clinical Features	n	%
Fever	40	40
Myalgia	15	15
Fever and myalgia	12	12
Headache	8	8
Nausea and vomiting	7	7
Fever and skin rashes	8	8
Petechiae	6	6
Fever and itching	4	4

Majority of the patients were males compared to females and the male to female ratio was 2:1. Fever was the most common presentation and was seen in 40 cases (40%) cases.

Table 3: Distribution of study population by hemoglobin and hematocrit level

Hemoglobin			Hematocrit		
Hb (gm/dl)	No. of cases	%	HCT (%)	No. of cases	%
6-8.9	10	10	20-26	-	-
9-11.9	40	40	27-36	45	45
12-14.9	35	35	37-46	35	35
15-17.9	15	15	47-56	20	20

Present study showed hemoglobin range of 6 gm% to 17 gm% (Table 3). Raised hematocrit (>47%) was noted in 15 (15%) of patients at presentation and the hematocrit ranged from 20-51%. The total leukocyte count ranged from 1500 cells/cumm to >11000 cells/cumm. Leucopenia with less than 4000 cells/cumm was present in 25% cases. In the present study out of 100 cases of dengue fever, 90% cases had thrombocytopenia and 10% cases had severe thrombocytopenia (< 20,000/cumm) with bleeding manifestations.

Discussion

Dengue fever is an infectious disease which is difficult to distinguish from other viruses prevalent in our region as there are no specific markers that can diagnose the disease early. Because it is a disease that can evolve with serious consequences and even be fatal, this study aimed at analyzing clinical and epidemiological data and laboratory dynamics in order to try to identify biomarkers that are predictive of severity. Dengue is caused by a virus belonging to the Flaviviridae family (single-stranded, positive, and non-segmented RNA virus). Infection with one serotype confers immunity to only that serotype, and hence, a person may be infected up to 4 times. Humans are the main reservoir of dengue virus. [2]

The most common age group affected in our study was 21–30 years and males outnumbered females which is comparable to that of studies done by Neeraja et al. [14] and Kumar et al., [15] and the reason for male preponderance is said to be due to

their clothing habits or outdoor activities. Dengue is hemorrhagic viral fever which can prove fatal therefore this study is aimed at analyzing hematological and biochemical parameters for early diagnosis of dengue fever. Thrombocytopenia, leucopenia, increased hematocrit, lymphocytosis with reactive/atypical/plasmacytoid lymphocytes along with altered liver function tests are the hematological and biochemical abnormalities that appear in dengue fever.

In our study majority of the patients were males compared to females and the male to female ratio was 2:1. Deshwal et al and Vibha et al too observed a male predominance in their studies with 72.8% and 70% male patients respectively. [16,17] In this study, we found that fever was the commonest symptom in dengue patients, followed by headache/retro-orbital pain and myalgia. Other prominent symptoms were arthralgia, skin rash, skin hemorrhage, loose motion, mucosal bleed and nausea/vomiting. These findings were comparable to those documented by others, though the frequencies of the symptoms varied slightly. In our study, itching, especially in the palms and soles, was noted in 01% of patients which was comparable to similar study by Deshwal et al. [16]

Present study showed hemoglobin (Hb) ranging from 6 gm% to >15 gm%, 40 cases showed Hb of 9-11.9 gm%, followed by 35 cases showed Hb of 12-14.9 gm%, 10 had Hb of 6-8.9 gm% and 15 had Hb of 15-17.9 gm%. In the study by Meena et al hemoglobin ranged from 7.5-17.5 g/dl; mean hemoglobin value was 12.6 g/dl. [18] Hemoglobin level more than 15gm%

was seen in 6% cases. Dongre et al observed hemoglobin level from 3.6 gm/dl to 16.7gm/dl with a mean of 11.9 gm/dl. [19] Studies done by Gajera et al and Butt et al showed hematocrit values raised in 28% and 50% cases respectively. In studies done by Gajera et al and Butt et al there are more cases of dengue hemorrhagic fever 30% and 100% respectively. [20,21]

In the present study, total leukocyte count ranged from 1500 to >11000 cells/mm³. In Meena et al study total leukocyte count ranged from 1310 to 16700 cell/mm³, with mean total leukocyte count of 4701 cells/cumm. [22] In studies done by Yaseen et al and Gajera et al TLC <4,000/mm³ was seen in 50% and 39% cases respectively and >11,000/mm³ was seen in 12% cases which was similar to our study. [20,21] Leucopenia has been reported among dengue patients in many studies leucopenia is the most prominent hematological change sometimes with counts of less than 2,000/mm³. However mild leucocytosis with neutrophilia is seen at the onset of the disease developing leucopenia later on. Lymphocytosis was the common finding with the presence of atypical and plasmacytoid lymphocytes on peripheral smear were representative of augmented immune response to control the spread of dengue virus infected cells. [23]

Conclusion

Hemoconcentration, leucopenia, thrombocytopenia, and raised liver enzymes SGOT and SGPT along with reactive/ plasmacytoid lymphocytes on peripheral smear gives enough clues to test for dengue serology so that dengue cases can be diagnosed in their initial stages. This facilitates early treatment and aggressive fluid replacement therapy with good nursing care so that fatality rates can be reduced. The disease was more severe in individuals aged 15 years and older with a more pronounced and persistent presence of liver abnormalities (AST, ALT) and hemoconcentration. The study results are

relevant in the characterization of biological markers in the evolution of the disease and can be used as markers for the most severe forms thereby enabling early help with the adaption of therapeutic conduct for specific patients.

References

1. Dorland WA, editor. Dorland's Illustrated Medical Dictionary. 30th ed. Philadelphia, PA: Elsevier Saunders; 2003.
2. Banerjee M, Chatterjee T, Choudhary GS, Srinivas V, Kataria VK. Dengue: a clinicohaematological profile. Medical Journal Armed Forces India. 2008 Oct 1;64(4):333-6.
3. Nathan MB, Dreger RD, Guzman M. Epidemiology, burden of disease and transmission. In: Dengue Guidelines for Diagnosis, Treatment, Prevention and Control. New ed. Geneva, PA: WHO Press; 2009.
4. Chaudhuri M. What can India do about dengue fever? BMJ. 2013 Feb 4;346.
5. Simmons CP, Farrar JJ. Current concepts. N Engl J Med. 2012; 366 :14 23-32.
6. World Health Organization. dengue guidelines for diagnosis, treatment, prevention and control. 2009.
7. Amin P, Acicbe Ö, Hidalgo J, Jiménez JI, Baker T, Richards GA. Dengue fever: report from the task force on tropical diseases by the world federation of societies of intensive and critical care medicine. Journal of Critical care. 2018 Feb 1; 43:346-51.
8. Guzman MG, Kouri G. Dengue: an update. Lancet Infect Dis 2002; 2(1): 33-42.
9. Kouri GÁ, Guzman MG, Bravo JR. Why dengue haemorrhagic fever in Cuba? An integral analysis. Transactions of the Royal Society of Tropical Medicine and Hygiene. 1987 Sep 1;81(5):821-3.
10. Basheer A, Iqbal N, Mookkappan S, Anitha P, Nair S, Kanungo R, Kandasamy R. Clinical and laboratory

- characteristics of dengue-orientia tsutsugamushi co-infection from a tertiary care center in south india. *Mediterranean journal of hematology and infectious diseases*. 2016;8(1).
11. Gupta N, Srivastava S, Jain A, Chaturvedi UC. Dengue in india. *The Indian journal of medical research*. 2012 Sep;136(3):373.
 12. Kao CL, King CC, Chao DY, Wu HL, Chang GJ. Laboratory diagnosis of dengue virus infection: current and future perspectives in clinical diagnosis and public health. *J Microbiol Immunol Infect*. 2005; 38 (1):5-16.
 13. Ageep AK, Malik AA, Elkarsani MS. Clinical presentations and laboratory findings in suspected cases of dengue virus. *Saudi medical journal*. 2006 Nov 1;27(11):1711-3.
 14. Neeraja M, Lakshmi V, Teja VD, Umabala P, Subbalakshmi MV. Serodiagnosis of dengue virus infection in patients presenting to a tertiary care hospital. *Indian J Med Microbiol*. 2006; 24:280-2.
 15. Kumar A, Rao CR, Pandit V, Shetty S, Bammigatti C, Samarasinghe CM. Clinical manifestations and trend of dengue cases admitted in a tertiary care hospital, Udupi district, Karnataka. *Indian journal of community medicine: official publication of Indian Association of Preventive & Social Medicine*. 2010 Jul;35(3):386.
 16. Deshwal R, Qureshi MI, Singh R. Clinical and laboratory profile of dengue fever. *J Assoc Physicians India*. 2015 Dec 1;63(12):30-2.
 17. Gajera VV, Sahu S, Dhar R. Study of Haematological Profile of Dengue Fever and its Clinical Implication. *Annals Applied Bio-Sci*. 2016;3 (3): 2455-396.
 18. Ahmed F, Hussain Z, Ali Z. Clinical and hematological profile of patients with dengue fever. *J Med Sci*. 2014;22(1):17-20.
 19. Dongre T, Karmarkar P. Hematological Parameters and Its Utility in Dengue: A Prospective Study. *IOSR-J Dental Med Sci*. 2015; 14(2):31-4.
 20. Gajera VV, Sahu S, Dhar R. Study of haematological profile of dengue fever and its clinical implication. *Annals of Applied Biosciences*. 2016 Aug 10;3 (3): A241-246.
 21. Butt N, Abbassi A, Munir SM, Ahmad SM, Sheikh QH. Haematological and biochemical indicators for the early diagnosis of dengue viral infection. *J Coll Physicians Surg Pak*. 2008 May 1; 18(5):282-5.
 22. Meena KC, Jelia S, Meena S, Arif M, Ajmera D, Jatav VS. A study of hematological profile in dengue fever at a tertiary care center, Kota Rajasthan. *Int J Adv Med*. 2016; 3(3): 621-4.
 23. Diane S., Baldé A. K., Inapogui, C. B., Kéita A., & Diawara M. Functional results of cataract surgery using the phacoemulsification technique & quot; The CADES/O experience. *Journal of Medical Research and Health Sciences*. 2022; 5(9): 2256–2263.