

An Observational Assessment of Physical Activity and Dietary Habits among Adolescents

Binish Inam

Tutor, Department of Community Medicine, Anugrah Narayan Magadh Medical College and Hospital, Gaya, Bihar, India

Received: 16-09-2022 / Revised: 22-11-2022 / Accepted: 05-12-2022

Corresponding author: Dr. Binish Inam

Conflict of interest: Nil

Abstract

Aim: The current study aimed to focus specifically on objectively assessed physical activity & and dietary habits among adolescents of Bihar.

Methods: The present study was conducted at Anugrah Narayan Magadh medical College and Hospital, Gaya, Bihar, India and cross-sectional data were collected. Data were obtained from adolescents (12–16 years) recruited from three secondary schools in Bihar. Staff at participating schools selected a subset of their classes for participation. All students from nominated classes (n = 242) were eligible and received written information on the project. Consent was sought from parents prior to the study and adolescent participants provided assent before completing written surveys during class.

Results: In total, 221 pupils provided consent and completed the questionnaire (90% response rate). Of these, 123 (54%) provided usable accelerometer data. Comparison of these 123 adolescents with those who did not provide usable accelerometer data (n = 102) showed no significant differences in age, SES, ethnicity or fruit and vegetable consumption. A significantly ($p < 0.01$) higher proportion of those with usable accelerometer data were girls, compared to boys (53% compared to 37%), and ate breakfast on more days per week (4.7 days compared to 3.5 days). The final sample composition was 123 adolescents, with 61 boys, 62 girls, 68 younger adolescents and 55 older adolescents. The mean age of younger adolescents was 14.3 years and of older adolescents was 14.6 years.

Conclusion: Many adolescents fail to meet multiple diet and physical activity recommendations, highlighting that physical activity and dietary behaviours do not occur in isolation. Future research should investigate how best to achieve multiple health behaviour change in adolescent boys and girls.

Keywords: Adolescence, Life Style, Non-Communicable Disease, Physical Activity.

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

Introduction

In young people, physical activity and healthy diets including regular breakfast consumption and adequate levels of fruit and vegetables, have important short- and long-term health protective effects. [1] Physical inactivity is a major risk factor for mortality [2] and is associated with

cardiovascular diseases (CVD), diabetes, certain types of cancer, and high cholesterol levels. Poor diet and physical inactivity is also associated with premature mortality. High blood pressure, high cholesterol levels, lack of fruit and vegetables in one's diet, obesity, lack of

physical activity, and smoking are all risk factors for Non-Communicable Diseases (WHO). [3] As many of these risk factors are related to nutrition and activity levels, maintaining good habits in these areas are essential to maintain individual health and prevent disease. [4] Conversely, physical activity is essential for reducing the risk of CVD, type-2 diabetes, stroke, obesity, depression, dementia, and benign prostatic hyperplasia.

A data from the Health Behaviour in School-aged Children (HBSC) study shows that less than two-fifths of young people eat fruit daily, and only about a third eat vegetables daily. [5] Also, recent data suggest that 30% of boys and 40% of girls do not meet the current physical activity guidelines [6]. Health risks associated with individual behaviours are broadly acknowledged, yet, arguably a key issue for public health is that there is potential for synergistic effects of multiple health behaviours on the risk of chronic conditions and health outcomes. [7,8]

The current study aimed to focus specifically on objectively assessed physical activity & and dietary habits among adolescents of BIHAR.

Materials & Methods

The present study was conducted at Anugrah Narayan Magadh medical College and Hospital, Gaya, Bihar, India and cross-sectional data were collected. Study procedures were approved by the Ethical Advisory Committee. Data were obtained from adolescents (12–16 years) recruited from three secondary schools in Bihar. Staff at participating schools selected a subset of their classes for participation. All students from nominated classes ($n = 242$) were eligible and received written information on the project. Consent was sought from parents prior to the study and adolescent participants provided assent before completing written surveys during class. Participants completed questionnaires

during Physical Education or Personal, Social and Health Education (PHSE) lessons, under the supervision of trained researchers and class teachers.

Measures

Demographic information, including date of birth and ethnicity, was provided by the school. Questionnaires collected information on demographic characteristics of adolescents including gender and home postcodes. Socioeconomic status (SES) was determined using the Index of Multiple Deprivation (IMD), a measure of compound social and material deprivation, calculated from a variety of data including income, employment, health, education, and housing. It is based on the postcode of the participant's home, and thus represents an area level approximation of SES.

Adolescent dietary behaviour

Adolescent food intake was assessed using a 30-item food frequency questionnaire (FFQ), based on the previously validated Youth/Adolescent Food Frequency questionnaire (YAQ). [9] Adolescents were asked how often they ate ten different fruits and twelve different vegetables in the past month. Responses to questions on the frequency of consumption of specific fruits and vegetables were summed to compute total frequency of fruit and vegetable consumption/day, respectively. For the purpose of this study each item was summed to calculate the frequency of consumption of 'fruits and vegetables' (FV) per day. Guided by the current recommendations for fruit and vegetable consumption (5 portions of FV/day), the total daily frequency of FV consumption was dichotomised into <5 times/day or ≥ 5 times/day.

Breakfast consumption was assessed with a single-item asking adolescents how often they ate breakfast in the past seven days. While there are no current national recommendations for frequency of

breakfast consumption, evidence suggests that young people who eat breakfast on most days of the week see health benefits (e.g. lower Body Mass Index) compared to those who skip breakfast.[10,11] A recent government initiative in the bihar recommends that parents encourage their children to eat breakfast regularly. Given such evidence, frequency of breakfast consumption/week was dichotomized into <5 days/week or ≥ 5 days/week.

Physical activity

Physical activity was assessed by Actigraph GT1M accelerometers (ActiGraph, Fort Walton Beach, FL) using a 5 second measurement interval (epoch). Participants were instructed to wear the accelerometer over their right hip for one-week. Exceptions included time spent sleeping, showering and during water-based activities. Duration of MVPA was computed only for adolescents who wore the accelerometer for a minimum of three days[12], defined as days on which accelerometer counts were between 10,000 and 20 million.[13] To estimate the time spent per day in moderate intensity physical activity (3.0–5.9 metabolic equivalent of rest [METs]) and vigorous intensity physical activity (6.0+ METs), age specific movement count thresholds were applied. Time-per-day in MVPA was derived by summing these values across valid days. The proportion of adolescents meeting the physical activity recommendations for young people was calculated according to whether they performed an average of ≥ 60 mins/day of MVPA. Time spent in MVPA was dichotomised as <60 mins/day or ≥ 60 mins/day

Statistical analyses

All analyses were conducted using SPSS statistical software version 16.0. Descriptive statistics were used to summarise sociodemographic, physical activity and dietary characteristics of the sample. Adolescents were categorised as

younger or older adolescents by dichotomising at the mean (14.4 years). Mann-Whitney tests were performed to examine gender and age-group differences in mean minutes/day spent in MVPA, fruit and vegetable consumption per day, and breakfast consumption per week. The proportion of adolescents achieving ≥ 60 mins/day of MVPA, and the proportion of adolescents consuming fruit and vegetables ≥ 5 times/day, and breakfast ≥ 5 times/ week were compared by gender and age-group using Pearson's chi-square (χ^2) tests of significance. A total risk behaviour score was calculated for each participant based on the total number of unmet health recommendations (range from zero to three). Pearson's chisquare tests were used to examine gender and age-group differences in the number of risk behaviours. The proportion of adolescents in each multiple-risk behaviour combination was determined to examine behaviour clustering patterns, and Pearson's chi-square tests were used to examine gender and age-group differences in the clustering patterns.

Results

Sample characteristics

In total, 221 pupils provided consent and completed the questionnaire (90% response rate). Of these, 123 (54%) provided usable accelerometer data. Comparison of these 123 adolescents with those who did not provide usable accelerometer data ($n = 130$) showed no significant differences in age, SES, ethnicity or fruit and vegetable consumption. However, a significantly ($p < 0.01$) higher proportion of those with usable accelerometer data were girls, compared to boys (53% compared to 37%), and ate breakfast on more days per week (4.7 days compared to 3.5 days). Thus, our figures likely underestimate the prevalence of not meeting the breakfast recommendations. The final sample composition was 123 adolescents, with 61 boys, 62 girls, 68 younger adolescents and

55 older adolescents. The mean age of older adolescents was 14.6 years and of younger adolescents was 14.3 years and of

Table 1: Distribution fruit and vegetables consumption per day, breakfast consumption per week and minutes per day spent in moderate-to-vigorous intensity physical activity (MVPA)

	Total (n = 123)	Boys (n = 61)	Girls (n = 62)	Younger adolescents (n = 68)	Older adolescents (n = 55)
MVPA minutes/day					
25th percentile	19.5		21.5	39.2	22.6
50th percentile	32.5	38.3	38.5	48.1	38.1
75th percentile	68.0	54.3	49.1	65.1	49.5
Frequency of FV consumption/day					
25th percentile	2.9	1.9	3.7	2.5	3.8
50th percentile	4.2	3.8	4.3	3.9	4.9
75th percentile	7.2	6.8	5.8	6.1	8.1
Frequency of breakfast consumption, days/week					
25th percentile	5	4	2.5**	5	5
50th percentile	6	7	7	7	6
75th percentile	7	8	8	8	8

Prevalence of meeting health recommendations

The proportion of boys, girls, younger and older adolescents meeting the physical activity, fruit and vegetable, and breakfast consumption recommendations are displayed in Table 2. A higher proportion of boys, compared to girls, ate breakfast on more than 5 days/week ($p < 0.01$). A higher proportion of younger adolescents, compared to older adolescents, met the physical activity recommendations and did not meet the recommendations for fruit and vegetable consumption ($p < 0.01$). Only 6% of the adolescent sample met the recommendations for all three of the health behaviours and almost 10% had three health risk behaviours. A higher proportion of girls, compared to boys, had three health risk behaviours ($p < 0.01$).

The clustering patterns of the three health risk behaviours are described in Table 3. For adolescents with two risk behaviours, the most prevalent cluster was formed by not meeting the physical activity and fruit and vegetables recommendations. A higher proportion of girls, compared to boys, had the cluster pattern of not meeting the recommendations for physical activity and breakfast consumption. ($p < 0.01$). A higher proportion of boys, compared to girls, and older adolescents, compared to younger adolescents, had the most prevalent single risk factor of not meeting the recommendations for physical activity ($p < 0.01$). A higher proportion of younger adolescents, compared to older adolescents, had the single risk factor of not meeting the recommendations for fruit and vegetable consumption ($p < 0.01$).

Table 2: Gender and age distribution of meeting health recommendations

	Total (n = 123)	Boys (n = 61)	Girls (n = 62)	Younger adolescents (n = 68)	Older adolescents (n = 55)
Meet >60 minutes MVPA per day, n (%)					
No	97 (78.8)	44 (72.4)	52 (83.1)	42 (69.4)	49 (88.5)
Yes	26 (22.2)	17 (27.6)	10 (16.8)	26 (30.6)	6 (11.5)
Meet >5 portions fruits/ vegetables, n (%)					
No	68 (55.2)	31 (51.2)	37 (59.6)	44(64.3)	25 (44.9)**
Yes	55 (44.8)	30 (48.3)	35 (40.4)	24 (35.7)	30 (55.1)
Meet >5 days a week eating breakfast, n (%)					
No	29 (23.3)	9 (14.9)	19 (31.5)	16 (23.5)	13 (23.1)
Yes	94 (76.7)	52 (85.1)	43 (68.5)**	52 (76.5)	42 (76.9)
Number of risk behaviours, n (%)					
0	8(6.2)	7 (10.3)	1 (2.2)	5 (7.1)	3 (5.1)
1	50 (40.3)	28 (46.0)	22 (34.8)	25 (36.7)	25 (44.9)
2	55(43.8)	23 (37.9)	30 (49.4)	33 (48.0)	21 (38.5)
3	10 (9.7)	5 (5.7)	9 (13.5)**	5 (8.2)	6 (11.5)

Table 3: Descriptive cluster pattern of multiple risk behaviours

Number of risk behaviours	Percent of sample	Boys (n = 61)	Girls (n = 62)	Younger adolescents (n = 68)	Older adolescents (n = 55)
3: All three risk behaviours	9.7	5.7	13.5**	8.2	11.5
2: MVPA < 60 minutes/day and < 5 fruit/ vegetables per day	33.0	31.0	34.8	36.7	28.2
2: MVPA < 60 minutes/day and < 5 days a week eating breakfast	6.2	1.1	11.2**	5.1	7.7
2: < 5 fruit/vegetables per day and < 5 days a week eating breakfast	5.7	5.7	5.6	8.2	2.6
1: MVPA < 60 minutes/day	29.0	34.5	23.6**	19.4	41.0**
1: < 5 fruit/vegetables per day	7.4	9.2	5.6	11.2	2.6**
1: < 5 days a week eating breakfast	2.8	2.3	3.4	4.1	1.3
0: No risk behaviours	6.2	10.3	2.2**	7.1	5.1

Discussion

This study described the prevalence and clustering patterns of three health behaviours (physical activity, fruit and vegetable consumption, breakfast consumption) in a sample of adolescents from the Bihar. Almost 54% of adolescents had multiple diet and physical activity risk behaviours, and only 6% achieved the recommendations for all three of the health behaviours. Poor diet and physical inactivity are established risk factors for chronic disease. In young

people, physical activity and healthy diets including regular breakfast consumption and adequate levels of fruit and vegetables, have important short- and long-term health protective effects. For example, physical activity in young people may benefit cardiovascular disease (CVD) risk factors, adiposity and bone health, which could influence health in adulthood. [14]

Higher levels of MVPA were found among boys, compared to girls, and in younger adolescents, compared to older adolescents. Such findings support

previous research [15,16] highlighting gender differences and age related declines in physical activity, using objective measures of physical activity among adolescents. Recent research using accelerometry has also shown that age and gender differences are evident when comparing children as young as six and eleven years of age. [17,18] This suggests that the primary school years may be critical for the development of disparities in physical activity behaviours. [19] Efforts to promote physical activity should begin in these critical years given that active children are more likely to become active adults. [20] Consistent with previous research, adolescent girls ate breakfast on fewer days per week compared to boys. [21] Despite the evidence that adolescents who skip breakfast are more likely to be overweight than those who regularly eat breakfast [23], skipping breakfast may be a chosen method of weight control for girls, and in some individuals may be associated with dieting, or disordered eating. [22]

Older adolescents ate more fruit and vegetables per day, and were more likely to meet the recommendations for fruit and vegetable consumption, compared to younger adolescents. In contrast, review level evidence has shown a negative association between age and fruit and vegetable consumption. [24] Contrasting findings may reflect a difference in the methodologies employed to assess fruit and vegetable consumption. Several studies have identified overestimation of fruit and vegetable intake when using food frequency questionnaires. [25,26] Girls were more likely to have the two behaviour risk cluster that paired physical activity and breakfast consumption. A possible explanation for such clustering is that adolescents who frequently skip breakfast have lower daily energy intakes, with higher daily energy intake being associated with more time spent being physically active. [27] Girls who skip

breakfast as part of a diet or method of weight control, may have less energy to be physically active. Efforts to promote both physical activity and regular breakfast consumption to adolescent girls, in particular, are needed. Consistent with previous research, girls had a higher number of risk factors related to physical activity and dietary behaviours compared to boys. [28] Such findings provide additional evidence for support of gender specific interventions promoting physical activity and dietary behaviours. Although health promotion programmes frequently target multiple behaviours, little is known about the best approaches to stimulating multiple behaviour change in adolescents. [28] Reviews of multiple behaviour interventions with young people have revealed changes in some but not all behaviours, with significant effects more likely for dietary as opposed to physical activity outcomes. [29] There is little evidence of covariation among diet and physical activity behaviours in adolescents over time. [30,31]

Conclusion

Many adolescents fail to meet multiple diet and physical activity recommendations, supporting previous evidence that physical activity and dietary behaviours do not occur in isolation. Differences in dietary and physical activity behaviours between adolescent boys and girls, as well as between older and younger adolescents, should be taken into consideration when assessing the efficacy of strategies promoting multiple health behaviour change.

References

1. Biddle SJ, Gorely T, Stensel DJ: Health-enhancing physical activity and sedentary behaviour in children and adolescents. *J Sports Sci.* 2004; 22:679-701.
2. Katzmarzyk PT, Janssen I, Ardern CI. Physical inactivity, excess adiposity

- and premature mortality. *Obesity Rev.* 2003; 4:257-90.
3. Waxman A. WHO global strategy on diet, physical activity and health. *Food Nutr Bull.* 2004; 25:292-302.
 4. Williams PT. Physical fitness and activity as separate heart disease risk factors: A meta-analysis. *Med Sci Sports Exerc.* 2001; 33:754-61.
 5. World Health Organization: Young people's health in context: selected key findings from the Health Behaviour in School- aged Children study.
 6. The Information Centre: Statistics on obesity, physical activity and diet: England, February. 2009.
 7. McGinnis JM, Foege WH: Actual causes of death in the United States. *J Am Med Assoc.* 1993; 270:2207-2212.
 8. McKenna M, Taylor WC, Marks J, Koplan J: Current issues and challenges in chronic disease control. In *Chronic disease epidemiology and control* Edited by: Brownson R, Remington P, Davis J. Washington DC: American Public Health Association, United Book Press; 1998.
 9. Rockett HR, Breitenbach M, Frazier AL, Witschi J, Wolf AM, Field AE, Colditz GA: *Prev Med.* 1997, 26:808-816.
 10. Barton BA, Eldridge AL, Thompson D, Affenito SG, Striegel-Moore RH, Franko DL, Albertson AM, Crockett SJ: breakfast and cereal consumption to nutrient intake and *J Am Diet Assoc.* 2005; 105:1383-1389.
 11. Delva J, Johnston LD, O'Malley PM: weight and Related Lifestyle Behaviors. *Racial/Ethnic and Am J Prev Med.* 2007; 33: s178-s186.
 12. Mattocks C, Ness A, Leary S, Tilling K, Blair SN, Shield J, Deere K, Saunders J, Kirkby J, Smith GD, et al.: *J Phys Act Health.* 2008; 5: S98-S111.
 13. Telford A, Salmon J, Jolley D, Crawford D: Reliability and validity of a self-report and proxy-report physical activity questionnaire: the Children's Leisure Activities Study Survey. *Pediatr Exerc Sci.* 2004; 16:64-78.
 14. Biddle SJ, Gorely T, Stensel DJ. Health-enhancing physical activity and sedentary behaviour in children and adolescents. *Journal of sports sciences.* 2004 Aug 1;22(8):679-701.
 15. Trost SG, Pate RR, Sallis JF, Freedson PS, Taylor WC, Dowda M, Sirard J. Age and gender differences in objectively measured physical activity in youth. *Medicine & Science in Sports & Exercise.* 2002 Feb 1;34(2):350-5.
 16. Kimm SY, Glynn NW, Kriska AM, Fitzgerald SL, Aaron DJ, Similo SL, McMahon RP, Barton BA. Longitudinal changes in physical activity in a biracial cohort during adolescence. *Medicine & Science in Sports & Exercise.* 2000 Aug 1;32(8):1445-54.
 17. Hesketh K, Crawford D, Salmon J. Children's television viewing and objectively measured physical activity: associations with family circumstance. *International Journal of Behavioral Nutrition and Physical Activity.* 2006 Dec;3(1):1-0.
 18. Bagley S, Salmon JO, Crawford D. Family structure and children's television viewing and physical activity. *Medicine and science in sports and exercise.* 2006 May 1;38(5):910-8.
 19. Brodersen NH, Steptoe A, Boniface DR, Wardle J. Trends in physical activity and sedentary behaviour in adolescence: ethnic and socioeconomic differences. *British journal of sports medicine.* 2007 Mar 1;41(3):140-4.
 20. Telama R, Yang X, Viikari J, Välimäki I, Wanne O, Raitakari O. Physical activity from childhood to adulthood: a 21-year tracking study. *American journal of preventive medicine.* 2005 Apr 1;28(3):267-73.
 21. Pearson N, MacFarlane A, Crawford D, Biddle SJ. Family circumstance and adolescent dietary behaviours. *Appetite.* 2009 Jun 1;52(3):668-74.

22. Rashidi A, Mohammadpour-Ahranjani B, Karandish M, Vafa MR, Hajifaraji M, Ansari F, Sadeghi S, Maddah M, Kalantari N, Akhavi-Rad MB. Obese and female adolescents skip breakfast more than their non-obese and male peers. *Open Medicine*. 2007 Dec 1;2(4):481-7.
23. Albertson AM, Franko DL, Thompson D, Eldridge AL, Holschuh N, Affenito SG, Bauserman R, Striegel-Moore RH. Longitudinal patterns of breakfast eating in black and white adolescent girls. *Obesity*. 2007 Sep;15(9):2282-92.
24. Rasmussen M, Krølner R, Klepp KI, Lytle L, Brug J, Bere E, Due P. Determinants of fruit and vegetable consumption among children and adolescents: a review of the literature. Part I: quantitative studies. *International journal of behavioral nutrition and physical activity*. 2006 Dec;3(1):1-9.
25. Kremers SP, Brug J, De Vries H, Engels RC. Parenting style and adolescent fruit consumption. *Appetite*. 2003 Aug 1;41(1):43-50.
26. Van Assema P, Brug J, Ronda G, Steenhuis I, Oenema A. A short dutch questionnaire to measure fruit and vegetable intake: relative validity among adults and adolescents. *Nutrition and Health*. 2002 Apr;16(2):85-106.
27. Berkey CS, Rockett HR, Gillman MW, Field AE, Colditz GA. Longitudinal study of skipping breakfast and weight change in adolescents. *International journal of obesity*. 2003 Oct;27(10):1258-66.
28. Sanchez A, Norman GJ, Sallis JF, Calfas KJ, Cella J, Patrick K. Patterns and correlates of physical activity and nutrition behaviors in adolescents. *American Journal of Preventive Medicine*. 2007 Feb 1;32(2):124-30.
29. Resnicow K, Robinson TN. School-based cardiovascular disease prevention studies: review and synthesis. *Annals of Epidemiology*. 1997 Oct 1;7(7):S14-31.
30. Rosenberg DE, Norman GJ, Sallis JF, Calfas KJ, Patrick K. Covariation of adolescent physical activity and dietary behaviors over 12 months. *Journal of Adolescent Health*. 2007 Nov 1;41(5):472-8.
31. Manfred, D. May There Exist Healthy Diseases? *Journal of Medical Research and Health Sciences*, 2022; 5(3):1801–1803.