

Assessment of the Impact of Corrected Maternal Anemia on Perinatal Outcome in Terms of Birth Weight and Gestational Maturity and its Comparison with Babies Born to Anemic Mothers

Bheemsen Kumar¹, Sanju Kumari², Gopal Shanker Sahni³

¹ Senior Resident, Department of Pediatrics, SKMCH, Muzaffarpur, Bihar, India

² Junior Resident, Department of Obstetrics and Gynecology, Nalanda Medical College and Hospital, Patna, Bihar India

³ HOD & Associate Professor, Department of Pediatrics, SKMCH, Muzaffarpur, Bihar, India

Received: 25-12-2022 / Revised: 05-01-2023 / Accepted: 28-01-2023

Corresponding author: Dr. Sanju Kumari

Conflict of interest: Nil

Abstract

Aim: The aim of the present study was to assess impact of corrected maternal anemia on perinatal outcome in terms of birth weight and gestational maturity and its comparison with babies born to anemic mothers and burden and severity of anemia in pregnant women.

Methods: The present study was conducted at Department of Pediatrics SKMCH Muzaffarpur, Bihar, India for one year. Total 120 present women of 20+4 weeks of gestation with singleton pregnancy attending OPD of department of obstetrics & gynecology at NMCH for one year and were enrolled after taking informed consent.

Results: A total of 120 women with pregnancy around 20+4 week's gestation were included in this study. As per hemoglobin levels and WHO definition of anemia, 6 (5%) antenatal mothers had severe anemia, 78 (65%) had moderate anemia and 36 (30%) had Hb levels >10gm/dl. 78 women with moderate anemia i.e. hemoglobin level between 7-10gm% constituted our study group. Out of 78 women 4 were lost to follow up, one who developed PIH, one with bleeding and 12 who did not show improvement in hemoglobin level due to irregular intake or intolerance were excluded from this study.

Conclusion: From the present study we concluded that; anemic mothers had higher chance of delivering preterm and low birth weight baby, which again is an important determinant of survival and quality of life a child.

Keywords: Maternal anemia, Perinatal outcome, Premature and low birth weight babies.

This is an Open Access article that uses a fund-ing model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

Introduction

Low birth weight has been widely studied and is an important risk factor for infant morbidity and mortality. [1-4] However, insufficient weight has received little attention [5-8], even though three decades ago, children with birth weights less than

3000 grams were considered to have a risk of mortality that was three times higher during the first year of life than that of children whose weights were above or equal to this cut-off point. [9,10]

The classic risk factors for low birth weight are associated with unfavorable biological, social and environmental conditions that may occur before or during the pregnancy period. [4,11-13] Nutritional determinants, such as pre-gestational weight and weight gain during pregnancy, influence birth weight. Thus, inadequate maternal caloric intake, which may be the result of a diet that is nutritionally poor, leads to lower absorption of essential micronutrients, such as vitamin B12 and iron, for fetal growth. [14]

Multiple factors lead to anaemia in pregnancy, nutritional iron deficiency anaemia (IDA) being the commonest. Physiological haemodilution and several factors affecting Hb and iron status in pregnancy lead to difficulties in establishing a definitive diagnosis. Fetal programming occurs when the normal pattern of fetal development is disrupted by an abnormal stimulus or 'insult' applied at a critical point in in-utero development.¹⁵ Correction of anaemia with haematinics provides adequate proof of a nutritional deficiency in its pathogenesis.

The aim of the present study was to assess impact of corrected maternal anemia on perinatal outcome in terms of birth weight and gestational maturity and its comparison with babies born to anemic mothers and burden and severity of anemia in pregnant women.

Methods

The present study was conducted at Department of Pediatrics SKMCH Muzaffarpur, Bihar, India for one year. Total 120 present women of 20+4weeks of gestation with singleton pregnancy attending OPD of department of obstetrics & gynecology at NMCH for one year and were enrolled after taking informed consent.

Exclusion Criteria: Antenatal mothers having associated conditions that may affect the fetal growth were excluded.

1. Women with systemic disease like primary cardiac, renal, hepatic disorder and hypothyroidism.
2. History of pathologic blood loss at any stage during study period.
3. Antenatal mother who developed PIH even after enrollment.
4. Those that did not show improvement of anemia after supplementation.
5. Mothers who had anemia other than nutritional.

Laboratory Investigation: To know the severity of anemia complete haemogram was done, and as per WHO criteria, on the basis of hemoglobin level¹¹ they were divided into 3 groups

Group A (Hb < 7) severe anemia

Group B (Hb 7- 10) moderate anemia

Group C (Hb > 10) mild/ no anemia

Women of Group B with hemoglobin levels of 7- 10 gm% became our study group. These were supplemented with oral hematinics as per protocol of management of anemia during pregnancy and these were followed till delivery for assessment of perinatal outcome. Oral haematinic was given in form of tablets with following composition Ferrous Ascorbate equivalent to elemental iron 100 mg, Folic acid 1.1 mg. These tablets were given one tablet twice daily after meals as per standard protocol of management of anemia during pregnancy. These patients were regularly followed till the time of delivery and improvement of anemia was assessed by hemoglobin levels done at regular intervals and at the time of delivery. Perinatal outcome in terms of prematurity and low birth weight babies was assessed in babies of mother who showed improvement of their anemia during follow up. Gestational age was assessed by modified Ballard scoring system (Ballard et al, 1991). Birth weight of babies of mothers enrolled for

the study was taken by same digital weighing machine used in labor room. Low birth weight baby was defined as per standard definition of birth weight of less than 2500 gms.

Babies of this group were compared with babies of 60 mothers who had anemia at the time of delivery. Their records were checked retrospectively to assess their anemia status during pregnancy. Only

those mothers were included in comparison group, whose hemoglobin levels were between 7-10 gm% at the time of delivery and who had previous lab records suggestive of moderate anemia. Those mothers who were anemic but had any other risk factor which was likely to affect the growth of the baby were excluded

Results

Table 1: Distribution of antenatal mothers according to their hemoglobin level

Groups	Range of Hb (gm/dl)	Study group n = 120	Mean Hb. (gm/dl)	HCT
A	< 7	6 (5%)	6.4	19.2
B	7- 10	78 (65%)	8.6	25.8
C	>10	36 (30%)	11.6	34.8

A total of 120 women with pregnancy around 20+4 week's gestation were included in this study. As per hemoglobin levels and WHO definition of anemia, 6 (5%) antenatal mothers had severe anemia, 78 (65%) had moderate anemia and 36 (30%) had Hb levels >10gm/dl. 78 women with moderate anemia i.e. hemoglobin

level between 7-10gm% constituted our study group. Out of 78 women 4 were lost to follow up, one who developed PIH, one with bleeding and 12 who did not show improvement in hemoglobin level due to irregular intake or intolerance were excluded from this study.

Table 2: Anemia status of antenatal mothers at the time of delivery

Groups	Supplemented Group (60)	Comparison Group (60)
Mean Hb Level (gm/dl) Before supplementation	~ 8.4	
Mean Hb level (gm/dl) After supplementation	~ 11.4	~ 8.2
HCT	34.2	24.6

In remaining 60 women improvement in mean hemoglobin was reported to be from 8.4 gm% to 11.4 gm%. Perinatal outcome in terms of birth weight and gestational age was assessed in these 60 women and was compared with 60 babies delivered to anemic mothers.

Table 3: Distribution of babies of anemic and non-anemic mothers as per their birth weight

Groups	Antenatal mothers (n) Birth Weight (>2500 gms)	Normal weight babies Birth Weight <2500 gms	Low birth weight babies
Non Anemic mothers	60	55 (91.66%)	5 (8.34%)
Anemic mothers	60	42 (70%)	18 (30%)

Improvement of maternal anemia was found to have significant positive effect on birth weight of baby in our study. 5 (8.34%) babies had low birth weight in supplemental group as compared to 30% in anemic group.

Table 4: Mean birth weight in two groups

Groups	Mean Birth wt (gms)
Non anemic group	2746
Anemic group	1876

Mean birth weight was 2746 gms in non-anemic group (supplemental group) as compared to 1876 gms in anemic group. Babies of non-anemic group showed improvement in birth weight of about 870 gms with maternal supplementation.

Table 5: Distribution of babies according to their gestation in anemic and non-anemic group

Groups	Preterm	Term	Post term	Total
Non anemic group	3 (5%)	56	1	60
Anemic Group	6 (10%)	54	0	60

Effect of improvement of maternal anemia on maturity of baby is depicted in the above table. No significant effect of maternal anemia on gestational age could be demonstrated in our study.

Discussion

Anaemia is hematological abnormality and has a significant public health problem globally. World Health Organization (WHO) reports that more than 1.62 billion people are affected globally. Of these, 56 million pregnant women (41.8%) were suffering from varying degrees of anaemia. [16] Despite anaemia being found worldwide, it's more prevalent in developing countries. [17] In India, of all infants who died before they completed 29 days post-birth, 48.1% suffered from LBW and premature birth. [20,21] Maternal anemia is reported to be one of the important factors of low birth weight of babies. The effect of maternal anemia on birth weight have been reported by many studies, but the level of hemoglobin at which this association is seen is reported to be, different in various studies. Most of the studies reported that hemoglobin level of less than 10 gm% had significant effect on birth weight of baby [18,20-22] but Manisha et al [19] found positive association only with severe anemia with hemoglobin level < 7gm%. Similar inverse relation with maternal hemoglobin level between 6-8 gm% and low birth weight was

reported by Ganesh et al. [23] In contrast K. Jagdish et al in their study observed no association between maternal anemia and birth weight of babies. [24]

In our study percentage of low birth weight babies was 27.8% in anemic mothers which was high as compared to 8.9% in non-anemic mothers. The overall prevalence of LBW as reported by various studies from India ranged from 28% to 40%. [21,23,24] Risk of having low birth weight baby increases when mother is anemic. [25,26] It was 1.9 to 4.8 times more as reported by Lone FW and Deshmukh. et al in their study. [27,28]

Maternal anemia has long been a known risk factor for preterm delivery along with its effect on birth weight. In our study 6.4% babies were born prematurely to non-anemic mothers as compared to 11.3% babies of anemic mothers. Incidence of preterm delivery was reported to be as high as 45% in moderate anemia group as compared to 50% in severely anemic mothers. [29] In our study intrauterine growth retardation was more responsible for low birth weight than prematurity. On the contrary Sharma S reported prematurity to be more associated with low birth weight. [30]

This study also reveals no significant association with initial Hb level, initial maternal weight on 1st visit and weight gain through 4 visits. But, study by Hosen

et al conclude that maternal weight at 1st visit (<45 kg) and weight gain (<7 kg) during pregnancy have been found to be consistently associated with LBW. [31,32]

Conclusion

From the present study we concluded that; anemic mothers had higher chance of delivering preterm and low birth weight baby, which again is an important determinant of survival and quality of life a child. This reiterates importance of preventing and controlling maternal anemia not only for mothers but also for the health of a baby.

References

1. Kramer MS. Determinants of low birth weight: methodological assessment and meta-analysis. *Bull World Health Organ.* 1987; 65(5):663–737.
2. World Health Organization. Meeting of Advisory Group on Maternal Nutrition and Low Birthweight Geneva: WHO; 2002.
3. Moraes IB. Risk factors for underweight at birth. Campinas, São Paulo: Universidade Estadual de Campinas. 2001.
4. UNICEF. Low Birthweight: Country, regional and global estimates. New York: UNICEF; 2004.
5. Raisanen S, Kancherla V, Gissler M, Kramer MR, Heinonen S. Adverse perinatal outcomes associated with moderate or severe maternal anaemia based on parity in Finland during 2006–10. *Paediatr Perinat Epidemiol.* 2014;28(5):372–80.
6. Mesa SLR, Sosa BEPS, Go´mez JA, Lo´pez NZ, Di´az CAG, Moreno CMR, Alarca NAC, Va´squez LEE. Maternal nutritional status and its relationship with birth weight of the newborn, pregnant women study Public Network of Medellı´n, Colombia *Perspect Nutr Humana.* 2012; 14(201):199–208.
7. de Araujo Filho AC, Sales IM, de Araujo AK, Almeida PD, da Rocha SS. Epidemiological aspects of neonatal mortality in a capital from northeastern Brazil/Aspectos epidemiol´gicos da mortalidade neonatal em capital do nordeste do Brasil/Aspectos epidemiol´gicos de la mortalidad neonatal en la capital del nordeste de Brasil. *Revista Cuidarte.* 2017 Sep 1;8 (3):1767-77.
8. Costa RSC, Caldevilla DE, Gallo PR, Sena BF, Leone C. Incidence and characteristics of insufficient birth weight Newborns from a cohort of neonates in a public regional Hospital of a metropolitan area. *Journal of Human Growth and Development.* 2013; 23(2):238–44.
9. Puffer RR, Serrano CV. Patterns of birthweights: a summary. *Bulletin of the Pan American Health Organization (PAHO);* 21 (2), 1987. 1987.
10. Rees JM, Lederman SA, Kiely JL. Birth weight associated with lowest neonatal mortality: infants of adolescent and adult mothers. *Pediatrics.* 1996 Dec;98(6):1161-6.
11. Carniel ED, Zanolli MD, Antˆnio Mˆ, Morcillo AM. Determinants for low birth weight according to Live Born Certificates. *Revista Brasileira de Epidemiologia.* 2008; 11:169-79.
12. Melo AS, Assuno PL, Gondim SS, Carvalho DF, Amorim MM, Benicio MH, Cardoso MA. Estado nutricional materno, ganho de peso gestacional e peso ao nascer. *Revista Brasileira de Epidemiologia.* 2007 Jun;10(2):249-57.
13. LTC G. Fr´u CM, Brandˆo M, Nunes ML. Determinantes sociais e biol´gicos da mortalidade infantil em coorte de base populacional em Passo Fundo, Rio Grande do Sul. *Cienc Sa´de Colet [Internet].* 2010;15(2): 363-70.
14. Bresani CC, Souza BA, Batista Filho M, Figueiroa JN. Anemia and iron deficiency in pregnant women: disagreements among the results of a cross-sectional study. *Revista*

- Brasileira de Saúde Materno Infantil. 2007;7:s15-21.
15. H.K.Cheema, Bajwa B S, Kaur K, Joshi H. Prevalence and possible risk factor of anemia in different trimester of pregnancy. International Journal of Contemporary Medical Research 2016;3(4):1194-1197.
 16. McLean E, Cogswell M, Egli I, Wojdyla D, De Benoist B. Worldwide prevalence of anaemia, WHO vitamin and mineral nutrition information system, 1993–2005. Public health nutrition. 2009 Apr;12(4):444-54.
 17. Parks S, Hoffman MK, Goudar SS, Patel A, Saleem S, Ali SA, Goldenberg RL, Hibberd PL, Moore J, Wallace D, McClure EM. Maternal anaemia and maternal, fetal, and neonatal outcomes in a prospective cohort study in India and Pakistan. BJOG: An International Journal of Obstetrics & Gynaecology. 2019 May;126(6):737-43.
 18. Kalaivani K. Prevalence & consequences of anaemia in pregnancy. Indian Journal of Medical Research. 2009 Nov 1;130(5):627-33.
 19. Nair M, Choudhury MK, Choudhury SS, Kakoty SD, Sarma UC, Webster P, Knight M. Association between maternal anaemia and pregnancy outcomes: a cohort study in Assam, India. BMJ Global Health. 2016 Apr 1;1(1):e000026.
 20. Causes of death lbw govt. census report 2010- 2013 release.
 21. Press Information Bureau Government of India Ministry of Health and Family Welfare 06-May-2016 14:18 IST Maternal and Child Mortality Rate.
 22. Sekhvat. L, Davar R, Hosseinidezoki S. Relationship between maternal hemoglobin concentration and neonatal birth weight. Hematology.2011 Nov; 16(6):373-6.
 23. Ganesh K S, Harsha Kumar HN, Jayaram S and Kotian M S. Determinants of Low Birth Weight: A Case Control Study in a district hospital Karnataka Indian J Pediatr 2010; 77 (1) 87-89.
 24. K Jagadish Kumar, N Asha, D Srinivasa Murthy, MS Sujatha,2 and VG Manjunath. Maternal Anemia in Various Trimesters and its Effect on Newborn Weight and Maturity: An Observational Study: Int J Prev Med. 2013 Feb;4(2):193–199.
 25. Agarwal G, Ahmad S, Goel K, Kumar V, Goel P, Garg M, Punj A. Maternal risk factors associated with low-birth-weight neonates in a tertiary care hospital, Northern India. J Community Med Health Educ. 2012;2(9):1000177.
 26. Agarwal K, Agarwal A, Agrawal VK, Agrawal P, Chaudhary V. Prevalence and determinants of “low birth weight” among institutional deliveries. Annals of Nigerian medicine. 2011 Jul 1;5(2):48.
 27. Lone FW, Qureshi RN, Emmanuel F. Maternal anaemia and its impact on perinatal outcome in a tertiary care hospital in Pakistan. Et Mediterr Health J. 2004 Nov;10(6):801-7.
 28. Kumar V, Deshmukh PR, Taywade M, Gupta SS. Magnitude and correlates of low birth weight at term in rural Wardha, Central India. Online J Health Allied Scs. 2016 Apr 28;15(1):2.
 29. Bangal VB, Kunal A, Ketki B, Tuse H. Study of maternal and perinatal outcome in moderate to severe degree iron deficiency anaemia in rural community. Int J Biomed Adv Res. 2016;7(2):88-93.
 30. Sharma SR, Giri S, Timalisina U, Bhandari SS, Basyal B, Wagle K, Shrestha L. Low birth weight at term and its determinants in a tertiary hospital of Nepal: a case-control study. PloS one. 2015 Apr 8;10(4):e0123962.
 31. Hosain M, Chatterjee N, Begum A, Saha SC. Factors associated with low birthweight in Rural Bangladesh. J Trop Pediatr. 2006;52(2):87-91.
 32. Martínez J. A. P., Saavedra A. J. G., Bohorquez G. D. B., Cordero J. F. B., Barros B. D., Ríos M. K. M., Duque L.

E. D., & Martínez L. J. M. Therapeutic Potential of Heparin in Sepsis Per Gram Negative. Journal of Medical

Research and Health Sciences. 2022; 5(4): 1881–1885.