

Cytological Diagnosis of Metastatic Lymph Node Lesions: An Observational Study

Amit Kumar¹, Jyoti Kumari²

¹Tutor, Department of Pathology, Bhagwan Mahavir Institute of Medical Science, Pawapuri, Nalanda, Bihar, India.

²Tutor, Department of Pathology, Bhagwan Mahavir Institute of Medical Science, Pawapuri, Nalanda, Bihar, India.

Received: 25-12-2022 / Revised: 20-01-2023 / Accepted: 03-02-2023

Corresponding author: Dr. Jyoti Kumari

Conflict of interest: Nil

Abstract

Aim: The aim of the study was to find out the cytological diagnosis of metastatic lymph node lesions.

Methods: The present study was conducted in the Department of Pathology, Bhagwan Mahavir Institute of Medical Science, Pawapuri, Bihar, India and study of all metastatic lymph node on FNAC samples reported over a period of nine months.

Results: The age of the patient ranged from 24 to 80 years with a mean of 60.4 years. The incidence was seen to peak at the age range above 60 years showing 144 cases (60%), followed by 84 cases (35%) in the age group 40 to 59 years. There were 12 cases (5%) below 40 years of age. The incidence of metastasis were more in female (144 cases, 60%) as compared to male (96 cases, 40%). The most common subtype of metastatic malignant tumor was adenocarcinoma and was observed in 156 cases. The metastatic lymph nodes were located in anterior and posterior cervical triangles, supraclavicular area, axilla, abdomen and inguinal region. The most common sites were the cervical triangles comprising 120 cases followed by 72 cases from supraclavicular nodes.

Conclusion: Cytology evaluation along with proper clinico-radiological correlation is quite useful in diagnosing metastasis with good degree of certainty. To conclude, in developing countries, like ours, FNAC is a cheap quick and reliable method to assess suspicious lymphadenopathy.

Keywords: Fine Needle Aspiration Cytology; Lymphadenopathy; Metastasis.

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

Introduction

The use of Fine Needle Aspiration Cytology (FNAC) for the diagnosis of metastatic malignancies in the lymph node is a well-established method. [1] Lymphadenopathy in an adult patient may be the first presenting clinical sign of non-hematologic malignancy. FNAC not only confirms the presence of metastatic

disease, but also gives the clue regarding the nature and origin of primary malignancy. FNAC is useful for the detection of recurrence and new metastasis. However, still a large percentage of cervical lymphadenopathy in adults turn out to be malignant. Cysts (congenital or acquired), abscesses, benign

and malignant tumors may mimic lymph node metastasis, especially with a case of known tumor. [2] Cystic metastasis or aspirate of low grade malignancies compose most of false negative cases. [3,4]

Lymphadenopathy in an adult patient may be the first presenting clinical sign of hematologic or non-hematologic malignancy. FNAC not only confirms the presence of metastatic disease, but also gives the clue regarding the nature and origin of primary malignancy. [5]FNAC is useful for the detection of recurrence and new metastasis. In developing country like India, infective lymphadenopathy is quite common, mostly due to high prevalence of tuberculosis. However, still a large percentage of cervical lymphadenopathy presenting in our hospital turn out to be malignant as this institute is a cancer care centre. Cysts (congenital or acquired), abscesses, benign and malignant tumors may mimic lymph node metastasis, especially with a case of known tumor.[6] Cystic metastasis or aspirate of low grade malignancies compose most of false negative cases.[7,8] This procedure is cheap, easily repeatable and well tolerated by the patients and can be performed on outpatient basis.[9]

In developing country like India, infective lymphadenopathy is quite common, mostly due to high prevalence of tuberculosis. However, still a large percentage of cervical lymphadenopathy in adults turn out to be malignant. Cysts (congenital or acquired), abscesses, benign and malignant tumors may mimic lymph node metastasis, especially with a case of known tumor.[6] Cystic metastasis or aspirate of low grade malignancies compose most of false negative cases.[7,8] This procedure is cheap, easily repeatable and well tolerated

by the patients and can be performed on outpatient basis.[9]

The aim of the study was to find out the cytological diagnosis of metastatic lymph node lesions.

Methods

The present study was conducted in the Department of Pathology, Bhagwan Mahavir Institute of Medical Science, Pawapuri, Bihar, India and study of all metastatic lymph node on FNAC samples reported over a period of nine months

Out of total 1500 cases of FNAC, 240 cases (16%) were of lymph nodes. Among these, there were a total of 48 nodes reported as “positive for metastasis” accounting for 20% of all lymph node FNACs and 2% of all FNAC cases. Other lymph nodes were reported as “reactive” and “infective” in 168 cases (70%), “positive for metastasis” in 48 cases (20%), “hematologic malignancy” in 5 cases (2%) and “unsatisfactory” in 19 smears (8%).

All FNACs were performed using a 22 or 23 gauge needle. An average of 1-2 passes and a minimum of 5 slides were made, Slides were routinely stained with both May Grunwald Giemsa and Papanicolaou (PAP) stains and wherever applicable, Periodic Acid Schiff stain was used. In case of deep seated lesions, Ultrasonography (USG) and Computerised Tomography (CT) guided FNAC was performed. Smears showing adequate cellular material was considered as “satisfactory” and were reported as “positive for metastasis” with further sub typing wherever possible. All the clinical and pathological data were collected and analyzed using Graphpad Prism version 5 software.

Results

Table 1: Demographic details

Age in years	N%
Below 40 years	12 (5%)
40-59 years	84 (35%)

Above 60 years	144 (60%)
Gender	
Male	144 (60%)
Female	96 (40%)

The age of the patient ranged from 24 to 80 years with a mean of 60.4 years. The incidence was seen to peak at the age range above 60 years showing 144 cases (60%), followed by 84 cases (35%) in the

age group 40 to 59 years. There were 12 cases (5%) below 40 years of age. The incidence of metastasis were more in female (144 cases, 60%) as compared to male (96 cases, 40%).

Table 2: Distribution of different pathological subtypes of metastasis

Pathological Subtypes	N%
Adenocarcinoma	156 (65%)
Squamous cell carcinoma	36 (15%)
Breast ductal carcinoma	12 (5%)
Papillary carcinoma thyroid	6 (2.5%)
Small cell carcinoma	6 (2.5%)
Non-small cell carcinoma	18 (7.5%)
Malignant melanoma	6 (2.5%)
Total	240 (100%)

The most common subtype of metastatic malignant tumor was adenocarcinoma and was observed in 156 cases. The metastasis had occurred from primary carcinomas in lung, stomach, colon and rectum. This was followed by squamous cell carcinoma (36 cases).

Table 3: Distribution of number of cases according to sites of lymphadenopathy

Sites of Lymphadenopathy	N%
Cervical triangles	120 (50%)
Supraclavicular	72 (30%)
Axillary	30 (12.5%)
Abdominal	3 (1.25%)
Inguinal	15 (6.25%)

The metastatic lymph nodes were located in anterior and posterior cervical triangles, supraclavicular area, axilla, abdomen and inguinal region. The most common sites were the cervical triangles comprising 120 cases followed by 72 cases from supraclavicular nodes.

Discussion

Fine needle aspiration cytology (FNAC) is used routinely as a first-line diagnostic test. The majority of studies reveal a malignant cause for palpable supraclavicular lymph nodes. [1,2] FNAC is a reliable diagnostic tool for lymphadenopathy in adult patients who are

is of considerable value in disease staging and documentation of metastasis in known and occult tumors. FNAC is a reliable diagnostic tool for lymphadenopathy in adult patients who are suspected for malignancy as it has less complication, is a simple invasive procedure and can be repeated easily. More than 90% of lymph node metastasis are diagnosed by initial aspiration. [9] Common metastatic tumors include malignancies from breast, oral cavity, thyroid, respiratory system, gastrointestinal tract, male and female genital tracts. [9]

suspected for malignancy as it has less complication, is a simple invasive

procedure and can be repeated easily. More than 90% of lymph node metastasis are diagnosed by initial aspiration. [6]

In the present study, adenocarcinoma was the most common metastatic tumor. In well differentiated adenocarcinoma, it showed cells with acinar and occasionally papillary arrangement and also singly scattered. The individual cells are usually large, cuboidal to columnar with moderate amount of cytoplasm and pleomorphic nuclei with prominent nucleoli. Cells even show vacuolated cytoplasm indicating intracellular mucin secretion. Background may show pink homogenous mucoid material if the mucin content of the tumor is high.

In other studies also the most common metastatic subtype were adenocarcinoma. [10,11] However, often it becomes difficult to distinguish between adenocarcinoma and poorly differentiated squamous cell carcinoma when the cell clusters show thick nuclear membrane and prominent nucleoli. [12,13] Cells with abundant clear cytoplasm also raise a suspicion of metastasis from the renal tumors. [14]

Metastatic squamous cell carcinoma was the second most common entity in our study. Tumor cells are seen mostly in sheets and singly scattered. The cells had dense cytoplasm with hyperchromatic nuclei in Giemsa stain and the cells show cytophilic or orangeophilic cytoplasm with pyknotic nuclei in PAP stain. In well differentiated squamous cell carcinoma, the tumor cells show individual cell keratinization. [14,15] The tumor cells often show necrotic material in the background. So in case of scanty cellularity with abundant necrotic material, a careful search for the tumor cells is required. Other studies showed squamous cell carcinoma as the most common metastatic tumor. [14,16]

Metastatic small cell carcinoma was seen in 12 cases where the patient had

suspicious mass lesion in the lung. Here the cells have scant cytoplasm with nuclei two to three times larger than small lymphocytes. Nuclei usually demonstrate the classical "salt and pepper" chromatin with indistinct nucleoli and frequent moulding. Streaking artefact along with karyorrhectic debris are seen in the background. [15,17] Sometimes, these background findings may make it difficult to differentiate from lymphoma where clinical findings (more generalised lymphadenopathy) may be helpful to differentiate. [14]

Melanomas can be seen anywhere in the body for example eyeballs, head, neck, great toe to name a few, and it is notorious to metastasize to any, specifically cervical or inguinal nodes. Our study showed 6 cases of metastatic melanoma, 3 each in cervical and inguinal lymph nodes. These smears showed discohesive pleomorphic cells with binucleate or multinucleate forms. The nuclei are large with characteristic prominent 1-2 macronucleoli. Intra and extracellular melanin pigment were seen both cases. Contrary with that other studies have observed melanin pigment in 25% of melanoma metastasis cases. [13,18]

Conclusion

Cytology evaluation along with proper clinico-radiological correlation is quite useful in diagnosing metastasis with good degree of certainty. To conclude, in developing countries, like ours, FNAC is a cheap quick and reliable method to assess suspicious lymphadenopathy. Furthermore, the histological architecture from cell blocks can be correlated with cytology, and such material can be used for appropriate histochemical and immunomarker studies, which can be useful in enhancing the diagnosis.

References

1. Steel BL, Schwartz MR, Ramzy I. Fine needle aspiration biopsy in the diagnosis of lymphadenopathy in 1,103

- patients. Role, limitations and analysis of diagnostic pitfalls. *Acta cytologica*. 1995 Jan 1;39(1):76-81.
2. Üstün M, Risberg B, Davidson B, Berner A. Cystic change in metastatic lymph nodes: A common diagnostic pitfall in fine-needle aspiration cytology. *Diagnostic cytopathology*. 2002 Dec;27(6):387-92.
 3. Verma K, Mandal S, Kapila K. Cystic change in lymph nodes with metastatic squamous cell carcinoma. *Acta cytologica*. 1995 May 1;39(3):478-80.
 4. Monchik JM, De Petris G, De Crea C. Occult papillary carcinoma of the thyroid presenting as a cervical cyst. *Surgery*. 2001 Apr 1;129(4):429-32.
 5. Khajuria R, Goswami KC, Singh K, Dubey VK. Pattern of lymphadenopathy on fine needle aspiration cytology in Jammu. *JK Sci*. 2006 Jul;8(3):157-9.
 6. Üstün M, Risberg B, Davidson B, Berner A. Cystic change in metastatic lymph nodes: A common diagnostic pitfall in fine-needle aspiration cytology. *Diagnostic cytopathology*. 2002 Dec;27(6):387-92.
 7. Verma K, Mandal S, Kapila K. Cystic change in lymph nodes with metastatic squamous cell carcinoma. *Acta cytologica*. 1995 May 1;39(3):478-80.
 8. Monchik JM, De Petris G, De Crea C. Occult papillary carcinoma of the thyroid presenting as a cervical cyst. *Surgery*. 2001 Apr 1;129(4):429-32.
 9. Thomas JO, Adeyi D, Amanguno H. Fine-needle aspiration in the management of peripheral lymphadenopathy in a developing country. *Diagnostic cytopathology*. 1999 Sep;21(3):159-62.
 10. Schafernack KT, Kluskens LF, Ariga R, Reddy VB, Gattuso P. Fine-needle aspiration of superficial and deeply seated lymph nodes on patients with and without a history of malignancy: Review of 439 cases. *Diagnostic cytopathology*. 2003 Dec;29(6):315-9.
 11. Nasuti JF, Mehrotra R, Gupta PK. Diagnostic value of fine-needle aspiration in supraclavicular lymphadenopathy: A study of 106 patients and review of literature. *Diagnostic cytopathology*. 2001 Dec; 25(6):351-5.
 12. El Hag IA, Chiedozi LC, Al Reyees FA, Kollur SM. Fine needle aspiration cytology of head and neck masses. Seven years' experience in a secondary care hospital. *Acta cytologica*. 2003 May 1;47(3):387-92.
 13. Fulciniti F, Califano L, Zupi A, Vetrani A. Accuracy of fine needle aspiration biopsy in head and neck tumors. *Journal of oral and maxillofacial surgery*. 1997 Oct 1;55(10):1094-7.
 14. Bagwan IN, Kane SV, Chinoy RF. Cytologic evaluation of the enlarged neck node: FNAC utility in metastatic neck disease. *Int J Pathol*. 2007;6(2):1-7.
 15. Singh HK, Silverman JF. Lung, chest wall and pleura. In: Orell SR, Sterrett GF, Whitaker D, editor. *Fine Needle Aspiration Cytology*. 4th ed. Elsevier: Churchill Livingstone; 2005; 245-9.
 16. Chute DJ, Stelow EB. Cytology of head and neck squamous cell carcinoma variants. *Diagn Cytopathol*. 2010; 38:65-80.
 17. Bedrossian CMW, Davila RM: Cytologic examination and fine needle aspiration. In Saldana MJ (ed.): *Pathology of pulmonary disease*. Lippincott-Raven, Philadelphia, 1994; 35.
 18. Schwarz R, Chan NH, MacFarlane JK: Fine needle aspiration cytology in the evaluation of head and neck masses. *Am J Surg*. 1990; 159:482-5.
 19. Pertuz J. G. V. Impaction Due to Sharp-Shaped Trauma in the Thoracic Abdominal Region: A Case Report. *Journal of Medical Research and Health Sciences*. 2022; 5(4): 1941-1945.