

Grading of Lymphocytic Thyroiditis Cytologically and Relationship to Hormonal Condition of the Thyroid

Bikash Kumar Singh¹, Bibhash Gogoi²

¹Assistant Professor, Department of Pathology, Lakhimpur Medical College, North Lakhimpur, Assam

²Associate Professor, Department of Pathology, Lakhimpur Medical College, North Lakhimpur, Assam

Received: 25-03-2023 / Revised: 18-04-2023 / Accepted: 30-05-2023

Corresponding author: Dr. Bibhash Gogoi

Conflict of interest: Nil

Abstract:

Background: The second most frequent thyroid lesion identified on FNAC after goitre is lymphocytic thyroiditis. Along with FNAC, other diagnostic factors include clinical characteristics, thyroid antibody titres, thyroid hormonal profile, and ultrasonography. The goal of the current study was to assess thyroid fine needle aspirates for the presence of lymphocytic thyroiditis, grade lymphocytic thyroiditis cases using predetermined cytological criteria, and determine the relationship between cytological grades and thyroid hormonal status.

Materials and Methods: From February 2021 to January 2022, this study was carried out at the Lakhimpur Medical College, Department of Pathology in North Lakhimpur, Assam. According to Bhatia et al., cytological criteria were used to further categorise lymphocytic thyroiditis patients that had been reported using the Bethesda system. Where data were available, cytological grades were associated with thyroid hormonal status.

Results: According to the Bethesda method, a total of 162 cases of lymphocytic thyroiditis were documented in our study. Further grading revealed that grade II thyroiditis accounted for the majority of cases (104, 64.2%), followed by grade I (48, 29.6%), and grade III (10, 6.2%). Out of the 78 cases with accessible thyroid hormonal profiles, the majority of patients (42; 53.85%) were hypothyroid. Euthyroid (21; 26.92%) and hyperthyroid (15; 19.23%) individuals were next in line. With a P value greater than 0.05, there was no statistically significant correlation between grades and hormone status.

Conclusion: The "gold standard" test for diagnosing lymphocytic thyroiditis is still the FNAC. However, in our investigation, there is no statistically significant link between the cytological grades and thyroid hormonal condition.

Keywords: Lymphocytic thyroiditis, Cytological grading, Thyroid hormonal status.

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

Introduction

With 42 million sufferers in India alone, thyroid disorders are among the most prevalent endocrine conditions globally.[1] The first line diagnostic method for assessing thyroid lesions is fine needle aspiration cytology (FNAC), which is capable of efficiently identifying neoplastic

and non-neoplastic lesions.[2] The majority of thyroid nodules are benign, and in about 60–70% of instances, the FNA results are interpreted as benign.[3] After benign follicular nodule, chronic lymphocytic thyroiditis is the thyroid lesion diagnosed on FNAC with the second-highest

frequency, and it is also the thyroid lesion with the second-highest frequency on FNA, behind goitre.[5] Hashimoto's thyroiditis (HT), also known as chronic lymphocytic thyroiditis or autoimmune thyroiditis, affects 30 to 60 people per 100,000 people annually and is more prevalent in women.[5] FNA has a diagnostic accuracy rate of 92%, making it very sensitive to the diagnosis of HT.6 The majority of patients initially have a symmetrical, widespread thyroid enlargement that is painless. In regions of the world where iodine is abundant, it is the most typical cause of hypothyroidism.[7] The diagnosis of this condition is aided by clinical characteristics, thyroid antibody titres, thyroid hormonal profile, and ultrasonography in addition to FNAC.[6, 8] The aims of the current study was to assess thyroid fine needle aspirates for the presence of lymphocytic thyroiditis, grade lymphocytic thyroiditis cases using predetermined cytological criteria, and determine the relationship between cytological grades and thyroid hormonal status.

Material and Methods

From February 2021 to January 2022, this study was carried out at the Department of Pathology, Lakhimpur Medical College, North Lakhimpur, Assam. Our department's records, thyroid fine needle aspirates from individuals who presented with thyroid enlargement during this time period were found. These patients stained smears with May-Grunwald-Giemsa (MGG) and Papanicolaou (PAP) were looked at under a light microscope.

According to the Bethesda System for Reporting Thyroid Cytopathology, the cases that were composed of numerous polymorphic lymphoid cells along with benign thyroid follicular cells and/or Hurthle cells were classified as lymphocytic thyroiditis (LT). It was not necessary to have a certain quantity of Hurthle or follicular cells to make the diagnosis of lymphocytic thyroiditis.[3]

These cases were further graded based on predefined cytological criteria as per Bhatia et al.[8] as:

1. Grade 0- No lymphoid cells.
2. Grade I (Mild)- Few lymphoid cells infiltrating the follicles/increased number of lymphocytes in the background.
3. Grade II (Moderate)- Moderate lymphocytic infiltration or mild lymphocytic infiltration with Hurthle cell change/giant cells/anisonucleosis.
4. Grade III (Severe)- Florid lymphocytic inflammation with germinal center formation, very few follicular cells left.

Where available, the thyroid hormonal profile of these patients was examined. Based on the patients' serum TSH, T4 and T3 readings, the patients were categorised as euthyroid, hypothyroid, and hyperthyroid. Where available, cytological grades and thyroid hormonal status were compared. Fischer Exact test of significance was used to statistically evaluate the relationship between grades and hormonal status, and a P value of <0.05 was deemed statistically significant.

Results

According to the Bethesda method, a total of 162 cases of lymphocytic thyroiditis were documented in our study. The age categories of 21–30 years (36; 22.22%), 41–50 years (32; 19.75%), and 31–40 years (48; 29.63%) had the highest proportion of patients, respectively. With ages ranging from 7 to 80, the mean age was 35.99 years. The study gender ratio was 9:125, with 146 (90.12%) female participants and 16 (9.88%) male participants.

These cases were then rated in accordance with Bhatia et al's standards. Grade II thyroiditis had the most cases, followed by grade I and grade III (Table 1) (Figures 1, 2, and 3).

Table 1: Distribution of lymphocytic thyroiditis cases into cytological grades

Lymphocytic Thyroiditis-cytological grading	No. of Cases	Percentage
Grade I (Mild)	48	29.6%
Grade II (Moderate)	104	64.2%
Grade III (Severe)	10	6.2%
Total	162	100.0%

Thyroid hormonal profiles were obtained for 78 out of 162 patients. The majority of patients (42; 53.85%) had hypothyroidism, which was followed by euthyroidism (21; 26.92%) and hyperthyroidism (15; 19.23%). Subclinical hypothyroidism was deemed to exist in cases with increased TSH but normal T4 and T3 levels (11; 14.10%). As demonstrated in Table 2, cytological grades and thyroid hormonal state were compared. The majority of individuals with thyroiditis of grade II also had hypothyroidism. Grade II thyroiditis, grade I thyroiditis, and no occurrences of

grade III thyroiditis were responsible for the majority of the euthyroid patients. There was no patient with grade III thyroiditis who had a normal hormonal condition and a thyroid profile was provided. There were eight cases of grade II lymphocytic thyroiditis associated with hyperthyroidism (Hashitoxicosis), six cases of grade I thyroiditis, and just one case of grade III thyroiditis (Table 2). However, there was no statistically significant link between the patients' thyroid hormone condition and cytological grades (P value = 0.21, greater than 0.05).

Table 2: Comparison of cytological grades with thyroid hormonal status

Cytological grade	Euthyroid	Hypothyroid	Hyperthyroid	Total
Grade I)	8	8	6	22
Grade II	13	32	8	53
Grade III	0	2	1	3
Total	21	42	15	78




Figure 1: Photomicrograph from a case of grade I lymphocytic thyroiditis showing a cluster of thyroid follicular cells infiltrated by few lymphocytes (MGG 400X).




Figure 2: Photomicrograph from a case of grade II lymphocytic thyroiditis showing mild lymphocytic infiltration with Hurthle cell change, anisonucleosis and increased lymphocytes in the background (MGG 400X).




Figure 3: Photomicrograph from a case of grade III lymphocytic thyroiditis showing florid inflammation comprising of polymorphic lymphoid cells, and few scattered follicular cells (MGG 400X).

Discussion

Hakaru Hashimoto, who bears his name, described the struma lymphomatosa, the first case of chronic thyroiditis.⁸ An autoimmune condition called Hashimoto's thyroiditis causes thyroid gland damage as well as gradual and increasing thyroid failure.⁷ The prevalence of HT appears to be rising recently. Ten times as many people now experience it as they did in the early 1990s. A high iodine diet, especially in coastal locations, has been related to this

rise in incidence.^[9,10] Characteristic cytological findings include the presence of lymphoid cells in the background and lymphoid cell infiltration of follicular cells. Hurthle cells, plasma cells, epithelioid cell granulomas, and multinucleated giant cells are examples of variable characteristics.^[11] Hypothyroidism typically comes on gradually. It may be preceded in some cases by temporary thyrotoxicosis, characterised by a decrease in TSH and an increase in free T4 and T3 levels. T4 and T3 levels drop as

hypothyroidism progresses, and TSH levels rise as a result.[7]

According to the Bethesda approach, thyroid fine needle aspirates were examined for cytological signs of lymphocytic thyroiditis, and 162 cases were identified as such in the current study. Our study's most prevalent age range for these individuals was 31 to 40 years old, which is consistent with research by Bhatia et al.,[8] Anila et al.,[6], and Shetty et al.[12] In contrast to a western study, where the median age of the individuals was 58 years old, these Indian studies demonstrate a younger age at which lymphocytic thyroiditis is diagnosed in the Indian population.[13] Female preponderance was comparable to that of the majority of research in the literature in the current investigation.[5, 6, 8, 12]

The majority of patients in our study had grade II thyroiditis, which was followed by grade I thyroiditis, while grade III thyroiditis affected the fewest patients. Studies by Bhatia et al.[8] and Shetty et al.[12] are comparable to this one. However, the majority of patients in a study by Anila et al.[6] had grade I thyroiditis, whereas the majority of cases in a study by Sood et al.[5] were grade III thyroiditis.

When thyroid hormone status was evaluated, the majority of individuals were hypothyroid, which is consistent with many other studies.[8,12,14] In our investigation, there were 11 (14.1%) patients with subclinical hypothyroidism. Shetty et al study[12] found that, greater than in our study, 39.6% of the cases had subclinical hypothyroidism. No statistically significant link was identified in our study when cytological grades were further correlated with patients' thyroid hormonal status, which is similar with many other studies in the literature.[6, 8, 12] Grade 3 lymphocytic infiltration does, however, statistically correlate with either TPO and TSH together or TSH alone, according to a research by Sood et al. The relationship between hormonal state and lymphocytic infiltration

was statistically significant (P- 0.02) in a research by Kumar et al. Hypothyroidism was more common (44%) than hyperthyroidism (6%), but there was no statistically significant link between the two (P- 0.36; P- 0.29), even among individuals with moderate to extensive infiltration. The frequency of individuals with hyperthyroidism was highest (47%) and statistically significant (P- 0.0006) among those with little invasion. The statistical relevance between the cytological grades of LT and TSH was found in a study by Megalamane et al.[16] (p-value <0.001). Clinical, radiographic, and hormonal criteria are overlapped in Hashimoto's thyroiditis and other thyroid pathologies, making them ineffective.[15] However, due to local antibody generation by intrathyroidal lymphocytes without spillover into the blood, around 7 to 33% of cases of Hashimoto's thyroiditis are antibody negative, particularly in the early stages of the disease. Therefore, FNAC is preferred as well as more economical than antibody screening to detect Hashimoto's thyroiditis,[10,17] especially in the early stages of the disease and if antibody testing facilities are not accessible, such as in our institution, or are prohibitively expensive.[15]

However, there are certain difficulties with using FNAC to diagnose lymphocytic thyroiditis. Smears with cytological signs of hyperplasia, such as in Grave's disease or copious colloid, are prone to miss the diagnosis of Hashimoto's thyroiditis.[9] It may not be able to differentiate between pure Grave's disease and Hashimoto's thyroiditis in the toxic phase, or Hashitoxicosis, based solely on cytological findings; this requires comparison with clinical information and hormonal profile.[11] Adenomatous/colloid/hyperplastic goitre is the primary differential diagnosis in I2 deficient regions when smears demonstrate hyperplastic follicular cells, Hurthle cells, a small population of

background lymphocytes, and the absence of lymphocytes in epithelial groups.[15]

Selective sampling from a hyperplastic nodule with a predominance of Hurthle cells and little to no lymphocytic infiltrate in the early stages of Hashimoto thyroiditis raises the possibility of a Hurthle cell tumour.[3] While poorly coordinated and cohesive cell clusters favour neoplasia, Hurthle cells in flat sheets favour thyroiditis.[11] The likelihood of lymphoma may increase if lymphoid components predominate, as in grade III thyroiditis.[12] Approximately 75% of thyroid gland primary lymphomas develop against a background of Hashimoto's thyroiditis.[11] It is possible to distinguish these patients from Non-Hodgkin lymphoma thanks to the polymorphic lymphoid population.[3,8] In cases when the diagnosis is unclear morphologically, immunophenotyping tests are critical to support the diagnosis. There is evidence that the incidence of cancer in HT can range from 0.4% to 28%. The likelihood of papillary cancer should be slightly increased since the follicular or Hurthle cells in LT occasionally exhibit modest atypia, nuclear expansion, grooves, and chromatin clearance.[3,12]

As a result, some of the potential pitfalls and uncertainties can be reduced, and the value of FNAC in the diagnosis of lymphocytic thyroiditis can be increased. This is because it is important to be aware of how cytological features of one thyroid pathology may overlap with those of another, and by using an integrated approach with clinicopathologic correlation.

Conclusion

Although the diagnosis of lymphocytic thyroiditis is aided by a combination of cytomorphology, clinical characteristics, thyroid hormonal profile, antibody testing, and ultrasonographic features, particularly in cytologically ambiguous cases, FNAC continues to be the "gold standard"

investigation. However, in our investigation, there is no statistically significant link between the cytological grades and thyroid hormonal condition.

References

1. AmbikaGopalakrishnan Unnikrishnan UshaV Menon Thyroid disorders in India: An epidemiological perspective Indian J Endocr Metab 2011.
2. Santosh Kumar Mondal Simanti Sinha Bijan Basak Dipanwita Nag Roy Swapan Kumar Sinha The Bethesda system for reporting thyroid fine needle aspirates: A cytologic study with histologic follow-up J Cytol 2013.
3. S Z Ali E S Cibas The Bethesda System for Reporting Thyroid Cytopathology. Definitions, Criteria and Explanatory Notes 2nd ed Springer Switzerland 2018.
4. Payal Mehra Anand Kumar Verma Thyroid Cytopathology Reporting by the Bethesda System: A Two-Year Prospective Study in an Academic Institution Pathol Res Int 2015. 2090-8091, 2042-003X10.
5. Neelam Sood Jitendra Singh Nigam Correlation of Fine Needle Aspiration Cytology Findings with Thyroid Function Test in Cases of Lymphocytic Thyroiditis J Thyroid Res 2014, 2090-8067, 2042-007210.
6. KR Anila Nileena Nayak K Jayasree Cytomorphologic spectrum of lymphocytic thyroiditis and correlation between cytological grading and biochemical parameters J Cytol 2016.
7. A Maitra V Kumar A K Abbas J C Aster The endocrine system Robbins and Cotran Pathologic Basis of Disease II Reed Elsevier India Private Limited 2014, 108589.
8. Alka Bhatia Arvind Rajwanshi Radharaman J Dash Bhagwant R Mittal Akshay K Saxena Lymphocytic thyroiditis – is cytological grading significant? A correlation of grades with clinical, biochemical, ultrasonographic and radionuclide parameters Cyto J 2007, 4100974-596310.

9. B N Gayathri M L Harendra Kumar R Kalyani Prasad K Krishna Fine needle aspiration cytology of Hashimoto's thyroiditis - A diagnostic pitfall with review of literature *J Cytol* 2011, 28421030970-9371.
10. Faiyaz Ahmad Ashutosh Kumar Jyotsana Khatri Ankita Mittal Seema Awasthi Shyamoli Dutta Cytological Diagnosis of Hashimoto's Thyroiditis Revealing the Increased Frequency than Expected: A Retrospective Study of 750 Thyroid Aspirates *Int J Med Res Professionals* 2016, 2314362454-6356, 2454-6364.
11. S R Orell G F Sterrett Orell & Sterrett's Fine Needle Aspiration Cytology 5th ed RELX India Private Limited New Delhi 2012, 11855.
12. Vijaya Chowdappa Archana Shetty Cytomorphological spectrum of Hashimoto's thyroiditis and its correlation with hormonal profile and hematological parameters *J Cytol* 2019, 363137410970-9371.
13. M. P. J. Vanderpump W. M. G. Tunbridge J. M. French D. Appleton D. Bates F. Clark The incidence of thyroid disorders in the community: a twenty-year follow-up of the Wickham Survey *Clin Endocrinol* 1995, 43155680300-0664, 1365-2265.
14. Shirish S. Chandanwale Rahul Nair Anushree Gambhir Supreet Kaur Aditi Pandey Abhinav Shetty Cytomorphological Spectrum of Thyroiditis: A Review of 110 Cases *J Thyroid Res* 2018, 2090-8067, 2042-007210.
15. N Kumar C Ray S Jain Aspiration cytology of Hashimoto's thyroiditis in an endemic area *Cytopathol* 2002, 1313190956-5507, 1365-2303.
16. Supreetha Megalamane Hemalatha Anantharamaiah Shashidhar Nagaraj Kurpad Correlation of Cytological Grading in Lymphocytic Thyroiditis with Thyroid Hormones and Antibodies - A Retrospective Study in the Era of Bethesda System of Thyroid Reporting *J Clin Diagn Res* 2018, 1281042249-782XJCDR.
17. C P Madhu V Swetha Fine Needle Aspiration Cytology of Hashimoto's Thyroiditis: Its Diagnostic Limitation *IJSS J Surg* 2017, 31279.