

Effect of Fenugreek (Methi) on Blood Pressure, Blood Sugar, Serum Cholesterol and Serum Triglycerides in Known Cases of Mild to Moderate Essential Hypertension.

Dhaneshwar Tripathi¹, Jitesh Aggarwal², Shokat Ali Bohra³, Krupa Akhiani⁴, Himani Dhingra⁵

¹Assistant Professor, Department of General Medicine, GBH General Hospital, Udaipur, Rajasthan, India

²Associate Professor, Department of General Medicine, GBH General Hospital, Udaipur, Rajasthan, India

³Associate Professor, Department of General Medicine, GBH General Hospital, Udaipur, Rajasthan, India

⁴Resident Doctor, Department of General Medicine, GBH General Hospital, Udaipur, Rajasthan, India

⁵Resident Doctor, Department of General Medicine, GBH General Hospital, Udaipur, Rajasthan, India

Received: 25-11-2023 / Revised: 25-12-2023 / Accepted: 30-01-2024

Corresponding author: Dr. Surekha Kadadi

Conflict of interest: Nil

Purpose: This study is carried out to assess the effect of FENUGREEK(METHI) on blood pressure, blood sugar, serum cholesterol and serum triglycerides in known cases of mild to moderate essential hypertension patient.

Methods: A total of 20 patients of mild to moderate hypertension was taken to study the effect of fenugreek in their blood pressure, blood sugar, serum triglycerides and serum cholesterol level. Initially a detailed clinical examination of patients was carried out and all the drugs were discontinued for 2 weeks(washout period). All the patients were administered 2.5gm of methi powder twice a day after food with a glass of water daily for 2 months

Conclusion: We concluded that patient of mild to moderate hypertension have no significant effect of FENUGREEK.

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

Introduction:

Susruta as early as 700 BC has described the medical use of 700 plants for the treatment of various disorders. [1] Incidentally many of these plants in some form or other are commonly used in Indian diet. Many of them recommended as disinfectant for the intestine, some are known to have important antibiotic properties and tonics. Tulsi, Methi, Haldi, Ginger are considered as outstanding medicines in ancient literature. [2] There is certainly a tenable rationale explaining the relationship between dietary fat and blood pressure in that linoleic acid is a potential precursor of prostaglandins, which in turn affect renal sodium excretion and vascular reactivity. [3] Looking to the important medicinal value and extensive use of some of these condiments, the present study has been planned to observe the effect of Fenugreek (METHI) on blood sugar, serum triglycerides, serum cholesterol and blood pressure in known patients of mild to moderate essential hypertension. [4-9]

Material and Methods

The study was conducted in 20 patients of mild to moderate essential hypertension. All the patient after their washout period were administered 2.5gm of methi powder twice a day after food with a glass of water daily for 2 months.

The blood samples were collected after overnight fast for blood sugar, cholesterol and triglycerides and 2 hours. These samples were collected before administration of drug, at 4 weeks and at 8 weeks. The blood pressure was measured in ideal resting condition in supine position and standing position before the administration of drug and thereafter at interval of 2 weeks for 8 weeks. The patients were allowed their routine activity and usual diet they were taking.

The blood samples were analysed for:

1. Blood sugar

- 2. Serum cholesterol and
- 3. Serum triglycerides

- 2. Patients having established essential hypertension with diastolic pressure above 90 mmHg and below 114mg of Hg

Criteria for the selection of hypertension patients:

Observation:

- 1. Adult of either sex between the age group 18-70 years.

Effect of Fenugreek(Methi) on Blood Pressure in Essential Hypertensive Patients





Figure 2. Mean (MBP), Systolic (SBP) and Diastolic (DBP) blood pressure in T2DM patients after successive intake of 2 g of fenugreek and their conventional drugs. (*: significant difference in comparison to baseline at 5%).


As the table shows out of 20 patients,9 patients showed slight increase in systolic and diastolic blood pressure. However, in 11 patients decrease in systolic as well diastolic blood pressure was observed. Thus the administration of fenugreek has no effect on mild to moderate hypertension.

Effect of fenugreek on blood sugar (fasting) in essential hypertension patient



According to above table the mean initial fasting blood sugar level was 122mg/dl which decrease to 118mg/dl. However, the decrease is statistically not significant.

Effect of Fenugreek on Serum Cholesterol and Serum Triglycerides Level in Essential Hypertensive Patient.



As the bar chart shows that administration of fenugreek had no significant effect on serum cholesterol level and serum triglycerides.

Summary and Conclusion:

The present study was conducted in 20 patients of mild to moderate essential hypertension. These were observed for blood pressure, blood sugar, serum cholesterol and triglycerides before and after 2 months of 2.5 gms of powdered fenugreek for 8 weeks.

Thus from the study it is deduced that fenugreek has no significant effect in patient of mild to moderate hypertension. The mechanism of action was not field of study in the project, however it can be postulated that the presence of fibre in fenugreek and other component like compact amylase, tightly bound carbohydrate, antinutrient like phytates, lectine and enzyme inhibitors may be having hypoglycaemic factor.

Bibliography

1. Bordia A., Sharma K.D., Parmar V.K. and Verma S.K.: Protective effect of garlic oil on

the change produced by three weeks of fatty diet on serum cholesterol, triglycerides, fibrinolytic activity and platelet adhesiveness in man. Indian heart Journal 1982,43:86.

2. The wealth of India. CSIR New Delhi 1976, X:305.
3. Nadkarni K.M.: India Metria Medica 1976,1:1240.
4. Vaishay Umadilal: Uses of methi 1974,1:1
5. Bowman D.E. Amylase inhibitor of navy beans. Science 1945,102:358.
6. Reid and Meier: Photochemistry, 1970; 9:513. Quoted from Wealth of India 1976, X:303.
7. Andrew et al.: J. chem. Soc.1952, 2744. Quoted from wealth of India 1976, X:303.
8. Kolousek and Coulson: J.Sci. Fd. Agric.1955,6,203. Quoted foom wealth of India1976, x:303.
9. Chakroborty, B., Parvin, S., Hossain, M. M., & Hossain, M. J. Self- Examination of Breast of the Students of Nursing College in Bangladesh. Journal of Medical Research and Health Sciences, 2022;5(12): 2339–2344.