

# Longitudinal Study of Operative Time, Intra-Op and Post-Op Bleeding of Transanal Suture Rectopexy for Hemorrhoids with Conventional Milligan Morgan Open Hemorrhoidectomy

Deepak Thakur<sup>1</sup>, Deepshikha Mishra<sup>2</sup>

<sup>1</sup>MBBS, MS, Department of General Surgery, SVNGMC Yavatmal, Maharashtra.

<sup>2</sup>MBBS, MS, Department of ENT, SVNGMC, Yavatmal, Maharashtra.

Received: 25-11-2023 / Revised: 23-12-2023 / Accepted: 26-01-2024

Corresponding Author: Dr. Deepshikha Mishra

Conflict of interest: Nil

## Abstract:

To compare the transanal suture rectopexy (TR) with conventional Milligan-Morgan hemorrhoidectomy (MH) in patients of grade II, III and IV hemorrhoids.

**Methods:** From March 2018 to October 2019, 60 patients of hemorrhoids from grade II to IV were randomly selected to undergo either the transanal suture rectopexy (n= 30) or the Milligan-Morgan hemorrhoidectomy (n=30). Results were analyzed during postoperative period and at 1 month. Operative time, post-operative complications, resolution of symptoms and recurrence were taken as variables. Data was analyzed by Open EPI version 2.3.

**Results:** Both the groups were comparable in terms of demographic data. Bleeding per rectum was the major symptom. TR group had more operative time and less intra-op blood loss, post-operative bleeding, pain, anal incontinence, hospital stay and recurrence than MH group. Data was statistically significant (p-value<0.05). In MH group, 16 patients developed postoperative bleeding while this was only 2 in TR group. In MH group 2 patients developed anal incontinence while there were none in TR group. In MH group 6 patients came with prolapse and 7 patients came with bleeding per rectum at 1 month follow up, while count was only 1 in TR group.

**Conclusion:** Transanal suture rectopexy is unique and simple stitching technique appropriate for treating all the grades of hemorrhoids. It offers cheaper, ambulatory and effective alternative to costly newer modalities for the treatment of hemorrhoids. It is better accepted by the patients. More studies are needed to declare it as a standard procedure for the cure of haemorrhoids.

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

## Introduction

Haemorrhoids, commonly called as piles are common ailments that present to a surgeon requiring hospital admissions. [1] Haemorrhoids are seen in the anal canal mainly as the bulged blood vessels around the supporting tissues. The main cause of haemorrhoids is considered to be constipation. [2] Haemorrhoids are classified into grades I-IV (or first to fourth degree) by Goligher. [3] Grade I patients are managed conservatively while grade II-IV require surgical intervention most of the time. Hemorrhoidectomy is the most definitive management for grade II-IV haemorrhoids. [4]

Hemorrhoidectomy is a routinely performed operation in the anorectal region. It involves excising both (internal & external) components of each hemorrhoid. In traditional hemorrhoidectomy, postoperative pain, the risk of complications such as post-operative bleeding, anal incontinence, wound infection, and the length of recovery time are challenges in the management of patients with

hemorrhoids. [5] Thus, there is need of better surgical technique in management of hemorrhoids. In this study we are comparing the outcome of a new technique of transanal suture rectopexy with the conventional open Milligan-Morgan hemorrhoidectomy. Only a few studies have been done on this new technique, so this study becomes important.

## Aims and Objectives

To compare transanal suture rectopexy for haemorrhoids with the conventional Milligan-Morgan hemorrhoidectomy with respect to operative time, intra-operative bleeding and post-operative bleeding

## Study Design and Methodology

This study was done at a tertiary level teaching hospital of the state of Maharashtra in India. It was a longitudinal study done between March 2018 and October 2019. All consenting patients admitted in

the general surgery department with haemorrhoids (grade II, III and IV) were included in the study. 60 patients were enrolled in the study, which were divided in two groups of 30 each by random table number method, as Transanal suture rectopexy (TR) and Conventional Milligan-Morgan open haemorrhoidectomy (MH) groups.

A complete detailed history, physical evaluation was done and the patients were informed about the details of the procedure and written informed consent was obtained. Diagnosis of haemorrhoids was done by per-rectal examination and proctoscopy. Haemorrhoids were graded as per Goligher classification. Inclusion criteria were grade II, III and IV haemorrhoids. Exclusion criteria included patients with previous anorectal surgery, anal fissure or fistula, acute thrombosed external hemorrhoids, anal stenosis, anal incontinence, rectal prolapse, malignancy, inflammatory bowel diseases, irritable bowel syndrome, previous pelvic radiotherapy.

### Operational Definitions

In transanal suture rectopexy, under spinal anaesthesia, in lithotomy position, anal canal is lubricated with lignocaine 2% jelly. By using Sim's speculum, the piles masses pushed upwards and replaced at their original anatomical position. Self-illuminating proctoscope with sliding valve and slit is introduced in the anal canal. Sliding valve is withdrawn and the dentate line is identified. 4 cm, first stitch is tied at 3'o clock position. Next stitches are double locked and 1-2 mm overlapping at the same level circumferentially. This prevents purse string effect. Each stitch is of submucosal depth and only a part of rectal muscle is taken. Stitches should not pass through and through the rectal wall. This results in fixation of mucosa, submucosa and protruding pile masses to rectal wall. The second suture line is applied circumferentially at 2 cm above the pectinate line. For suturing a 30 mm, 5/8 circle, 2-0 polyglactin round body suture is used. [6-7]

In conventional Milligan-Morgan haemorrhoidectomy, under spinal anaesthesia, in lithotomy position, anal canal is massaged with lignocaine 2% jelly and digital dilation is done. The internal haemorrhoids are pulled down by using babcock's forceps, this exposes the haemorrhoidal pedicles. After traction is applied on the haemorrhoid, transfixation and

ligation of the pedicle is done by 2-0 vicryl, with the knot tied on the lumen side. By using cautery, the pedicle is cut, leaving sufficient cuff.

The other haemorrhoids are removed in a similar fashion leaving intact bridge of perianal skin and anal mucosa between each dissection side, should be not less than one centimeter wide. This prevents post-operative anal stenosis. [8-9]

**Post-operative care:** Patients were managed in the surgery ward. Oral antibiotics and analgesics were given. The patients were discharged on the next day, if there were no complications. Post-operative complications were noted before discharge

### Statistical Analysis and Results

Data was collected from hospital records and case files of the patients. The data was entered in Excel sheets and was analyzed with the help of Open EPI version 2.3 through its statistical program, using mean and standard deviation for quantitative data like operative time, post-operative bleeding, and hospital stay. Unpaired t-test was used for quantitative data. Fisher exact test was used for qualitative data. P-value of  $\leq 0.05$  was considered statistically significant.

In patients undergoing Milligan Morgan procedure (MH), mean age was  $42.06 \pm 6.78$  and while, it was  $45.13 \pm 6.32$  for patients undergoing transanal suture rectopexy (TR). There were 48 males and 12 females. Male: Female ratio was 4:1. In MH group, 25 were male and 5 were female and TR group, 23 were male and 7 were female. It was observed that in both the groups, there was male preponderance of cases.

Out of total 60 patients, 12 were diagnosed to have grade II disease, 38 patients presented with grade III disease and remaining 10 belonged to grade IV disease. 53 out of 60 patients complained of bleeding per rectum, which the most common presenting complaint. 50 out of 60 patients suffered from constipation. 48 out of 60 patients complained about prolapsed piles.

Pain was complained by 37 out of 60 patients. All the patients were given spinal anaesthesia. There were 4 patients of grade II, 19 patients of grade III and 7 patients of grade IV haemorrhoids in MH group. In TR group, this count was 8, 19 and 3 for grade II, III and IV haemorrhoids respectively.

**Table 1: Analysis of Demography**

Characteristics	Milligan-Morgan	Transanal Suture Rectopexy
Age In Years	42.06 ± 6.78	45.13 ± 6.32
Gender	M=25; F=5	M=23; F=7
Grade	II-4; III-19; IV-7	II-8; III-19; IV-3

Mean operative time for MH and TR groups was  $34.36 \pm 1.13$  min and  $37.4 \pm 4.18$  min respectively. Calculated p-value was 0.0000001 (significant). Statistically, it can be concluded that TR has more operative time as compared to MH. Intra-operative bleeding was calculated by gravimetric method (weighing pre and post-operative blood soaked pads). Mean blood loss for MH and TR groups was  $62.66 \pm 10.8$  ml and  $37.66 \pm 7.27$  ml respectively.

Calculated p-value was 0.03629 (significant). It can be concluded that TR has less intra-operative bleeding as compared to MH group.

**Table 2: Analysis of Quantitative Data**

Variables	Group	Mean	Standard Deviation	P-Value
Operative Time	Milligan Morgan	34.36	1.13	<0.05
	Transanal Suture Rectopexy	37.4	4.18	
Intra-Operative Bleeding	Milligan Morgan	62.66	10.8	<0.05
	Transanal Suture Rectopexy	37.66	7.27	

(Unpaired t-test; results obtained from Open EPI, Version 3).

In this study, 18 patients had bleeding in post-operative period (before discharge), out of which 16 belonged to MH and only 2 belonged to TR group. Calculated p-value was 0.00007236 (significant). It was concluded that MH patients have higher post-op bleeding than TR patients. Post-op bleeding was managed conservatively by gauze packing and laxatives.

**Table 3: Analysis of Qualitative Data**

Variables	Group	Frequency	Percentage	P-Value
Bleeding In Post-Op Period	Milligan Morgan	16	53.3	<0.05
	Transanal Suture Rectopexy	2	6.6	

(Fischer exact test; results obtained from Open Epi, Version 3)

## Discussion

Hemorrhoids or piles are one of the most ubiquitous conditions that afflict problem for general population but it is difficult to detect its exact prevalence in the population, owing to its social stigma. [10] Various treatment modalities are present for hemorrhoids but high cost, post-op complications and recurrence of pre-operative symptoms are the limiting factors. [10] We require a surgical technique which is cheap, with minimal post-operative complications and negligible recurrence rate.

At 3, 7 and 11'o clock, in lithotomy position, three anal cushions are present. They are nothing but clusters of dilated veins of superior rectal venous plexuses. The symptoms of hemorrhoids are due to prolapse of these anal cushions. [6]

All the earlier surgeries for hemorrhoids aim at ligating the blood vessel supplying the prolapsed pile masses or excision of pile mass. When we ligate the blood vessel at one point, in due course of time collaterals develop, which results in recurrence rate of about 18-25%. [6-7] In this new modality for the treatment of all the grades of hemorrhoids, a simple suturing technique is applied to achieve the required results. Two circumferential suture lines are taken in pain free area, 2 cm and 4 cm proximal to the dentate line. Sutures are of submucosal depth so that only a part of rectal muscle layer is taken to anchor the lax pile mass along with anal cushions.

Our aim was to compare a new modality Transanal suture rectopexy with the conventional Milligan Morgan open hemorrhoidectomy. Transanal suture rectopexy proved to be significantly better than the Milligan Morgan open hemorrhoidectomy with respect to intra-operative blood loss and control of bleeding per rectum in post-op period but it has more

operative time. It is cheaper than other procedures because no additional sophisticated instrument is required. [6-7]

## Conclusion

Transanal suture rectopexy is new modality for treatment of haemorrhoids. It is a simple stitching technique whose learning curve is lesser than other new procedures. Best thing is that when it is applied to different grades of haemorrhoids, it shows similar excellent results. The grade of hemorrhoids and the surgeon do not have effect on result and complications. This new modality is successful in controlling bleeding per rectum and prolapsed pile masses. Conventional techniques are associated with post-operative complications. This new surgery has more operative time with respect to Milligan-Morgan haemorrhoidectomy but, it has lesser intra-op blood loss and negligible post-operative bleeding. This new procedure is an attractive option for both surgeon and patients. Use of Chivate's proctoscope makes this surgery easier for surgeons. The new method still requires more studies to determine long term outcomes and complications.

## References

- Hollingshead J. R. F. & Phillips R. K. S. Haemorrhoids: Modern diagnosis and treatment. Postgraduate Medical Journal 2015; 92(1083)4-8.
- Navarra, L., R. Pietroletti, G. Maggi, S. Leardi, and M. Simi. "Diagnosis and treatment of haemorrhoids in the elderly: results from 291 patients." Techniques in Coloproctology 3.3 1999; 127-130.
- Ohning, Gordon & Machicado, Gustavo & Jensen, Dennis. Definitive Therapy for Internal Hemorrhoids-New Opportunities and Options.

- Reviews in gastroenterological disorders. 2009; 9: 16-26.
4. Pattana-arun J, Wesarachawit W, Tantiphlachiva K, Atithansakul P, Sahakitrungruang C, Rojanasakul A. A comparison of early postoperative results between urgent closed hemorrhoidectomy for prolapsed thrombosed hemorrhoids and elective closed hemorrhoidectomy. *J Med Assoc Thai.* 2009;92(12):1610-5.
  5. Danson Y, Kok-Yang Tan. Hemorrhoidectomy-making sense of the surgical options. *World J Gastroenterol.* 2014 Dec 7; 20(45):16976–16983.
  6. Chivate, S.D., Ladukar, L., Ayyar, M. et al. *Indian J Surg* (2012) 74: 412.
  7. Jain, Amit & Khan, Irfan & Poonia, Kuldeep & Jain, Richa. (2016). Comparison of Transanal Suture Rectopexy with Stapled Haemorrhoidopexy in Patients of Haemorrhoids: A Randomized Control Trial. *IOSR Journal of Dental and Medical Sciences.* 2016;15: 112-116.
  8. Milligan ETC, Morgan C, Naughton Jones, Office RR 1937” *Surgical anatomy of the anal canal and the operative treatment of haemorrhoids” Lancet* ii.1119.
  9. Milligan ETC 1939” *Hemorrhoids” BMJ* 2:412.
  10. A. Farook, A. Periasamy, S. Arvind, I. Kannan. Study on prevalence and surgical management of haemorrhoid’s in a tertiary care teaching hospital in a rural area: *Int Surg J.* 2018 Apr; 5(4):1263-1266.