

**Prevalence of Thyroid Dysfunction and Non-Alcoholic Fatty Liver Diseases (NAFLD) in Patients with Type 2 Diabetes Mellitus**Sunil Aren<sup>1</sup>, Dinesh Kumar Gurjar<sup>2</sup>, Ajeet Singh<sup>3</sup><sup>1,2,3</sup>Department of General Medicine, Shree Jagganath Pahadia Govt. Medical College, Bharatpur, Rajasthan, India

Received: 25-01-2024 / Revised: 23-02-2024 / Accepted: 18-03-2024

Corresponding Author: Dr Dinesh Kumar Gurjar

Conflict of interest: Nil

**Abstract:****Background:** The aim of this study was to assess the level of thyroid dysfunction in patients with type 2 diabetes mellitus and to identify the association of thyroid dysfunction with diabetic complications.**Methods:** All patients were evaluated for thyroid status; assessment of T3, T4 and TSH levels and Blood glucose levels and HbA1c.**Results:** In present study, 22.00% patients were present with subclinical hypothyroidism, 17.33% patients were present with hypothyroidism and 3.33% patients present with hyperthyroidism.**Conclusion:** Screening for thyroid disease among patients with diabetes mellitus should be routinely performed for early detection and treatment of thyroid dysfunction to delay the complications of diabetes.**Keywords:** Thyroid, Type-2DM, Complication.

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

**Introduction**

One of the oldest diseases known to man is Diabetes mellitus (DM). It's first case was reported in the Egyptian manuscript dating to about 3000 years ago. In 1936, Type 1 and Type 2 DM were clearly distinguished. [1] Diabetes mellitus is a collection of common metabolic disorder mainly considered by hyperglycaemia which results, commencing from defective insulin secretion or insulin action or both together. [2]

Patients with type 2 diabetes mellitus are more prone to develop thyroid disorders. Many diabetic patients show features of thyroid dysfunction over a period of time. [3] Insulin resistance plays an important role in the development of hypothyroidism in patients with type 2 diabetes mellitus. Hypothyroidism in diabetic patients leads to an aggravation of dyslipidemia, hypertension and cardiovascular disease.[3] Thus, it is necessary to recognize and treat hypothyroidism in diabetic patients to prevent worsening of diabetic complications.[4] Hypothyroidism can be diagnosed with the help of a simple blood test which is readily and easily available. This can be performed by the primary care physician involved in treating diabetic patients. Early treatment of thyroid dysfunction in diabetic patients will help in normalizing their glycemic status and lipid profile.

**Materials and Methods**

- Type of Study: A Hospital based study

**Inclusion Criteria:** All patients with Type 2 diabetes Mellitus.**Exclusion Criteria:**

- Patients with history of alcohol consumption for any duration of time were excluded.
- Persons with previous history of jaundice, ascites and signs of liver cell failure will be excluded.
- Patients with history of intake of Methotrexate, Amiodarone, Glucocorticoids, Synthetic estrogens, Nucleoside Analogues (ddI, AZI) or any other hepatotoxic drug will be excluded.
- Patients with history of Chronic Renal Failure and Ischemic Heart Disease.
- Pancreatitis.
- Steroid induced Diabetes will be excluded from the study.

**Methodology**

1. A detailed history and examination was conducted as per the proforma.
2. All patients were undergo evaluation for diabetes.

- All patients were evaluated for thyroid status; assessment of T3, T4 and TSH levels and Blood glucose levels and HbA1c.
- The laboratory evaluation of thyroid functions were conducted by estimation of serum T3, T4 and TSH levels by chemi-luminescence assay method. 2 ml of blood would be drawn and will be centrifuged and the serum (500microml) collected from that and will be incubated with the

reagent (separate for T3, T4 and TSH) for about 1 hour at room temperature.

- HbA1c and FBS levels was checked.

### Results

In our study, maximum patients (56.00%) were from less than 46-60 Yrs followed by 22.67% patients were less than 45 yrs and 21.33% patients were more than 60 yrs age. 54.00% patients were female and 36.00% patients were male.

**Table 1: Blood sugar wise distribution**

Variable	Mean	SD
FBS (mg/dl)	152.70	79.06
PPBS (mg/dl)	239.71	114.08
Hb1Ac (%)	7.00	1.61
Free T3 (pg/ml)	2.38	0.88
Free T4 (ng/ml)	1.35	0.49
TSH ( $\mu$ U/L)	4.39	6.46

In present study, FBS was  $152.70 \pm 79.06$  mg/dl, PPBS was  $239.71 \pm 114.08$  mg/dl and Hb1Ac was  $7.00 \pm 1.61\%$ . In present study, free T3 was  $2.38 \pm 0.88$  pg/ml, free T4 was  $1.35 \pm 0.49$  ng/ml and TSH was  $4.39 \pm 6.46$   $\mu$ U/L.

**Table 2. Distribution of various thyroid parameters**

Thyroid biochemical parameters	No of cases	Percentage	
Free T3 (pg/ml)	Low (<1.4)	7	4.67
	Normal (1.4 to 4.2)	142	96.67
	High (>4.2)	1	0.67
Free T4 (ng/ml)	Low (<0.8)	15	10.00
	Normal (0.8 to 2)	133	88.67
	High (>2)	2	1.33
TSH ( $\mu$ U/L)	Low (<0.3)	6	4.00
	Normal (0.3 to 4)	107	71.33
	High (>4.00)	37	24.67

4.67 % patients had low free T3 and 0.67 % patients had low free T3. 10.00% patients had low free T4 and 1.33% patients had high free T4. 4.00% patients had low TSH and 24.67% patients had high TSH.

**Table 3. Distribution of thyroid dysfunction**

Classification	No of cases	Percentage
Normal	86	57.33
Hypothyroidism	26	17.33
Hyperthyroidism	5	3.33
Subclinical Hypothyroidism	33	22.00
Total	150	100.0

In present study, 22.00% patients were present with subclinical hypothyroidism, 17.33% patients were present with hypothyroidism and 3.33% patients present with hyperthyroidism.

### Discussion

Diabetes mellitus is one of the most important health problems in the population worldwide and in spite of advances in treatment, a huge number of patients present with complications owing to poor glycaemic control. One of the vital factors that contribute to deprived glycaemic control is thyroid dysfunction, which tends to happen with diabetes mellitus. This study tries to find out the prevalence of thyroid

dysfunction in people with type 2 diabetes mellitus in our region. There is a complex interaction between DM and thyroid disorder. Because of a closely involved relation between insulin and thyroid hormone cellular metabolism, any abnormal levels of either of them may result in a functional derangement of the other.

In present study, 22.00% patients were present with subclinical hypothyroidism, 17.33% patients were present with hypothyroidism and 3.33% patients present with hyperthyroidism.

DM and thyroid diseases are the two endocrine diseases seen commonly in the population. There is an

inter-dependence between insulin and thyroid hormones for normal cellular metabolism so that DM and thyroid diseases can mutually influence the other disease process. [5]

On one hand, thyroid hormones contribute to the regulation of carbohydrate metabolism and pancreatic function, and on the other hand, diabetes affects TFT to variable extents. Studies have found that diabetes and thyroid disorders tend to coexist in the majority of the patients.

Thyroid disorders can have a major impact on glycaemic control and untreated thyroid disorders affects the management of patients with diabetes. Consequently, a systematic approach to thyroid testing in patients with diabetes is recommended earlier which will prevent the occurrence of cardiovascular complications and other diabetic complications such as nephropathy and retinopathy. [6]

Thus, the detection of abnormal hormone levels in addition to other biochemical variables in the early stage of diabetes will help patients to improve their health and reduce the mortality and morbidity.

Vaghasiya K et al [7] was found that 50 diabetic patients (cases) and 50 non-diabetic patients (control) were screened for thyroid dysfunction by thyroid function tests (TFT). Abnormal thyroid function was found in 14 (28%) DM cases and 50 control had normal thyroid function, which is statistically significant. They have found that there is variation in the TSH, T3 and T4 levels, found in diabetics and diabetics with thyroid disorders. Patients with thyroid disorders had a higher level of TSH compared to those without thyroid diseases whereas there was not much of a difference in T4 and T3 levels.

The prevalence of thyroid disorder was 45% in Type 2 diabetics, among them 28% had hypothyroidism, and 17% had hyperthyroidism in a study conducted by Pasupthi et al. [8]

Similar type of study was done by Udiong from Nigeria which showed 46.5% prevalence of thyroid

disorder out of which hypothyroidism was seen in 26.6% and 19.9% had hyperthyroidism. [9]

### Conclusion

Screening for thyroid disease among patients with diabetes mellitus should be routinely performed for early detection and treatment of thyroid dysfunction to delay the complications of diabetes

### References

1. Shekhar Chandra Yadav, Alwin Saldhana, Biswajit Majumdar. Status of Thyroid profile In Type-2 Diabetes Mellitus. Journal of Nobel Medical College. 2012; 2: 64.
2. Banarasidas Bhanot, Park K. Textbook of preventive and social medicine. 19th ed. Jabalpur India; 2007; 327-32.
3. Diabetes in india. Available at [https://en.wikipedia.org/wiki/Diabetes\\_in\\_India](https://en.wikipedia.org/wiki/Diabetes_in_India). Last accessed on 26 Sept. 2020.
4. Vibha Uppal, Chittranjan Vij, Gurdeep Kaur Bedi. Thyroid Disorders in Patients of Type 2 Diabetes Mellitus. Indian J Clin Biochem. 2013 Oct; 28(4): 336–341.
5. Satish R, Mohan V. Diabetes and thyroid diseases. Rev Int J Diabetes Dev Countries. 2003; 23:120-3.
6. Hage M, Zantout MS, Azar ST. Thyroid disorders and diabetes mellitus: review article. J Thyroid Res. 2011; Article ID 439463:7.
7. Vaghasiya K, Patel C, Shahane K, Gamit N, Zala A, Shah A. Estimation of thyroid profile in patients with diabetes mellitus in New Civil Hospital, Surat. Int J Basic Clin Pharmacol. 2015; 4:708-13
8. Pasupathi P. Screening for thyroid dysfunction in the diabetic/ nondiabetic population. Thyroid Sci. 2008;3(8): CSL-6.
9. Udiong CE, Udoh AE, Etukudoh ME. Evaluation of thyroid function in diabetes mellitus in Calabar, Nigeria. Indian J Clin Biochem. 2007; 22(2):74-8.