

## Study of Clinical Profile of Patients of Acute Coronary Syndrome Presenting in Emergency Medicine Department

Pushkar Pradip Shah<sup>1</sup>, Seema Shripad Karhade<sup>2</sup>

<sup>1</sup>MBBS, MD (Medicine), FACEE Associate Professor, Department of Emergency Medicine, Smt Kashibai Navale Medical College and General Hospital, Narhe, Pune. 411041

<sup>2</sup>MBBS, MD (Anaesthesia), FACEE Professor & HOD, Department of Emergency Medicine, Smt Kashibai Navale Medical College and General Hospital, Narhe, Pune. 411041

Received: 25-01-2024 / Revised: 23-02-2024 / Accepted: 25-03-2024

Corresponding Author: Dr. Pushkar Pradip Shah

Conflict of interest: Nil

### Abstract:

**Introduction:** Acute coronary syndrome (ACS) is most common cause of mortality in patients with coronary artery disease. Cardiovascular risk factors for ACS are on the rise in people of Indian origin, and ACS is now the leading cause of death.

**Aims and Objectives:** To study clinical profile, risk factors and complications of patients of acute coronary syndrome (ACS) presenting in Emergency Medicine Department.

**Material and Methods:** An observational study spanning 1 year (1<sup>st</sup> January 2023 to 31<sup>st</sup> December 2023) was conducted in Emergency Medicine Department (EMD) of SKN Medical College & General Hospital, Pune. Data sourced from hospital records with approval.

**Result:** The study analyzed 158 ACS patients. Most of them were males (72%). Most patients were from age group of 40-60 years (45%). STEMI was commonest presentation (68%). Chest pain/discomfort was most common presenting symptom seen in 82% followed by sweating in 61%. Among the risk factors studied, hypertension was commonest (39%) followed by smoking in 30%. Tachyarrhythmias were seen in 11% while cardiogenic shock was seen in 9%.

**Conclusion:** ACS is more common in males with STEMI being commonest presentation and chest pain/discomfort being commonest presenting symptom. Complications can be seen during treatment in EMD and are tachyarrhythmias, cardiogenic shock, and bradyarrhythmias.

**Keywords:** Acute coronary syndrome, ST-elevation myocardial infarction, Non-ST elevation myocardial infarction, Unstable angina, Emergency Medicine Department.

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

### Introduction

Coronary artery disease (CAD) is the leading cause of mortality and morbidity in the world [1] and acute coronary syndrome (ACS) is the most common cause of mortality in patients with CAD. Cardiovascular diseases have become one of the major health problem reaching epidemic proportions. Previous studies have reported that there is a rising incidence of ACS in the young. The currently available evidence, young patients represent 0.4-19% of all ACS cases, depending on the cut-off age used [2-5].

Acute coronary syndrome (ACS) is a constellation of clinical features that results from coronary artery occlusion, which is commonly due to the formation of a thrombus on a ruptured atherosclerotic plaque [6]. The spectrum of ACS includes ST-segment elevation myocardial infarction (STEMI), non-STEMI (NSTEMI), and unstable angina (UA) [7]. Hypertension, diabetes mellitus, dyslipidemia,

obesity, smoking, and a sedentary lifestyle are modifiable risk factors that predispose to ACS. Non-modifiable risk factors include genetic factors, age, gender, race, family history of ischemic heart disease (IHD) and a lower socioeconomic status [8]. Depending on the profile of ACS presentation, percutaneous coronary intervention (PCI) and thrombolytic therapy (fibrinolysis) are the two currently available therapeutic interventions capable of restoring coronary perfusion [9].

Cardiovascular risk factors for ACS are on the rise in people of Indian origin, and ACS is now the leading cause of death [10-14]. The present study was undertaken to study the clinical profile, risk factors and complications of ACS in patients presenting to Emergency Medicine Department (EMD).

## Method

### Research Design

This observational study was conducted in SKNMC & GH, Pune from 1 January 2023 to 31 December 2023 for duration of 1 year. All the patients of ACS coming to our hospital's emergency medicine department during the research period were considered. The necessary data was taken from the medical facility's records after obtaining necessary approvals. ACS was defined as STEMI, NSTEMI, or UA as per established clinical, ECG, and enzymatic definitions for the same. Standard history was taken and data was recorded in case paper. Hypertension was defined as a blood pressure (BP) of >140/90mm Hg. Diabetes mellitus (DM) was defined as a fasting blood sugar (FBS) level of >126mg/dL or postprandial blood sugar (PPBS) level of >200mg/dL or HbA1C > 6.5%. Dyslipidemia was defined as fasting low-density lipoprotein (LDL) levels of >100 mg/ dL or high-density lipoprotein (HDL) levels of < 40 mg/dL or triglyceride (TG) > 150 mg/dL. Patients of STEMI were thrombolysed in EMD ICU as per protocol and then shifted to Cardiac ICU. Prompt medical treatment was started in all cases. Any complications developed during EMD stay was documented.

### Inclusion and Exclusion criteria

#### Inclusion

- Patients of ACS presenting to the emergency medicine department from January 2023 to December 2023.

#### Exclusion


- Patients of ACS not presented in EMD.
- Patients who developed ACS during hospital stay.
- Patients presented with ACS beyond the study period.

#### Statistical analysis


- The study was conducted using SPSS 27 and MS Excel.

#### Result


- A total of 158 patients were enrolled in this study. A number of parameters were observed, including age, gender, risk factors, presenting symptoms, diagnosis and complications in EMD.
- In our study out of 158 patients, 113 (72%) were males and 45 (28%) were females.




According to age group, patients were classified in three age groups - <40 years, 40-60 years, >60 years. 71 (45%) patients were from 40-60 years age group. 67 (42%) were from >60 years age group while 20 (13%) were from <40 years age group




Patients were classified according to type of ACS and STEMI was most common diagnosis, seen in 107 (68%) patients. NSTEMI was seen in 27 (17%) patients and UA was seen in 24 (15%) patients.




Most common presenting symptom was chest pain/discomfort seen in 129 (82%) patients followed by sweating seen in 97 (61%) patients. Breathlessness, epigastric pain, and vomiting were seen in 24 (15%), 17 (11%), and 8 (5%) patients respectively.



Hypertension was most common risk factor present in 62 (39%) patients. Smoking was present in 47 (30%) patients. DM, dyslipidemia, and family history of IHD were the other risk factors present in 40 (25%), 25 (16%), and 8 (5%) patients respectively.



In our study, we also noted complications arising in ACS patients during EMD stay. Most common complication seen was tachyarrhythmia in 17 (11%) patients. Other complications were bradyarrhythmia and cardiogenic shock seen in 10 (6%) and 14 (9%) patients respectively. 4 (2.5%) patients died in EMD during treatment.



**Discussion**

In our study, males (72%) outnumbered females (28%). Similar result was found in study by Naveen Kumar et al in which 83% patients were males and 17% were females [15]. Among ACS patients of all age group, 13% were from < 40 years of age in our study. A study by VK Kadam had similar prevalence with 11% patients from < 40 years age group [16]. In another study by Kesavan et al 9.1% of ACS patients were from <40 years age group [17]. Hence, we can conclude that the percentage of young ACS patients presenting to EMD is increasing in newer studies indicating transition from middle age to young population.

STEMI patients were 68% in our study while NSTEMI and UA were 17% and 15% respectively. A study by VK Kadam had 63% STEMI, 15% NSTEMI and 22 % UA patients [16]. Both the study had STEMI in majority of patients. This was similar to the CREATE registry, which is the largest registry of patients with ACS in India [18]. However, ACCEPT registry and GRACE registry enrolling patients mainly in the western hemisphere found NSTEMI/UA as predominant presentation of the ACS with STEMI coming in second [19,20] Hence, we can conclude that in India, STEMI is main presentation.

Chest pain/discomfort was seen as predominant presenting symptom in 82% patients followed by sweating in 61%. Similar findings were noted in a study by P.Yadav et al with 94% patients having chest pain as predominant symptom followed by sweating in 78% [21]. Other less common symptoms were epigastric pain and vomiting seen in patients with inferior wall myocardial infarction and diabetics.

Hypertension was most common risk factor associated with ACS in our study found in 39% patients followed by smoking in 30%. In a study by VK Kadam, hypertension and smoking were found in 49% and 48% patients respectively [16]. Another study by P.Yadav found hypertension and smoking (tobacco) in 33% and 65% patients respectively [21]. This underlines hypertension and smoking as prominent risk factors for ACS. Other risk factors studied were DM, dyslipidemia, and family history of IHD which were seen in 25%, 16% and 5% patients respectively. Same risk factors were seen in 16%, 14% and 12% patients respectively in a study by P.Yadav highlighting importance of these risk factors [21].

Tachyarrhythmia was most complications arising during EMD stay in our study seen in 11%. Accelerated idioventricular rhythm (AIVR) was most common tachyarrhythmia and was seen during thrombolysis of STEMI. Cardiogenic shock was seen in 9% patients. Bradyarrhythmias were noted in 6% patients. All of these were noted in inferior wall myocardial infarction (IWMI). 2.5 % patients died in EMD during treatment in our study. These patients were presented late and had complications on presentation. In a study by VK Kadam, tachyarrhythmias were found in 48% patients followed by bradyarrhythmias and cardiogenic shock in 30% and 17% patients respectively [16]. In a study by P.Yadav, arrhythmias were seen in 60% while cardiogenic shock was seen in 35% patients [21]. This difference between our study and other studies is due to the fact that we studied complications during the stay of patient in EMD only while in other studies complications during entire hospital stay. This is one of the limitation of our study, other being no medium and long term follow up.

### Conclusion

Despite limitations described above, we can draw some conclusions about the clinical profile in ACS. The number of patients presented in < 40 years of age were though small but not insignificant. Majority of patients were male. STEMI is commonest presentation of ACS. Chest pain/discomfort is commonest presenting symptom however patients can present with other symptoms and in the absence of chest pain. These are mostly IWMI and patients with DM. Hypertension

contribute as major risk factor followed by smoking. Patients can develop complications during treatment in EMD including mortality.

### References

1. Finegold JA, Asaria P, Francis DP. Mortality from ischaemic heart disease by country, region, and age: Statistics from World Health Organisation and United Nations. *Int J Cardiol* 2013;168: 934-45
2. Chen YL, Bhasin A, Youssef AA, Wu CJ, Yang CH, Hsieh YK, et al. Prognostic factors and outcomes in young Chinese patients with acute myocardial infarction undergoing primary coronary angioplasty. *Int Heart J* 2009; 50:1-11.
3. Chan MY, Woo KS, Wong HB, Chia BL, Sutandar A, Tan HC. Antecedent risk factors and their control in young patients with a first myocardial infarction. *Singapore Med J* 2006 ;47:27-30.
4. Morillas P, Bertomeu V, Pabón P, Ancillo P, Bermejo J, Fernández C, et al. Characteristics and outcome of acute myocardial infarction in young patients. The PRIAMHO II study. *Cardiology* 2007;107:217-25.
5. Hoit BD, Gilpin EA, Henning H, Maisel AA, Dittrich H, Carlisle J, et al. Myocardial infarction in young patients: An analysis by age subsets. *Circulation* 1986;74:712-21.
6. ECG diagnosis and classification of acute coronary syndromes. Birnbaum Y, Wilson JM, Fiol M, de Luna AB, Eskola M, Nikus K. *Ann Non-invasive Electrocardiol.* 2014;19:4-14.
7. 2011 ACCF/AHA focused update of the guidelines for the management of patients with unstable angina/ non-st-elevation myocardial infarction (updating the 2007 Guideline). A report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. Wright RS, Anderson JL, Adams CD, et al. *Circulation.* 2011 ;123:2022-2060.
8. Symptoms of men and women presenting with acute coronary syndromes. Arslanian-Engoren C, Patel A, Fang J, Armstrong D, Kline-Rogers E, Duvernoy CS, Eagle KA. *Am J Cardiol .* 2006;98:1177-1181.
9. Acute coronary syndromes. Smith SW, Whitwam W. *Emerg Med Clin North Am.* 2006; 24: 53-89, vi.
10. Reddy KS, Yusuf S. Emerging epidemic of cardiovascular disease in developing countries. *Circulation* 1998;97:596-601.
11. Gupta M, Singh N, Verma S. South Asians and cardiovascular risk: What clinicians should know. *Circulation* 2006;113:e924-9.

12. Jafar TH, Jafary FH, Jessani S, Chaturvedi N. Heart disease epidemic in Pakistan: Women and men at equal risk. *Am Heart J* 2005; 150 :221-6.
13. Mammi MV, Pavithran K, AbduRahiman P, Pisharody R, Sugathan K. Acute myocardial infarction in north Kerala – a 20 year hospital based study. *Indian Heart J* 1991;43:93-6.
14. Yusuf S, Reddy S, Ounpuu S, Anand S. Global burden of cardiovascular diseases: Part I: General considerations, the epidemiologic transition, risk factors, and impact of urbanization. *Circulation* 2001;104:2746-53.
15. Naveen K, Sarit S, et al. Clinical and angiographic profile of patients presenting with first acute myocardial infarction in a tertiary care center in northern India. *Indian heart J* 2008 May-Jun;60(3):210-4
16. VK Kadam. Clinical profile and outcomes of patients presenting with acute coronary syndrome in a tertiary care hospital. *MGM Journal of Medical Sciences* 2019 July – September; 6 (3):113-7
17. Kesavan S, Suryaprakash A, et al. Clinico angiographic profile of young patients presenting with acute coronary syndrome at a tertiary care center in western India. *Indian Heart J* 2010; 62:462-543
18. Xavier D, Pais P, Devereaux PJ, Xie C, Prabhakaran D, Reddy KS, et al. Treatment and outcomes of acute coronary syndromes in India (CREATE): A prospective analysis of registry data. *Lancet* 2008;371:1435-42
19. Piva e Mattos LA, Berwanger O, Santos ES, Reis HJ, Romano ER, Petriz JL, et al. Clinical outcomes at 30 days in the Brazilian Registry of Acute Coronary Syndromes (ACCEPT). *Arq Bras Cardiol* 2013;100:6-13.
20. Fox KA, Eagle KA, Gore JM, Steg PG, Anderson FA; GRACE and GRACE2 Investigators. The Global Registry of Acute Coronary Events, 1999 to 2009—GRACE. *Heart* 2010; 96:1095-101
21. Yadav P, Joseph D, Joshi P, Sakhi P, Jha RK, Gupta J. Clinical profile and risk factors in acute coronary syndrome. *Nat J Community Med* 2010;150:150-2