e-ISSN: 0975-1556, p-ISSN:2820-2643

Available online on www.ijpcr.com

International Journal of Pharmaceutical and Clinical Research 2025; 17(1); 1122-1125

Review Article

Herbal Nano-Emulsion for the treatment of Acne Vulgaris A Review

Prerna Jaiswal[1], Ruchi Khare Shrivastava[2], Aakash Singh Panwar[3]

[1,2,3]Institute of Pharmaceutical Sciences, SAGE University Indore Kailod Kartal Bypass Road Indore (M.P.)

Received: 18-12-2024 / Revised: 03-01-2025 / Accepted: 17-01-2025

Corresponding Author: Aakash Singh Panwar

Conflict of interest: Nil

Abstract:

Acne vulgaris is a widespread dermatological condition characterized by the presence of comedones, papules, pustules, and cysts, affecting a substantial portion of the global population, particularly adolescents and young adults. The pathogenesis of acne involves a multifactorial process, including hyperseborrhea, follicular hyper keratinization, microbial colonization, and inflammation. Conventional treatments, such as antibiotics, retinoids, and hormonal therapies, are often associated with adverse effects, including skin irritation, dryness, and antibiotic resistance. Consequently, there is a growing interest in exploring alternative therapeutic approaches that offer efficacy with minimal side effects.

Herbal medicines have been traditionally used for centuries to treat various skin conditions, including acne. The natural compounds in herbs exhibit a broad spectrum of biological activities, such as anti-inflammatory, antimicrobial, antioxidant, and sebum-regulating properties. However, the clinical application of herbal extracts is often limited by their poor solubility, stability, and bioavailability. Recent advances in nanotechnology have introduced nanoemulsions as a promising delivery system for herbal medicines, addressing these limitations and enhancing their therapeutic potential.

Nanoemulsions are colloidal dispersions of oil and water stabilized by surfactants, with droplet sizes typically ranging from 20 to 200 nm. The unique physicochemical properties of nanoemulsions, such as increased surface area, kinetic stability, and enhanced solubility, make them ideal carriers for delivering lipophilic and hydrophilic herbal compounds. This review aims to comprehensively explore the formulation, characterization, and therapeutic efficacy of herbal nanoemulsions in the treatment of acne vulgaris.

Introduction

Acne vulgaris, commonly referred to as acne, is one of the most prevalent dermatological conditions, affecting millions of people worldwide. It is particularly common during adolescence, but it can persist into adulthood, significantly impacting the quality of life due to its physical and psychological effects. The pathogenesis of acne involves a complex interplay of factors, including increased sebum production, follicular hyper keratinization, proliferation of *Propionibacterium acnes* (recently reclassified as *Cutibacterium acnes*), and inflammation. Traditional treatment options, such as antibiotics, retinoids, and hormonal therapies, although effective, often come with a host of side effects and limitations, such as skin irritation, dryness, and the development of antibiotic resistance.[1,2] These drawbacks necessitate the exploration of alternative therapeutic approaches that are both effective and have a lower risk of adverse effects.

Herbal medicine, with its long-standing history in traditional and folk medicine, offers a promising alternative for acne treatment. Various plant extracts have been recognized for their potent anti-inflammatory, antimicrobial, antioxidant, and sebum-regulating properties. However, the clinical application of herbal extracts is often hampered by their poor solubility, stability, and bioavailability. These limitations can be significantly addressed by advancements in nanotechnology, particularly through the formulation of nanoemulsions. Nanoemulsions, as advanced drug delivery systems, enhance the solubility, stability, and bioavailability of herbal compounds, providing a novel and effective approach for acne treatment.[3]

The Pathogenesis of Acne Vulgaris: Acne vulgaris is a multifactorial inflammatory disease of the pilosebaceous unit, which consists of the hair follicle and its associated sebaceous gland. The pathogenesis of acne involves a complex interplay of factors including genetic, hormonal, microbial, and environmental influences. Here, we delve into the major factors contributing to the development of acne:[4]

Sebum Production: Sebum is an oily substance secreted by the sebaceous glands, primarily composed of triglycerides, wax esters, squalene, and free fatty acids. The production of sebum is regulated by androgens (male hormones), which are present in both males and females. During puberty, increased androgen levels stimulate the sebaceous glands to produce more sebum. This hyperseborrhea, or overproduction of sebum, is a hallmark of acne and provides an ideal environment for the growth of acne-causing bacteria.[6,5]

Follicular Hyperkeratinization

Follicular hyperkeratinization refers to the excessive proliferation and shedding of keratinocytes, the predominant cells in the epidermis, within the hair follicle. This process leads to the formation of a keratin plug, or microcomedone, which obstructs the opening of the hair follicle. Factors contributing to hyperkeratinization include:

- Androgens: These hormones not only increase sebum production but also enhance the proliferation of keratinocytes within the follicle.
- Insulin-like Growth Factor-1 (IGF-1): Elevated levels of IGF-1, often influenced by diet and genetics, can promote keratinocyte proliferation and sebum production.
- Cytokines: Pro-inflammatory cytokines like IL-1 can stimulate keratinocyte proliferation and contribute to the hyperkeratinization process.[7]

Microbial Colonization: The human skin is home to a diverse microbiome, including bacteria, fungi, and viruses. Cutibacterium acnes (formerly Propionibacterium acnes) is a Gram-positive anaerobic bacterium that plays a crucial role in the development of acne. C. acnes reside in the sebaceous glands and contributes to acne through several mechanisms:

- Lipase Activity: C. acnes produce lipases that hydrolyze triglycerides in sebum into free fatty acids. These fatty acids are pro-inflammatory and can disrupt the skin barrier, promoting inflammation.
- Biofilm Formation: The bacteria can form biofilms, which are aggregates of microorganisms embedded in a self-produced matrix. Biofilms protect C. acnes from the immune system and antibiotic treatments, contributing to chronic inflammation and persistence of acne.
- Activation of Toll-like Receptors (TLRs): C. acnes can activate TLRs on keratinocytes and sebocytes, particularly TLR2, leading to the production of pro-inflammatory cytokines such as IL-1, IL-8, and TNF-α.

Inflammation: Inflammation is a central component of acne pathogenesis and is both a cause and consequence of the other pathogenic factors. The inflammatory process in acne involves:

- Cytokine Release: Activation of TLRs by C. acnes and other stimuli results in the release of pro-inflammatory cytokines (IL-1, IL-6, IL-8, and TNF-α) and chemokines that recruit immune cells to the site of infection.
- Inflammatory Cell Infiltration: Neutrophils, macrophages, and lymphocytes infiltrate the follicle, releasing enzymes and reactive oxygen species (ROS) that contribute to tissue damage and the formation of inflammatory lesions such as papules, pustules, and nodules.
- Inflammatory Mediators: Lipid mediators such as leukotrienes and prostaglandins, produced from arachidonic acid, further propagate inflammation.

Hormonal Influences: Hormonal changes, particularly increases in androgens, are pivotal in acne development. Androgens stimulate sebaceous gland activity and sebum production. Conditions such as polycystic ovary syndrome (PCOS) that alter androgen levels can exacerbate acne. Additionally, hormonal fluctuations during the menstrual cycle can influence acne severity in females.

Genetic Factors: Genetic predisposition plays a significant role in acne susceptibility. Family studies have shown that individuals with a family history of acne are more likely to develop the condition. Genetic factors influence sebaceous gland size and activity, hormone levels, and immune responses to microbial colonization and inflammation.

Dietary Factors: Emerging evidence suggests that diet may influence acne. Diets high in glycemic load, dairy products, and certain fats can increase IGF-1 levels, which, in turn, stimulate sebum production and keratinocyte proliferation. Although the exact relationship between diet and acne is complex and not fully understood, dietary modifications can be beneficial for some individuals.[8,9,10]

Environmental Factors: Environmental factors, including pollution, humidity, and exposure to certain chemicals, can exacerbate acne. For instance:

- Pollution: Particulate matter and other pollutants can occlude pores, increase oxidative stress, and promote inflammation.
- Humidity and Sweat: High humidity and excessive sweating can clog pores and create a conducive environment for bacterial growth.
- Cosmetics: Comedogenic cosmetics can block pores and contribute to the formation of comedones.[11]

Stress

Psychological stress has been linked to acne exacerbation. Stress can influence hormone levels (e.g., cortisol) and immune function, leading to increased sebum production and inflammatory responses. Stress management techniques and psychological support can be important adjuncts in acne treatment.[12]

Immune Response: The immune response to C. acnes and other stimuli is a critical determinant of acne severity. An exaggerated immune response can lead to severe inflammation and tissue damage, resulting in more significant lesions and scarring. Understanding individual variations in immune responses can help tailor treatments to reduce inflammation and minimize scarring.

Understanding the pathogenesis of acne vulgaris is essential for developing effective treatments. Acne is a multifactorial disease involving:[13]

- **Increased Sebum Production:** Sebum, an oily substance produced by sebaceous glands, plays a crucial role in acne development. Androgens, particularly during puberty, stimulate the sebaceous glands to produce more sebum. Excessive sebum can create an environment conducive to the growth of *C. acnes*.[14]
- **Follicular Hyperkeratinization:** This process involves the abnormal shedding of keratinocytes within the hair follicle, leading to the formation of a keratin plug. The plug obstructs the sebaceous duct, resulting in the accumulation of sebum and the formation of comedones (whiteheads and blackheads).
- **Microbial Colonization:** *C. acnes* is a Gram-positive anaerobic bacterium that resides in the sebaceous glands. It contributes to acne by metabolizing sebum into pro-inflammatory fatty acids and releasing various pro-inflammatory mediators.
- **Inflammation:** The inflammatory response in acne is mediated by various cytokines and inflammatory cells. *C. acnes* activate toll-like receptors (TLRs) on keratinocytes and sebocytes, leading to the production of pro-inflammatory cytokines such as IL-1, IL-8, and TNF-α.1[5,16,17]

The interplay of these factors leads to the characteristic lesions of acne, including comedones, papules, pustules, and cysts. The chronic nature of acne and its potential for causing permanent scarring necessitate effective treatment strategies.[17]

Conclusion

Acne vulgaris is a multifactorial skin disorder that affects individuals globally, significantly impacting their physical appearance, psychological well-being, and quality of life. Conventional therapies, while effective to some extent, are often associated with limitations such as side effects, antibiotic resistance, and inconsistent efficacy, necessitating the development of safer and more effective alternatives. Herbal bioactives have emerged as promising candidates for acne treatment due to their natural origin, multi-targeted mechanisms of action, and reduced risk of adverse effects. Bioactive compounds such as flavonoids, terpenoids, phenolic acids, and alkaloids exhibit potent antimicrobial, anti-inflammatory, antioxidant, and sebum-regulating properties, addressing the key pathogenic factors of acne vulgaris. The integration of herbal bioactives with advanced drug delivery systems, such as nanoemulsions, has further enhanced their therapeutic potential. Nanoemulsions offer advantages such as improved bioavailability, enhanced skin penetration, controlled release, and protection of bioactives, making them an ideal platform for the delivery of herbal compounds. This innovative approach ensures targeted action, minimizes systemic side effects, and improves patient compliance, thereby overcoming the challenges associated with conventional acne treatments. In conclusion, the combination of herbal medicine and nanoemulsion technology represents a significant advancement in acne therapy. This approach holds the potential to provide a safe, effective, and patient-friendly solution for the management of acne vulgaris. However, further research, including rigorous clinical studies, is essential to validate the efficacy, safety, and long-term benefits of herbal nanoemulsions. With continued scientific exploration and optimization, herbal nanoemulsions may become a mainstream therapeutic option, offering a natural and innovative solution for individuals suffering from acne vulgaris.

Reference

1. Shakeel F, Baboota S, Ahuja A, Ali J, Shafiq S, Khan S. Nanoemulsions as vehicles for transdermal delivery of aceclofenac. AAPS PharmSciTech. 2007;8(4)
2. Kumar R, Philip A. Modified transdermal technologies: breaking the barriers of drug permeation via the skin. Trop J Pharm Res. 2007;6(1):633-644.
3. Lawrence MJ, Rees GD. Microemulsion-based media as novel drug delivery systems. Adv Drug Deliv Rev. 2000;45(1):89-121.
4. Sharma N, Mishra S, Sharma S, Deshpande RD, Sharma RK. Preparation and optimization of nanoemulsions for targeting drug delivery. Int J Drug Dev Res. 2013;5(4):37-48.
5. Tadros T, Izquierdo P, Esquena J, Solans C. Formation and stability of nano-emulsions. Adv Colloid Interface Sci. 2004;108-109:303-318.
6. Solans C, Izquierdo P, Nolla J, Azemar N, Garcia-Celma MJ. Nano-emulsions. Curr Opin Colloid Interface Sci. 2005;10(3-4):102-110.
7. Gupta A, Eral HB, Hatton TA, Doyle PS. Nanoemulsions: formation, properties and applications. Soft Matter. 2016;12(11):2826-2841.

8. Sarker DK. Engineering of nanoemulsions for drug delivery. Curr Drug Deliv. 2005;2(4):297-310.
9. McClements DJ. Nanoemulsions versus microemulsions: terminology, differences, and similarities. Soft Matter. 2012;8(6):1719-1729.
10. Forgiarini A, Esquena J, González C, Solans C. Studies of the relation between phase behavior and emulsification methods with nano-emulsion formation. Prog Colloid Polym Sci. 2001;118:184-189.
11. Rao J, McClements DJ. Formation of flavor oil microemulsions, nanoemulsions and emulsions: influence of composition and preparation method. J Agric Food Chem. 2011; 59(10): 5026-5035.
12. Panwar A.S., Rathore P., Ghodke A. Physiology of Aging and Mechanism of Herbal Anti-Aging Drugs: A Review from a Pharmaceutics Perspective Journal of Medical Case Reports and Reviews 2024 7 (10) 1418-1423
13. Panwar A. S., Rathore P. Impact of formulation excipients on skin barrier functions: A review International Journal of Pharmaceutical Chemistry and Analysis 11 (1) 41-44
14. Anton N, Benoit JP, Saulnier P. Design and production of nanoparticles formulated from nanoemulsion templates—a review. J Control Release. 2008;128(3):185-199.
15. Salvia-Trujillo L, Rojas-Graü MA, Soliva-Fortuny R, Martín-Belloso O. Physicochemical characterization of lemongrass essential oil-alginate nanoemulsions: effect of ultrasound processing parameters. Food Bioprocess Technol. 2013;6(9):2439-2446.
16. Shakeel F, Ramadan W, Shafiq S, Ahmed MA. Nanoemulsions as potential vehicles for transdermal and dermal delivery of hydrophobic compounds: an overview. Expert Opin Drug Deliv. 2011;8(7):957-971.
17. Solans C, Kunieda H. Industrial applications of nanoemulsions. Curr Opin Colloid Interface Sci. 2013;18(3):235-244.