

The Effects Of Washing Practices And Storage On Antioxidant Activity Of Some Selected Fruits

Rastkari Noushin^{1*}, Farajvand Nasrin², Alimohammadi Mahmood², Yunesian Masud^{1,2}, Samadi Nasrin³

¹Center for Air Pollution Research (CAPR), Institute for Environmental Research (IER), Tehran University of Medical Sciences (TUMS), Tehran, Iran

²Department of Environmental Health Engineering, School of Public Health, Tehran University of Medical Sciences (TUMS), Tehran, Iran.

³Department of Drug and Food Control, Faculty of Pharmacy and Pharmaceutical Quality Assurance Research Center, Tehran University of Medical Sciences (TUMS), Tehran, Iran

Available Online: 1st January, 2015

ABSTRACT

Effects of washing practices using a commercial detergent or disinfectants alone or in combination with each other on the radical-scavenging activity and reducing power of the fresh fruits were examined. The fresh fruits were subjected to a triple wash treatment of washing in tap water for mud removal, washing in water containing a detergent (dishwashing liquid) or disinfectant individually, and rinsing in tap water. The results show that there were significant variations in antioxidant activity across fresh fruits (ranging from 15.02 to 57.15 mmol/L Trolox equivalents per kg of fresh weight). The antioxidant capacities of the all samples were soaked in Kanz Disinfecting Liquid showed lowest antioxidant activity ($P < 0.05$). Pretreatment with detergent before washing with calcium hypochlorite or benzalkonium chloride showed significant reductions of antioxidant activity (ranging from 8 to 27%, $P < 0.05$). Levels of total antioxidant activity showed different changes throughout the storage period at 4 °C depend on storage duration.

Key words: Total antioxidant activity, washing practices, Storage, Disinfectants, Fresh fruit, 2, 2-diphenyl-1-picrylhydrazyl (DPPH), Ferric reducing antioxidant power (FRAP)

INTRODUCTION

Various epidemiological studies have reported that diets rich in fruits, vegetables and grains are associated with a lower risk of several degenerative diseases, such as cancers and cardiovascular diseases¹. Nutritional recommendations to reduce the risk of coronary heart disease have focused largely on the intake of nutrients that affect the risk factors, including plasma lipid and lipoprotein levels, blood pressure, and body weight, etc. Over the past two decades, considerable evidence has been gathered in support of the hypothesis that the free radicals and free-radical-mediated oxidative processes and specific end products play a key role in atherogenesis². A practical way of fighting against degenerative diseases may be to improve body antioxidant status. Antioxidants constitute a diverse group of compounds with different properties. They operate by inhibiting oxidant formation, intercepting oxidants once they have formed, and repairing oxidant-induced injury. Although there is an effective antioxidant defense mechanism to protect the body against oxidant attacks, sometimes it cannot cope with oxidant load in the body, and additional dietary antioxidants are needed³⁻⁵. Fresh vegetables and fruits are excellent sources of antioxidant components. Fruits are an important part of our diet and as a source of energy, fiber, vitamins, minerals and antioxidants⁶. Washing of fruits and vegetables is a

common method to eliminate soil, pesticide residues, debris and to reduce microbial load⁷. Sanitization treatments can play an important role in reducing the indigenous microflora and pathogens on fresh fruits and vegetables. Application of a detergent before disinfection may help remove microorganisms from the surface of fresh produce⁸⁻⁹. Most of the investigations concerning the efficacy of disinfectants for reduction of pathogenic bacteria have been conducted on inoculated fresh fruits such as cantaloupe, berries, and apples¹⁰⁻¹³ and vegetables such as lettuce and cabbage¹⁴⁻¹⁷. But the efficacy of these treatments on the antioxidant activity of the fresh fruits has not been clearly defined. Therefore, this study was carried out to determine the effects of domestic washing practices with detergents and fresh produce disinfectants or their combination, on antioxidant activity of the fresh fruits.

MATERIALS AND METHODS

Fruits: Seven types of fresh fruits including strawberry (*Fragaria ananassa L.*), cherry (*Prunus avium L.*), grape (*Vitis vinifera L.*, White table grapes), peach (*Prunus Persica L.*), apricot (*Prunus armeniaca L.*), pear (*Pyrus communis L.*) and apple (*Malus domestica L.*) were purchased during harvest period from the main field of fruits and vegetables in Tehran, Iran, before passing 24 h from harvesting, packed in plastic bags, and transported to

the laboratory within 1 h. Approximately 8 kg of each mature fruit was selected, from which damaged fruits were discarded. The fruits were treated on the day of purchase and used directly for analyses at time zero. Treated fruits were stored at 4 °C and analyzed after several storage times (3 and 5 days).

Preparation and decontamination of fruits: For each replication of disinfection or washing procedure, about 2 kg of each type of fruits was added to 6 L of tap water with gentle stirring for about 20 s to ensure complete immersion and mud removal. An adequate sample of fruit was separated immediately for antioxidant activity determination before any treatment. Fruits were removed from water, and subjected to the one of the procedures presented in the treatment step in Figure 1 (step 2, procedures 2.1 to 2.4). Each of the treatment procedures was replicated three times.

Preparation of disinfectant solutions: The following disinfectants with their respective concentrations and

chloride (Hygen, Mosmer Co., Tehran, Iran) at 92 ppm for 15 min, calcium hypochlorite (perchlorine, commercial grade) at 300 ppm for 15 min and Kanz Disinfecting Liquid for vegetables & fruits (which is sold in Iran local markets and its active ingredients are: benzalkonium chloride and cocoamidopropyl betaine) at 110 ppm for 20 min. The concentration of benzalkonium chloride was determined following the method proposed by the U.S. Pharmacopeia¹⁸. Calcium hypochlorite concentration was determined by titration with sodium thiosulfate¹⁹. A dishwashing liquid (Golrang Group, Tehran, Iran) containing alkyl benzene sulfonate, sodium lauryl ether sulfate, isothiazolones, and coconate fatty acid diethanol amide was used as the detergent. Kanz Disinfecting Liquid was used alone not in combination with detergent.

Sample preparation: Samples were taken after each processing step and used directly for analyses at time zero. Treated fruits were stored at 4 °C and analyzed after 3 and 5 days storage. For the analysis, the extract was




Fig 1. Fresh fruit decontamination protocols.

exposure times (according to the manufacturer's recommendations) were used in the study: benzalkonium

Table 1. Antioxidant content expressed as mmol L⁻¹ Trolox equivalents per kg of fresh weight of antioxidant capacity (DPPH) (mean ±SD) of fresh and treated fruits.

Fruit ^a	Storage time (days) ^b	Treatments ^{c,d}							
		1	2	3	4	5	6	7	8
Strawberry	1	51.12±2.17	50.67±1.13	49.61±1.51	49.73±1.19	40.52±1.07	48.99±2.07	41.04±1.18	37.96±1.41
	3	42.73±1.77	41.97±1.10	42.10±1.41	44.08±1.54	35.20±1.19	44.63±2.26	36.05±1.29	33.53±1.92
	5	42.08±1.52	42.05±1.81	41.94±1.66	43.89±1.21	34.71±0.87	45.08±1.19	35.60±1.09	32.92±1.88
Cherry	1	47.75±1.25	48.06±1.37	46.85±1.86	46.51±1.02	40.10±1.57	47.18±1.11	39.79±2.02	36.10±1.21
	3	42.75±1.07	43.06±1.21	39.25±1.02	38.51±1.79	35.10±1.47	40.18±1.23	36.79±1.12	31.10±1.36
	5	38.83±1.72	38.96±1.51	35.68±0.87	35.18±1.47	30.16±1.17	35.01±0.97	29.59±1.17	27.39±1.27
Peach	1	44.97±2.18	44.81±1.88	45.63±1.95	45.70±1.24	38.88±1.33	44.42±2.41	38.74±1.67	37.70±1.77
	3	39.89±1.80	38.85±1.45	40.30±1.34	40.60±1.18	34.07±0.97	39.74±0.85	33.84±1.33	28.78±1.68
	5	35.03±1.21	35.92±1.35	36.45±1.97	35.89±1.18	30.86±1.88	36.15±1.46	30.06±1.28	24.81±2.21
Pear	1	39.07±1.11	39.23±1.24	38.07±1.31	38.00±1.87	36.21±1.61	37.97±2.09	35.88±1.74	32.84±1.66
	3	38.66±1.47	39.15±1.77	38.45±1.14	37.61±1.57	34.57±2.11	36.88±2.32	35.01±1.49	31.91±1.96
	5	38.98±2.55	38.92±0.96	37.92±1.47	37.91±1.22	34.05±1.18	36.21±1.71	34.17±1.55	31.42±1.41
Apple	1	37.83±1.41	38.40±1.28	38.99±2.57	37.57±1.67	35.93±1.51	38.33±1.63	34.96±1.74	32.68±1.15
	3	36.98±1.49	37.60±1.71	38.33±1.23	36.36±1.49	35.18±1.35	37.50±1.61	34.01±0.82	31.77±1.43
	5	36.13±1.15	36.97±1.94	37.55±1.81	36.96±1.41	34.88±1.58	37.14±2.05	33.42±1.17	31.04±1.51
Grape	1	16.39±1.37	16.32±1.22	16.20±1.02	16.97±0.96	12.02±1.33	15.66±2.07	13.08±1.32	10.81±1.45
	3	15.86±1.23	14.98±1.56	16.83±0.93	16.23±1.41	11.59±1.26	14.96±1.14	12.65±1.57	9.76±1.17
	5	15.78±1.11	15.32±1.49	15.68±1.56	15.96±0.85	12.14±1.34	14.22±1.28	13.16±1.19	9.97±1.05
Apricot	1	15.02±1.49	14.96±1.13	15.09±0.77	15.19±1.11	12.55±1.32	15.12±2.37	12.75±2.02	8.85±1.95
	3	11.60±1.76	10.71±1.55	12.29±1.37	12.01±1.19	8.60±2.15	11.36±0.89	9.38±1.19	5.46±1.62
	5	8.48±1.44	8.85±1.32	9.02±2.17	8.35±2.03	5.62±0.84	9.41±1.11	6.61±1.21	3.47±1.04

^a Results showed that there were significant variation in antioxidant activity across the selected fresh fruits (P<0.05).

^b Homogeneous subgroups are discussed in the results and discussion section.

^c Different treatments: 1) only mud removal, 2) Soak fruits in 6 lit tap water for 10 min, 3) Soak fruits in detergent solution (2 mL in 6 lit tap water) for 10 min, 4) Soak fruits in 6 lit calcium hypochlorite solution (300 ppm) for 15 min, 5) Soak fruits in 6 lit benzalkonium chloride solution (92 ppm) for 15 min, 6) First, soak fruits in detergent solution (2 mL in 6 lit tap water) for 10 min and then in 6 lit calcium hypochlorite solution (300 ppm) for 15 min, 7) First, soak fruits in detergent solution (2 mL in 6 lit tap water) for 10 min and then in 6 lit calcium hypochlorite solution (300 ppm) for 15 min, 8) Soak fruits in 6 lit Kanz disinfecting liquid solution (110 ppm) for 20 min.

^d No significant differences in total antioxidant activity were found on selected fruits after decontamination with treatments 1,2,3,4 and 6 (P > 0.05). Decontamination with treatments 5,7 and 8 showed significant reductions of antioxidant activity (P<0.05).

prepared by a method described by Scalzo et al.²⁰ and Gao and Mazza²¹. Twenty g of the homogenized sample was extracted with 60 ml of ethanol (Merck no.1009832500) and was placed for 2 h at room temperature in darkness. The mixture was centrifuged, and the clear phase was separated and filled to 100 ml with the extraction solvent. This extract was used for the determination of the antioxidant capacity.

2, 2-diphenyl-1-picrylhydrazyl (DPPH) assay: A slightly modified method of Alamanni and Cossu²² strawberry was used in which a stable radical (DPPH) is reduced by the antioxidants of the sample, which leads to a decolonization. Ten ml of a 0.1 mmol/L DPPH (Fluka Chemical Co.) solution in ethanol were mixed with 0.1 ml sample extract. After 30 min, the absorbance at 517 nm

was measured with Lambda 25 spectrophotometer (Perkin Elmer co.). Trolox (6-hydroxy-2, 5, 7, 8-tetramethylchroman-2-carboxylic acid) was used as reference compound, and the antioxidant capacity is expressed in mmol/L trolox equivalents per kg of fresh weight (mmol/L T.E /kg fw).

Ferric reducing antioxidant power (FRAP) assay: The method described by Benzie and Strain²³ modified by Guo et al.²⁴, which is based on the reduction of a ferric (III) complex to its colored ferrous (II) form in the presence of antioxidants. The FRAP reagent was prepared freshly before measuring and contained 5 ml of a 10 mmol/L TPTZ (2,4,6-tri(2-pyridyl)-s-triazine) solution (Merck) in 40 mmol/L hydrochloric acid, 5 ml of a 20 mmol/L ferric (III) chloride solution, and 50 ml of 0.3

Table 2. Antioxidant content expressed as mmol L⁻¹ Trolox equivalents per kg of fresh weight of antioxidant capacity (FRAP) (mean±SD) of fresh and treated fruits.

Fruit ^a	Storage time (days) ^b	Treatments ^{c,d}							
		1	2	3	4	5	6	7	8
Strawberry	1	57.15±1.47	56.77±1.76	56.21±1.37	57.10±2.05	46.72±1.15	55.99±1.93	46.14±1.41	40.86±1.58
	3	48.73±1.31	49.07±1.30	48.50±1.11	49.08±1.29	41.20±2.21	50.03±0.94	40.05±1.51	37.63±1.48
	5	48.48±1.74	48.75±1.58	48.94±1.61	48.89±1.75	40.71±1.23	49.68±1.14	39.60±1.47	38.12±1.56
Cherry	1	52.75±1.29	52.06±1.37	51.95±1.86	50.95±1.02	48.10±1.57	51.28±1.11	47.79±2.02	40.11±1.21
	3	45.16±1.28	44.15±1.08	43.16±0.85	42.34±1.41	36.16±1.33	44.69±1.57	35.46±1.91	32.77±1.67
	5	40.83±1.55	39.56±2.15	38.68±1.43	38.18±1.29	31.16±1.34	37.81±1.65	30.09±1.11	28.39±0.95
Peach	1	49.27±1.52	48.81±1.24	48.63±1.16	47.98±1.12	39.48±1.03	47.52±1.25	38.54±1.13	36.71±1.29
	3	42.29±1.46	41.65±0.89	40.90±2.25	40.60±2.01	35.77±1.43	38.74±1.23	34.34±1.82	30.74±1.36
	5	37.23±1.02	37.52±1.74	36.95±1.91	35.89±1.35	31.86±1.16	35.05±1.44	30.06±1.49	26.88±2.08
Pear	1	43.67±1.44	43.91±1.21	44.07±1.50	43.25±1.66	41.31±0.94	42.97±1.18	40.88±1.41	33.44±1.50
	3	42.96±1.27	42.85±1.75	43.45±1.45	42.61±1.07	39.77±2.50	43.08±1.58	40.01±1.22	34.01±1.73
	5	43.28±1.36	42.72±1.04	42.92±2.11	42.12±1.75	40.65±1.48	42.64±1.59	39.87±1.12	33.72±1.20
Apple	1	41.83±1.49	41.40±1.06	40.99±1.18	40.27±1.36	37.93±1.65	39.93±1.18	36.96±2.13	32.58±1.35
	3	41.78±1.39	41.60±1.72	40.33±1.52	41.06±1.64	37.98±1.48	39.50±1.67	36.01±1.68	31.77±1.54
	5	40.43±2.25	41.67±1.61	39.59±0.65	39.96±1.26	38.08±1.14	38.94±1.05	36.42±1.25	32.14±1.14
Grape	1	23.19±1.61	22.52±1.26	22.08±1.82	21.97±1.81	17.32±1.70	21.86±1.06	17.58±1.76	11.81±1.32
	3	23.06±1.98	23.13±1.99	21.53±1.70	21.23±1.31	17.59±1.85	21.46±0.87	17.35±1.39	10.96±1.33
	5	22.78±1.68	22.32±1.11	21.18±1.78	20.95±0.99	17.46±1.77	19.82±1.15	16.96±1.25	11.17±1.01
Apricot	1	18.80±1.75	18.66±1.27	17.82±1.91	17.19±2.25	15.15±1.33	17.52±1.41	14.75±1.19	11.85±1.22
	3	14.32±1.64	14.01±1.76	12.89±1.28	12.31±1.88	10.63±1.91	13.06±2.35	10.88±1.97	8.66±1.77
	5	10.52±1.32	10.15±1.49	9.52±1.54	10.35±1.62	8.72±1.24	9.81±1.05	9.01±0.97	5.77±1.41

^a Results showed that there were significant variation in antioxidant activity across the selected fresh fruits (P<0.05).

^b Homogeneous subgroups are discussed in the results and discussion section.

^c Different treatments: 1) only mud removal, 2) Soak fruits in 6 lit tap water for 10 min, 3) Soak fruits in detergent solution (2 mL in 6 lit tap water) for 10 min, 4) Soak fruits in 6 lit calcium hypochlorite solution (300 ppm) for 15 min, 5) Soak fruits in 6 lit benzalkonium chloride solution (92 ppm) for 15 min, 6) First, soak fruits in detergent solution (2 mL in 6 lit tap water) for 10 min and then in 6 lit calcium hypochlorite solution (300 ppm) for 15 min, 7) First, soak fruits in detergent solution (2 mL in 6 lit tap water) for 10 min and then in 6 lit calcium hypochlorite solution (300 ppm) for 15 min, 8) Soak fruits in 6 lit Kanz disinfecting liquid solution (110 ppm) for 20 min.

^d No significant differences in total antioxidant activity were found on selected fruits after decontamination with treatments 1,2,3,4 and 6 (P > 0.05). Decontamination with treatments 5,7 and 8 showed significant reductions of antioxidant activity (P<0.05).

mol/L acetate buffer (pH 3.6). The extract (0.1 ml) was mixed with 0.3 ml of water and 3 ml of reagent. After 8 min, the absorbance was measured at 593 nm. Trolox was used as the reference compound, and the antioxidant capacity is expressed in mmol/L T.E /kg fw.

Statistical analysis: All statistical analyses were carried out by SPSS for Windows (version 11.5). The data determined were expressed as the mean of three replicate determinations and presented as means ± SD. Data from each treatment or control group were analyzed for differences by using repeated-measures analysis. Tukey's all-pairwise-comparison test was used to identify differences between different groups. P<0.05 was considered statistically significant. Pearson's bivariate

correlation coefficients were calculated to compare the DPPH and FRAP assay. Fresh fruits data is expressed on fresh weight basis.

RESULTS

Antioxidant capacity and reducing power of fresh fruits extracts: The DPPH scavenging activity of fresh fruits ranged from 15.02 to 51.12 mmol/L Trolox equivalents per kg of fresh weight (mmol/L T.E /kg fw), with high activity being found in strawberry, cherry, peach, pear and apple, while low activity was observed in grape and apricot (Table 1). Similarly, the FRAP activity of fresh fruits showed a wide range from 18.8 to 57.15 mmol/L T.E /kg fw, the highest activity being in strawberry (57.15 mmol/L

T.E /kg fw) and the lowest in apricot (18.8 mmol/L T.E /kg fw) (Table 2). A significant correlation was found between DPPH and FRAP assays ($r = 0.84$, $r^2 = 70.56\%$) in fresh fruits. The good correlation between the results from DPPH and FRAP assays in dry fruits has been previously reported²⁵.

Effects of washing treatment on total antioxidant activity of fruits: The antioxidant capacities of untreated samples were measured with the DPPH and FRAP assay and the antioxidant activity of fresh fruits showed a wide range from 15.02 to 51.12 and 18.8 to 57.15 mmol/L T.E /kg fw, respectively (Table 1 and 2). The differences between the means for fruits washed in tap water, detergent, calcium hypochlorite or benzalkonium chloride solution alone were not significant. The samples which were washed with the combination of detergent and one kind of fresh produce disinfectants in two sequential steps showed significant reductions of antioxidant activity ($P < 0.05$). The antioxidant capacities of the all samples were soaked in Kanz Disinfecting Liquid measured with the DPPH and FRAP assay, showed lowest antioxidant activity ranging from 8.85 to 37.96 and 11.85 to 40.86 mmol/L T.E /kg fw, respectively (Table 1 and 2).

Effects of storage time on total antioxidant activity of fruits: In the present study, a significant ($P < 0.05$) decrease in total antioxidant activity were observed for cherry, apricot and peach samples from days 0 to 3 and from days 3 to 5 (Table 1 and 2).

DISCUSSION

A variety of methods specific to their chemical properties have been used to quantify antioxidant activity in plant foods²⁶. Since DPPH and FRAP radical scavenging assays are widely used due to their simplicity, stability, accuracy and reproducibility^{25, 27} we conducted these methods to quantify antioxidant activity in the present study.

The selected fruits were subjected to a triple wash treatment: washing in tap water for mud removal, washing in water containing a detergent or disinfectant and rinsing in tap water. The differences between the means for fruits washed in tap water, detergent, calcium hypochlorite or benzalkonium chloride solution alone were not significant. In previous study⁷, no significant effects of washing treatments (Tap rinsed, water dipped and chlorine dipped) on levels of total antioxidant activity were observed. The samples which were washed with the combination of detergent and one kind of fresh produce disinfectants in two sequential steps showed significant reductions of antioxidant activity. These reductions were higher in fruits with slim skin and soft flesh such as grape (27%), strawberry (21%), cherry, apricot (17%) and peach (16%). Apple and pear showed less reduction (7 and 8 %, respectively). This observation could possibly be explained by the fact that these fruits have thick skin and fibrous structure, which can reduce the penetration of detergent and disinfectants into the soft tissues.

The antioxidant capacities of the all samples were soaked in Kanz Disinfecting Liquid showed lowest antioxidant activity. The probable cause could be that the using of detergent and disinfectants in one step and at the same time

has synergetic effect on the reductions of antioxidant activity. Samadi et al. have reported that application of a detergent at the permitted level before disinfection did not improve antimicrobial activity of the disinfectant significantly²⁸. However, the detergent could have other useful biological effects, such as detachment of parasites from vegetables. In summary, we can conclude that the necessity of application of a detergent at the permitted level before disinfection need further studies.

Levels of total antioxidant activity showed different changes throughout the storage period depend on the type of fruits (ranging from 0 to 35%). No significant effects of storage time were observed on activity of apple, grape and pear samples. Similar observations were previously done on some fruits²⁹⁻³⁰ or vegetables³¹ stored at room temperature or in the refrigerator. The results of these studies indicated that in some fruits and vegetables the storage did not affect negatively the antioxidant capacity. Sacchetti et al. have also shown that the antioxidant capacity of apple derivatives were not negatively affected by storage, and in some cases, an increase of the antioxidant capacity was observed during storage³². Kevers and et al. have observed that antioxidant capacity decreased during storage in apricot (25%) and decreased by >50% in spinach, banana, broccoli, and leek⁶. In apricot, Bartolini et al also observed a decrease of the antioxidant capacity during storage at low temperature³³. Storage also affected the antioxidant activity of strawberry from days 0 to 3, and the decrease observed from days 3 to 5 was not significant ($P > 0.05$). These results also were similar to the ones asserted in the literature³⁴.

CONCLUSION

Antioxidative capacity is an important fruit quality parameter. The results of this study indicate that each type of fruits had a different antioxidant activity, which may contribute by different antioxidant components, such as α -tocopherol, β -carotene, vitamin C, selenium or phenolic compounds. Washing and sanitizing treatment can play an important role in reducing microbial populations on fresh fruits and vegetables, but the efficacy of these treatments on the antioxidant activity of the fresh fruits has not been clearly defined. The results of this study indicated that washing with chlorine did not negatively affect the total antioxidant activity, but in combination with detergents (in two sequential steps or at the same time) has negative synergistic effects on total antioxidant activity. Application of a detergent at the permitted level before disinfection may help remove microorganisms from the surface of fruits and vegetables but further studies are needed to determine its efficacy. Levels of total antioxidant activity showed different changes throughout the storage period depend on the type of fruits. However this study has some limitations. the current study only examined total antioxidant activity among different fruits without qualification and quantification of antioxidant substances separately, therefore it was impossible to draw a cause and effect in the observed reduction. Further study of this possible association is warranted. If the results of this research would be confirmed in future prospective

studies, May be it is nessecary in order to maximize the contents of antioxidant active substances in fruits, the number of processing steps are minimized as far as possible and long holding times should be avoided.

ACKNOWLEDGMENTS

This research has been supported by Tehran University of Medical Sciences (TUMS) and Health Services grant (project no. 91-01-63-17301). Hereby, the cooperation of the University, School of public health and also the Institute for Environmental Research (IER) is highly appreciated.

REFERENCES

1. Amin I, Norazaidah Y, Hainida KIE. Antioxidant activity and phenolic content of raw and blanched Amaranthus species. *Food Chemistry* 2006; 94(1): 47-52.
2. Lee J-Y, Hwang W-I, Lim S-T. Antioxidant and anticancer activities of organic extracts from Platycodon grandiflorum A. De Candolle roots. *Journal of Ethnopharmacology* 2004; 93(2-3): 409-415.
3. Erguder B, Avci A, Devrim E, Durak I. Effects of cooking techniques on antioxidant enzyme activities of some fruits and vegetables. *Turkish Journal of Medical Sciences* 2007; 37(3): 151.
4. Locatelli M, Travaglia F, Coisson JD, Martelli A, Stévigny C, Arlorio M. Total antioxidant activity of hazelnut skin (Nocciola Piemonte PGI): Impact of different roasting conditions. *Food Chemistry* 2010; 119(4): 1647-1655.
5. Turkoglu A, Duru ME, Mercan N, Kivrak I, Gezer K. Antioxidant and antimicrobial activities of *Laetiporus sulphureus* (Bull.) Murrill. *Food Chemistry* 2007; 101(1): 267-273.
6. Kevers C, Falkowski M, Tabart J, Defraigne JO, Dommès J, Pincemail J. Evolution of antioxidant capacity during storage of selected fruits and vegetables. *Journal of Agricultural and Food Chemistry* 2007; 55(21): 8596-8603.
7. Kenny O, Obeirne D. The effects of washing treatment on antioxidant retention in ready-to-use iceberg lettuce. *International Journal of Food Science & Technology* 2009; 44(6): 1146-1156.
8. Olmez H, Kretzschmar U. Potential alternative disinfection methods for organic fresh-cut industry for minimizing water consumption and environmental impact. *LWT-Food Science and Technology* 2009; 42(3): 686-693.
9. Sapers GM, Miller RL, V. Pilizota, Matrazzo AM. Antimicrobial treatments for minimally processed cantaloupe melon. *Journal of Food Science* 2006; 66(2): 345-349.
10. Bastos MSR, N. F. F. Soares, N. J. Andrade, Arruda AC, Alves aRE. The effect of the association of sanitizers and surfactant in the microbiota of the cantaloupe (*Cucumis melo* L.) melon surface. *Food Control* 2005; 16: 369-373.
11. Kaye VS, K. H. McWatters, Beuchat aLR. Efficacy of gaseous chlorine dioxide as a sanitizer for killing Salmonella, yeasts, and molds on blueberries, strawberries, and raspberries. *Journal Food Protection*® 2005; 68: 1165-1175.
12. Liao CH, Sapers GM. Attachment and growth of Salmonella Chester on apple fruits and in vivo response of attached bacteria to sanitizer treatments. *Journal of Food Protection*® 2000; 63(7): 876-883.
13. Wisniewsky MA, B. A. Glatz, M. L. Gleason, Reitmeier aCA. Reduction of Escherichia coli O157: H7 counts on whole fresh apples by treatment with sanitizers. *Journal of Food Protection*® 2000; 63(6): 703-708.
14. Hellstrom S, R. Kervinen, M. Lyly, R. Ahvenainen-Rantala, H. a, Korkeala. Efficacy of disinfectants to reduce *Listeria monocytogenes* on pre-cut iceberg lettuce. *Journal of Food Protection*® 2006; 69(7): 1565-1570.
15. Hilgren JD, Salverda JA. Antimicrobial Efficacy of a Peroxyacetic/Octanoic Acid Mixture in Fresh-Cut-Vegetable Process Waters. *Journal of Food Science* 2000; 65(8): 1376-1379.
16. Nascimento MS, N. Silva, M. P. L. M. Catanozi, Silva KC. Effects of different disinfection treatments on the natural microbiota of lettuce. *Journal of Food Protection*® 2003; 66(9): 1697-1700.
17. Vandekinderen I, J. Van Camp, B. De Meulenaer, K. Veramme, Q. Denon, Ragaert P. The effect of the decontamination process on the microbial and nutritional quality of fresh-cut vegetables. *Acta Hort* 2007; 746(173-179).
18. Rockville M. U.S. Pharmacopeia. In; 2006: 3281-3282.
19. Velazquez LC, N. B. Barbini, M. E. Escudero, C. L. Estrada, Guzman AMSd. Evaluation of chlorine, benzalkonium chloride and lactic acid as sanitizers for reducing *Escherichia coli* O157: H7 and *Yersinia enterocolitica* on fresh vegetables. *Food Control* 2009; 20(3): 262-268.
20. Scalzo J, Politi A, Pellegrini N, Mezzetti B, Battino M. Plant genotype affects total antioxidant capacity and phenolic contents in fruit. *Nutrition* 2005; 21(2): 207-213.
21. Gao L, Mazza G. Characterization, quantitation, and distribution of anthocyanins and colorless phenolics in sweet cherries. *Journal of Agricultural and Food Chemistry* 1995; 43(2): 343-346.
22. Alamanni M, Cossu M. Radical scavenging activity and antioxidant activity of liquors of myrtle (*Myrtus communis* L.) berries and leaves. *Italian journal of food science* 2004; 16(2): 197-208.
23. Benzie IF, Strain J. The ferric reducing ability of plasma (FRAP) as a measure of "antioxidant power": the FRAP assay. *Analytical biochemistry* 1996; 239(1): 70-76.
24. Guo C, Yang J, Wei J, Li Y, Xu J, Jiang Y. Antioxidant activities of peel, pulp and seed fractions of common fruits as determined by FRAP assay. *Nutrition Research* 2003; 23(12): 1719-1726.

25. Vijaya Kumar Reddy C, Sreeramulu D, Raghunath M. Antioxidant activity of fresh and dry fruits commonly consumed in India. *Food Research International* 2010; 43(1): 285-288.
26. Stratil P, Klejdus B, Kuban V. Determination of total content of phenolic compounds and their antioxidant activity in vegetables evaluation of spectrophotometric methods. *Journal of Agricultural and Food Chemistry* 2006; 54(3): 607-616.
27. Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C. Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radical Biology and Medicine* 1999; 26(9): 1231-1237.
28. Samadi N, Abadian N, Bakhtiari D, Fazeli MR, Jamalifar H. Efficacy of detergents and fresh produce disinfectants against microorganisms associated with mixed raw vegetables. *Journal of Food Protection*® 2009; 72(7): 1486-1490.
29. Ayala-Zavala JF, Wang SY, Wang CY, González-Aguilar GA. Effect of storage temperatures on antioxidant capacity and aroma compounds in strawberry fruit. *LWT-Food Science and Technology* 2004; 37(7): 687-695.
30. Gil MI, Aguayo E, Kader AA. Quality changes and nutrient retention in fresh-cut versus whole fruits during storage. *Journal of Agricultural and Food Chemistry* 2006; 54(12): 4284-4296.
31. Jimenez A, Romojaro F, Gomez JM, Llanos MR, Sevilla F. Antioxidant systems and their relationship with the response of pepper fruits to storage at 20 C. *Journal of Agricultural and Food Chemistry* 2003; 51(21): 6293-6299.
32. Sacchetti G, Cocci E, Pinnavaia G, Mastrocola D, Rosa MD. Influence of processing and storage on the antioxidant activity of apple derivatives. *International Journal of Food Science & Technology* 2008; 43(5): 797-804.
33. Bartolini S, Viti R, Zanol G. Apricot cultivars and cold storage affect the total antioxidant capacity and glutathione content in fruit. *XIII International Symposium on Apricot Breeding and Culture* 2005; 717: 359-362.
34. Hartmann A, Patz C-D, Andlauer W, Dietrich H, Ludwig M. Influence of processing on quality parameters of strawberries. *Journal of Agricultural and Food Chemistry* 2008; 56(20): 9484-9489.