

## Research Article

# Antiasthmatic Activity of Methanolic Extract of *Sphaeranthus indicus*

M. S. Prajapati\*, M. B. Shah, A. K. Saluja, U. D. Shah, S. K. Shah

Dept. of Pharmacognosy and Phytochemistry, A. R. College of Pharmacy & G. H. Patel Institute of Pharmacy,  
Vallabh vidyanagar -388120 (Gujarat)

### ABSTRACT

*Sphaeranthus indicus* auct. Non L. (Asteraceae) commonly known as Gorkhmundi. It is used for various ailments in Indian system of traditional medicines. Methanolic extract of aerial parts was evaluated for antiasthmatic activity. Histamine induced bronchospasm in guinea pigs and egg albumin induced bronchospasm in guinea pig model were performed and various parameters were investigated like PCD time, serum bicarbonate level, differential leukocytes count and histopathological changes in lung. Biochemical estimation like glutathione (GSH) level, malondialdehyde (MDA) level and total proteins were also checked. The result showed that it significantly increased preconvulsion dyspnoea (PCD) time, decreased differential leukocyte counts and serum bicarbonate level. It decreased the oxidative burden by reducing the MDA level and increasing the GSH level which was imbalanced in asthmatic condition, also shown significant action on total protein level. The drug also reduced the inflammation and dilated the bronchioles. All these data suggested that *S. indicus* is having promising antiasthmatic activity.

**Key words:** *S. indicus*, Histamine, Antiasthmatic, Asteraceae and Bronchospasm

### INTRODUCTION

Asthma is a chronic inflammatory disorder of the airways in which many cells and cellular elements play a role, in particular, mast cell, eosinophils, T-lymphocytes, macrophages, neutrophils, and epithelial cells<sup>1</sup>. The pathophysiological changes occur in asthma is like airway obstruction due to combination of various factors include spasm of airway smooth muscle, edema of airway mucosa, increased mucus secretion, cellular infiltration of the airway walls and injury and desquamation of the airway epithelium<sup>2,3</sup>.

*Sphaeranthus indicus* auct. Non L. (Asteraceae) commonly called as Gorkhmundi is found to be grown in rice fields, dry waste places and cultivated throughout India in rainy season, Sri Lanka, Africa and Australia from sea level to 1200m altitude<sup>4</sup>. It is a much branched, strongly scented annual erect herb most commonly grow as a weed. The plant flowers in October- January and fruits are formed from December – April<sup>5</sup>. According to Ayurveda, this herb is hot, laxative, digestible, tonic, fattening, alterative, anthelmintic and alexipharmic. It is used in tuberculosis, bronchitis, anemia, piles, asthma, leucoderma, etc. All the parts of the plant have medicinal value. The juice of the plant is styptic and said to be useful in liver and gastric disorders<sup>6</sup>. The sesquiterpene glycoside isolated from *S.indicus* is having immunostimulant effect<sup>7</sup>. The plant is reported to possess anti-inflammatory, antimicrobial activity and wound healing property<sup>8,9</sup>. Considering the ayurvedic uses and reported activities of the drug, it is proposed to investigate its antiasthmatic potential using various experimental models.

### MATERIAL AND METHODS

All chemicals and solvents used were of analytical grades. Ketotifen fumarate was obtained as a gift sample from FDC Ltd., Mumbai. Histamine hydrochloride was purchased from Chitichem, Baroda. Egg albumin, sodium dodecyl sulfate, acetic acid, hydrochloric acid, thiobarbituric acid (TBA), trichloroacetic acid, disodium hydrogen phosphate, 5, 5'-dithiobis-2-nitrobenoic acid, sodium-potassium tartrate, sodium hydroxide, sodium carbonate, Folin-phenol reagent were of analytical grade.

### Plant material and Extraction

Fully grown, fresh entire herb of *Spharaenthus indicus* auct. Non L. was collected in the month of December from Vallabh Vidyanagar. The herb was authenticated by a taxonomist at the Department of Bioscience, Sardar Patel University, and Vallabh Vidyanagar. A voucher specimen (no argh-7) was deposited in the Department of Pharmacognosy, A. R. College of Pharmacy, Vallabh Vidyanagar. The aerial parts were cleaned, dried at room temperature, powdered to coarse powder (60 #) and stored in air tight container. 5 kg of dried aerial parts of *Sphaeranthus indicus* was exhaustively extracted by subjecting to reflux below 80°C (3 X 5 L). The combined methanol extract was distilled and evaporated to dryness (25 %w/w). A weighed quantity of the extract was taken and suspension was prepared using Tween 80 and was used for further study.

### Animals

Guinea pigs (450-600 gms) were procured from Animal Vaccine Institute, Gandhinagar. All animals were housed at ambient temperature (21±10C) and relative humidity (55±5 %) with fixed 12h/12h light/dark cycle. Animals had free access to standard pellet diet and water given *ad*

\*Author for correspondence: [manishasprajapati@gmail.com](mailto:manishasprajapati@gmail.com)

libitum. The experimental protocol was approved by Institutional Animal Ethical Committee as per the guidance of committee for the purpose of Control and Supervision of Experiments on Animals (CPCSEA), Ministry of Social Justice and Empowerment, Government of India.

**Table: 1** Effect of *Sphaeranthus indicus* on PCD time in guinea pigs after acute treatment.

Treatment group	% increase in time of PCD
Normal control group	5.17 ± 2.54
Ketotifen group (1mg/kg)	87.35±3.80 <sup>@</sup>
MSI group (150mg/kg)	64.26±2.62 <sup>@</sup>
MSI group (300mg/kg)	36.34±4.8 <sup>@</sup>

MSI group (450mg/kg) 29.63±3.9<sup>@</sup>  
<sup>@</sup>Significantly different from control group (p < 0.05)  
 MSI- Methanolic extract of aerial part of *S. indicus*

### In vivo antiasthmatic activity

#### Histamine induced bronchospasm in guinea pig

The guinea pigs (Animal breeding centre) were kept in a closed chamber and exposed to an aerosol of 0.1 percent histamine hydrochloride and time for preconvulsion dyspnoea was noted. As soon as preconvulsion dyspnoea (PCD) commenced, animals were removed from the chamber and placed in fresh air to recover. This time for PCD was taken as basal value. After 15 days of wash out period the same animals were randomly divided into five groups each containing six animals. Group I received distilled water (normal control); Group II received Ketotifen (1 mg/kg, P.O.) (Standard control) and Group III, IV and V received methanolic extract of *S. indicus* (MSI group) in divided doses 150, 300 and 450 mg/kg respectively. The suspension of drugs was administered orally. Two hours after the drugs treatment, animals were exposed to histamine aerosol and times for PCD were noted. The effect of drug was calculated by the following formula<sup>10</sup>

$$\% \text{ increase in PCD time} = [1 - T1/T2] \times 100$$

Where, T1 = time for PCD onset on day 0, T2 = time for PCD onset after drug treatment.

#### Egg albumin induced bronchospasm in conscious Guinea pigs

Hartley strain Guinea pigs of either sex weighing 400-600 gm were selected. The animals were examined for the following parameters:

Eosinophil and macrophages count in the bronchoalveolar fluid

Serum bicarbonate level

Measurement of MDA level


Measurement of GSH level

Measurement of total protein level

Histopathology

The animals were randomly divided into three groups each containing six animals. Group I received distilled water (normal control), Group II received egg albumin

(sensitized group) and Group III received methanolic extract of *Sphaeranthus indicus* (MSI group) (150mg/kg).



**Fig. 1** Histopathological studies

#### Eosinophil and macrophages count in the bronchoalveolar fluid

The suspension of drugs was administered orally. The guinea pigs of all the groups except Group I were sensitized with egg albumin (1 ml, 10% w/v). The animals of Group III was dosed once daily for fifteen days with the suspension of drug. Two hours after the last dose of drug administration on fifteen days all the animals except Group I animals were challenged with egg albumin (0.5 ml, 2% w/v) through saphenous vein. After 3 hours of the challenge of the egg albumin or just prior to death of animals, whichever was earlier, the tracheobronchial tree was lavaged with 10 ml of saline and the fluid so collected was centrifuged at 2000 rpm for 5 minutes and the pellet was resuspended in 0.5 ml saline. 0.2 ml of geimsa stain in buffered saline (pH 6.5) was added to it. After 5 minute the number of each type of leukocytes in 0.5 ml fluids was determined under the microscope 450 X magnifications. The results obtained were compared with unsensitized and untreated egg albumin sensitized guinea pig.<sup>11</sup>

#### Serum bicarbonate level

Serum is suitable specimens for measurement of bicarbonate level. The whole blood should be collected and handled anaerobically to minimize exposure to air. Serum bicarbonate is stable for one hour when stored under anaerobic conditions in an ice bath. Lable one test

tube or cuvette for a reagent blank. Calibrator, controls and unknown sample. Add 1 ml of completely dissolved reagent to each tube or Cuvette. Add 0.01 ml of H<sub>2</sub>O to the reagent blank and 0.01 ml of calibrator, controls and unknown sample to the appropriately labeled tube. Mix and incubate for 5min and measure the absorbance at 380 nm.<sup>12</sup>

**Table: 2 Effect of MSI on differential leukocyte counts of the bronchoalveolar lavage (BAL) fluid**

	Normal control group	Sensitized group	MSI group (150mg/kg)
TLC/cmm	7134.16± 101.79	13440.16 ± 315.9 <sup>#</sup>	11422.5 ± 537.18 <sup>§</sup>
Polymorphs count/cmm	20.5 ± 0.83	37.74 ± 0.93 <sup>#</sup>	34.09 ± 1.29 <sup>§</sup>
Lymphocyte count/cmm	49.79 ± 0.66	58.86 ± 0.76 <sup>#</sup>	52.18 ± 1.44 <sup>§</sup>
Eosinophil count/cmm	6.21 ± 0.31	14.05 ± 0.62 <sup>#</sup>	8.98 ± 0.57 <sup>§</sup>
Monocyte count/cmm	4.83± 0.33	13.41 ± 0.87 <sup>#</sup>	7.01 ± 0.91 <sup>§</sup>

<sup>#</sup> Significantly different from control (p<0.001)

<sup>§</sup> Significantly different from sensitized (p<0.05)

MSI- Methanolic extract of aerial part of *S. indicus*

**Table: 3 Effect of MSI on serum bicarbonate level**  
**Serum bicarbonate level (meq/l)**

Normal control group	40.44 ± 0.70 meq/l
Sensitized group	51.31 ± 1.02 meq/l
MSI group (150mg/kg)	39.89 ± 1.20 <sup>#</sup> meq/l

<sup>#</sup> Significantly different from sensitized (p<0.05)

MSI- Methanolic extract of aerial part of *S. indicus*

#### Measurement of Malondialdehyde (MDA) level

One gm of lung tissue homogenized in 10 ml ice cold Phosphate buffer. The prepared homogenates were centrifuged and used for the determination of antioxidant parameters.

Supernatant (0.2 ml) was mixed with 0.2 ml of 8 percent W/V sodium dodecyl sulfate, 1.5 ml of 20 percent acetic acid in 0.27 M hydrochloric acid, 1.5 ml freshly prepared of thiobarbituric acid (TBA) (1percent W/V) and 0.6 ml of distilled water. The mixture was heated in a water bath at 95°C for 45 minute, cooled and 2 ml of the mixture was mixed with 2 ml of 10 percent trichloroacetic acid. The resulting mixture was centrifuged at 1000 rpm for 5 min. the intensity of pink

color developed was read against blank at 532 nm. The amount of (MDA) (thiobarbituric acid reactive material) was calculated and reported as µg of MDA / mg protein.<sup>13,14</sup>

#### Measurement of GSH level

The supernatant (2 ml) was mixed with 10 percent chilled trichloroacetic acid. The mixture was kept in ice bath for 30 min and centrifuged at 1000 g for 10 min at 4 °C. Supernatant (0.5ml) was mixed with 2.0 ml 0.3 M disodium hydrogen phosphate and 0.25 ml 5, 5'-dithiobis-2-nitrobenoic acid (40 mg/100 ml in 1 percent sodium citrate) was added just before measuring the absorbance at 412 nm. Standard curve for GSH was prepared using glutathione. Results are expressed as µg of GSH /mg protein.

#### Measurement of total protein level

0.2 ml of crude homogenate was added to 4.0 ml of solution – C (solution A- 2 g of sodium hydroxide, 10g of sodium carbonate, 0.1 g of sodium-potassium tartrate in 500 ml of distilled water; solution B- 0.5 g of cuprous sulphate in 100 ml of distilled water; Solution C – (mixture of 10 ml of solution A and 0.2 ml of solution B) and 0.6 ml distilled water was added and allowed to stand for 15 min at 37 C. Folin-phenol reagent 0.4 ml was added and incubated for 30 min. Absorbance was read at 540 nm. Amount of protein was calculated in mg/ml of tissue homogenate from the graph of standard albumin.

#### Histopathology of Lung

Histopathological study of bronchi and bronchioles were carried out to study the effects of MSI on bronchocontractive effects brought about by sensitization of the animals.

#### Statistical Analysis

All the values were expressed as mean ± SEM of six observations. The statistical analysis was performed using Student-t-test. Value of p less than 5 % (p < 0.05) was considered statistically significant.

#### Results

MSI in different concentrations were tested for its antiasthmatic activity in two different *in vivo* models and various parameters were checked.

Guinea pigs were administered with Ketotifen (1mg/kg) and MSI (150, 300, 450 mg/kg) and PCD time was recorded. With histamine aerosol, the percentage increase in PCD time of *Sphaeranthus indicus* (MSI) gives significant result in a dose of 150 mg/kg (64.26 ± 2.64) (p < 0.05). (Table 1)

**Table: 4 Effect of MSI on total protein, malondialdehyde and glutathione level**

	Protein (mg/ml)	MDA (µg/mg/protein)	GSH (µg/mg/protein)
Normal control group	6.80 ± 0.056	0.076 ± 0.0021	16.17 ± 0.019
Sensitized group	9.57 ± 0.017 <sup>§</sup>	0.14 ± 0.0002 <sup>§</sup>	2.7 ± 0.006 <sup>§</sup>
MSI group (150mg/kg)	6.76 ± 0.037 <sup>#</sup>	0.064 ± 0.0017 <sup>#</sup>	11.92 ± 0.08 <sup>#</sup>

<sup>#</sup> Significantly different from sensitized (p<0.05)

<sup>§</sup> Significantly different from control (p<0.001) (student's t test)

MSI- Methanolic extract of aerial part of *S. indicus*

The smear of the bronchoalveolar lavage (BAL) fluid showed different types of White Blood Cells. The total leukocyte count, eosinophil, monocyte, polymorphs and lymphocyte count revealed a change after sensitization of guinea pigs with egg albumin. There was significant ( $p < 0.001$ ) increase in the total leukocyte, polymorph, eosinophil, monocyte and lymphocyte counts in the sensitized animals ( $13440.16 \pm 315.9$ ), ( $37.74 \pm 0.93$ ), ( $14.05 \pm 0.62$ ), ( $13.41 \pm 0.87$ ), ( $58.86 \pm 0.76$ ) as compared to control animals ( $7134.16 \pm 101.79$ ), ( $20.5 \pm 0.83$ ), ( $6.21 \pm 0.31$ ), ( $4.83 \pm 0.33$ ), ( $49.79 \pm 0.66$ ) respectively. The treatment with MSI produced a significant ( $p < 0.05$ ) decrease in the elevated total leukocyte, polymorph, eosinophil, monocyte and lymphocyte counts ( $11422.5 \pm 537.18$ ), ( $35.77 \pm 1.50$ ), ( $8.98 \pm 0.57$ ), ( $7.01 \pm 0.91$ ), ( $52.18 \pm 1.44$ ) as compared to sensitized animals respectively. (Table 2)

Significant ( $p < 0.001$ ) increase in the serum bicarbonate level in the sensitized animals ( $51.31 \pm 1.02$  meq/l) as compared to control animals ( $40.44 \pm 0.70$  meq/l). The treatment with MSI produced a significant ( $p < 0.05$ ) decrease in the elevated serum bicarbonate level ( $39.89 \pm 1.20$  meq/l) as compared to sensitized animals. (Table 3)

In Guinea pig sensitized with egg albumin showed significant ( $0.14 \pm 0.0002$ ) ( $p < 0.001$ ) increase in level of the lipid peroxidation product malondialdehyde (MDA) compared to normal control group which indicated high oxidation procedure in asthma stage. While sensitized animals treated with MSI showed significant ( $0.064 \pm 0.0017$ ) ( $p < 0.05$ ) decrease in level of the lipid peroxidation product malondialdehyde compare to sensitized animal. (Table 4)

Further sensitized animals showed significant ( $2.7 \pm 0.006$ ) ( $p < 0.001$ ) decrease in the level of glutathione which was reversed by the treatment with MSI significantly ( $11.92 \pm 0.08$ ) ( $p < 0.05$ ) increases the GSH level. Moreover elevated level in protein in sensitized animals were reduced by (Table 4)

In histopathological study of lungs of sensitized guinea pigs showed retained of normal vasculature and cellular structure with dilated blood vessels and very few of chronic inflammatory cells compared to sensitized guinea pigs which showed pneumonic patch formed by mixed inflammatory cells and showed bronchoconstriction. (Fig. 1)

#### Discussion

Bronchial asthma is commonly characterized by increased airway reactivity to spasmogen. The MSI was found to significantly increase the PCD time against aerosol as compared to normal control animals indicating its bronchodilating activity. The total leukocyte, eosinophil, polymorph, monocyte and lymphocytes counts were increased after allergen challenge in the BAL fluid, thus gave an indication of inflammatory asthma<sup>15,16</sup>. In the sensitized plus treated animals with MSI it was found that extract of *S. indicus* significantly reduced total leukocyte, neutrophil, lymphocyte counts, eosinophil and monocyte counts. This indicated the protective effect by preventing release

of several preformed mediators there by preventing direct damage of airway epithelium. In our study, oxidative stress has been increased in sensitized group of animal. The increased oxidative burden is result both of increased oxidative stress as evidenced by increased MDA and of the decreased antioxidant capacity as evidence by the lowered glutathione while in case of treated with MSI decreased the oxidative burden by reducing the MDA level and increasing the GSH level which is imbalanced in asthmatic condition<sup>17</sup>. In asthmatic condition bicarbonate level was increased due to increased tension of CO<sub>2</sub> in lung. Our drug significantly decreased the increased bicarbonate level which is beneficial in asthma. Further in histopathological study of lung, the treatment with the drug prevented inflammation and bronchoconstriction. Our all findings supported the use of *S. indicus* as traditional medicine in asthma.

#### References

1. M. B. Rosenblatt, E. B. Weiss, M. S. Segal, M. Stein, Bronchial Asthma: Mechanisms and therapeutics, *History of bronchial asthma* 2<sup>nd</sup> ed. Boston, Little, Brown; 1976; 5-17.
2. M. D Bethesda,; NHLBI, National Asthma Education and Prevention Program, Expert Panel Report 2 *Guidelines for the Diagnosis and Management of Asthma*. NIH Publication No. 1997; 97-4051.
3. National Institutes of Health, National Heart, Lung and Blood Institute. National Asthma Education and Prevention Program. Expert Panel Report: Guidelines for the diagnosis and management of asthma update on selected topics. *J Allergy Clin Immunol* 2002; **110**: S142-219.
4. A. Chatterjee and S. Chandraprakashi, "The Treatise on Indian Medicinal Plants", NISCAIR, Delhi 2003
5. G. L. Shah, Flora Of Gujaratat State Part-II, P-619.
6. K. R. Kirtikar and B. D. Basu, "Indian Medicinal Plants" Lalitmohan Basu. Allahabad, India. 1981; Vol-II, 1346-1348.
7. Shekhani, M. S., Shah, P. M., Khan, K. M., Attaurrahman, Yasmin, A., Kazmi, S. U. Immunostimulant sesquiterpene glycoside from *S.indicus* 1990; **29** 8; 2573-2576.
8. Dubey, K. S., Ansari, A. H., Harduha, M. *Asian journal of chemistry*, 2000; **12** 2; 577-578
9. Sadaf, F., Saleem, R., Ahmed, M., Ahmad, S. I., Navaid-UL-Zafar *Journal of Ethnopharmacology*, **107**, 2, 19; 161-163 (2006).
10. H. Gerhard Vogel (Ed.) Drug Discovery and Evaluation Pharmacological Assays Second Completely Revised, Updated, and Enlarged Edition, P-362
11. H. Gerhard Vogel (Ed.) Drug Discovery and Evaluation Pharmacological Assays Second Completely Revised, Updated, and Enlarged Edition, P-379
12. J. F. Zilva, P. R. Pannall, Hydrogen ion Homeostasis: Blood gas level in clinical chemistry

- in diagnosis and treatment. Lloyd-lake London 1979; chapter IV: 78-113.
13. J. A. Buege, S. D. Aust, Microsomal lipid peroxidation method. *Enzymology* 1978; **52**: 302–310.
  14. C. W. Owens, R.V. Belcher, *Biochem. J.* 1965; **94**, 705–711.
  15. Busse, W. W. and Lemanske, R. F. Jr. *Asthma N Engl. J. Med.* 2001; **344**: 350-362
  16. Fireman, P. *Allergy Asthma Proc.* 2003; **24** 2: 79-83
  17. Hulya Ercan, Esra Birben, Evrim A. Dizdar, Ozlem Keskin, Cagatay Karaaslan, Ozge U. Soyer, Raziye Dut, Cansin Sackesen, Tanju Besler, and Omer Kalayci, “Oxidative stress and genetic and epidemiologic determinants of oxidant injury in childhood asthma”, *J allergy clin immunol.* 2006; **18** (5), 1097-1104