

Chemical Constituents of *Rheum ribes* L.

Consolacion Y Ragasa^{1,2,*}, Jariel Naomi B Bacar¹, Maria Margarita R Querido¹, Maria Carmen S Tan¹, Glenn G Oyong³, Robert Brkljača⁴, Sylvia Urban⁴

¹Chemistry Department, De La Salle University, 2401 Taft Avenue, Manila 1004, Philippines

²Chemistry Department, De La Salle University Science & Technology Complex Leandro V. Locsin Campus, Biñan City, Laguna 4024, Philippines

³Biology Department, De La Salle University, 2401 Taft Avenue, Manila 1004, Philippines

⁴School of Science (Discipline of Applied Chemistry and Environmental Science), RMIT University (City Campus), Melbourne 3001, Victoria, Australia

Received: 10th Sept, 16; Revised: 12th Dec, 16; Accepted: 20th Dec, 16; Available Online: 15th January, 2017

ABSTRACT

Chemical investigation of the dichloromethane extract of *Rheum ribes* has led to the isolation of β -sitosteryl-3 β -glucopyranoside-6'-O-fatty acid esters (**1**), β -sitosterol (**2**), phytol fatty acid esters (**3**), triacylglycerols (**4**) and chlorophyllide a (**5**). The structures of **1-5** were identified by comparison of their NMR data with literature data.

Keywords: *Rheum ribes* L., Polygonaceae, β -sitosteryl-3 β -glucopyranoside-6'-O-fatty acid esters, β -sitosterol, phytol fatty acid esters, triacylglycerols, chlorophyllide a

INTRODUCTION

Rheum ribes L. of the family *Polygonaceae*, locally known as "Rivas" is a native plant of Iran which grows in several provinces including Khorasan¹ and in the Middle East². A study reported that palmitic acid [27.08%], n-icosane [9.9%], n-tetracosane [7.34%], linoleic acid [6.56%], and ethyl linoleate [4.76%] were the main components of the oil of *Rheum ribes*. Extracts and fractions from the plant inhibited the growth of the protozoan, *Trichomonas vaginalis*³. Through GC-MS, the constituents of the extract from the flowers of *R. ribes* were characterized as possessing a high quantity of unsaturated fatty acids (66.0 %) and some long chain hydrocarbons. The main components of the hexane extract were 9-octadecenoic acid (ω -9) (42.8 %), 9, 12-octadecadienoic acid (linoleic acid or ω -6) (19.6 %), hexadecanoic acid, (palmitic acid) (8.6 %), 1,2-benzenedicarboxylic acid diisooctyl (5.7 %), dodecane (3.7 %) and γ -linolenic acid (3.6 %). The major constituents of the distilled oil were germacrene D (22.3 %), α -pinene (13.5 %), terpinolene (12.4%), p-cymene (10.6 %), bicyclogermacrene (9.6 %) and limonene (8.6 %). The essential oil and hexane extract exhibited a moderate effect on some Gram-positive and Gram-negative bacteria. The hexane extract of this plant's flowers possessed considerable antioxidant activity⁴. The essential oil of the stalks and flowers of *R. ribes* L., growing in Iran, were extracted via hydro-distillation and analyzed by GC-MS. Thirty constituents representing 93.84% of Rheum oil were identified. The oil was found to be rich in hydrocarbons, especially long-chain n-alkanes (80.81%). The most abundant components in the oil included tricosane (26.29%), heneicosane (26.07%), pentacosane (10.63%), heptacosane (10.37%) and palmitic

acid (3.64%). The essential oil was also evaluated for general toxicity using a bioassay brine shrimp lethality method. The toxicity profile of the oil indicated some degree of toxicity in comparison with podophyllotoxin⁵. *R. ribes* was also reported to contain the anthraquinones, physcion and rhein and the stilbene; rhaponticin or rhapontin⁶. The anthraquinones, chrysophanol, parietin and emodin, the flavonoids quercetin, fisetin, quercetin 3-O-rhamnoside, quercetin 3-O-galactoside and quercetin 3-O-rutinoside were isolated from the shoots of *R. ribes*⁷. Another study reported the isolation of four anthraquinone derivatives (chrysophanol, physcion, rhein and aloe-emodin), two anthraquinone glucosides (physcion-8-O-glucoside and aloe-emodin-8-O-glucoside), the dianthron glucoside, sennoside A, and the stilbene glucoside, rhaponticin from the roots and rhizomes⁸. The ethyl acetate extracts of *R. ribes* shoot and root dry powder were shown to be potential scavengers of DPPH radicals (IC₅₀ value of 206.28 μ g/mL for the shoots and 10.92 μ g/mL for the roots). *R. ribes* inhibited the survival of HL-60 cells in a concentration- and time-dependent manner⁹. We report herein the isolation of β -sitosteryl-3 β -glucopyranoside-6'-O-fatty acid esters (**1**), β -sitosterol (**2**), phytol fatty acid esters (**3**), triacylglycerols (**4**) and chlorophyllide a (**5**) from *Rheum ribes*. The structures of **1-5** are shown in Fig. 1.

MATERIALS AND METHODS

General Experimental Procedure

¹H (500 MHz) NMR spectra were acquired in CDCl₃ on a 500 MHz Agilent DD2 NMR spectrometer with referencing to solvent signals (δ 7.26). Column chromatography was performed, with silica gel 60 (70-230




Figure 1: Chemical structures of β -sitosterol-3 β -glucopyranoside-6'-O-fatty acid esters (1), β -sitosterol (2), phytol fatty acid esters (3), triacylglycerols (4) and chlorophyllide a (5) from *Rheum ribes*.

mesh). Thin layer chromatography, was performed with plastic backed plates coated with silica gel F254 and the plates were visualized by spraying with vanillin/H₂SO₄ solution followed by warming.

General Isolation Procedure

A glass column 18 inches in height and 1.0 inch internal diameter was packed with silica gel. The crude extracts were fractionated by silica gel chromatography using increasing proportions of acetone in dichloromethane (10% increment) as eluents. Twenty milliliter fractions were collected. All fractions were monitored by thin layer chromatography. Fractions with spots of the same R_f values were combined and rechromatographed in

appropriate solvent systems until TLC pure isolates were obtained. A glass column 12 inches in height and 0.5 inch internal diameter was used for the rechromatography. Five milliliter fractions were collected. Final purifications were conducted using Pasteur pipettes as columns. One milliliter fractions were collected.

Plant Material

Plant specimens were collected from the Zagros mountain areas of West Azerbaijan, Iran, with the kind assistance of Mr. Saeed Karami Ishghlo of Payame Noor University of Bukan. The identity of the samples was confirmed in comparison with voucher specimens in the Central Herbarium of Tehran University.

Isolation of the Chemical Constituents

The freeze-dried sample (53.9 g) was soaked in CH₂Cl₂ for three days and then filtered. The filtrate was concentrated under vacuum to afford a crude extract (0.4825 g) which was chromatographed using increasing proportions of acetone in CH₂Cl₂ (10% increment) as eluents. The CH₂Cl₂ fraction was rechromatographed (2 ×) using 2.5% EtOAc in petroleum ether to yield **3** (2 mg) after washing with petroleum ether. The 10% acetone in CH₂Cl₂ fraction was rechromatographed (3 ×) using 5% EtOAc in petroleum ether to yield afford **4** (5 mg). The 20% acetone in CH₂Cl₂ fraction was rechromatographed (3 ×) using CH₃CN:Et₂O:CH₂Cl₂ (0.5:0.5:9, v/v) to afford **2** (5 mg) after washing with petroleum ether. The 70% acetone in CH₂Cl₂ fraction was rechromatographed (4 ×) using CH₃CN:Et₂O:CH₂Cl₂ (2:2:6, v/v) to afford **1** (4 mg) after trituration with petroleum ether. The 80% to 90% acetone in CH₂Cl₂ fractions were combined and rechromatographed (3 ×) using CH₃CN:Et₂O:CH₂Cl₂ (2.5:2.5:5, v/v) to afford **5** (3 mg) after washing with petroleum ether, followed by diethyl ether.

RESULTS AND DISCUSSION

Silica gel chromatography of the dichloromethane extract of *Rheum ribes* L. afforded **1-5**. The NMR spectra of **1** are in accordance with data reported in the literature for β-sitosterol-3β-glucopyranoside-6'-O-fatty acid esters¹⁰; **2** for β-sitosterol¹¹, **3** for phytol fatty acid esters¹², **4** for triacylglycerol¹³, and **5** for chlorophyll a¹⁴. The fatty acids attached to the triacylglycerol were identified as linolenic acid, linoleic acid and oleic acid based on resonance intensities for the methyl triplet at δ 0.96 (t, *J* = 7.8 Hz), the double allylic methylenes at δ 2.78 and the olefinic protons at δ 5.34 (m) for the linolenic acid; methyl triplet at δ 0.86 (t, *J* = 6.6 Hz), the double allylic methylene at δ 2.80 and the olefinic protons at δ 5.34 (m) for the linoleic acid; and the ; methyl triplet at δ 0.86 (t, *J* = 6.6 Hz) and the olefinic protons at δ 5.34 (m) for the oleic acid¹⁵.

Literature search revealed that **1-2** and **4-5** exhibited diverse biological activities. α-Sitosterol-3α-glucopyranoside-6'-O-palmitate (**1**) was reported to exhibit cytotoxicity against Bowes (melanoma) and MCF7 (breast) cancer cell lines with IC₅₀ values of 152 μM and 113 μM, respectively¹⁶. Furthermore, **1** exhibited cytotoxicity against human stomach adenocarcinoma (AGS) cell line with 60.28% growth inhibition¹⁷. In another study, **1** isolated from *Orostachys japonicus* was found to exhibit potent anti-complement activity (IC₅₀ = 1.0 ± 0.1 μM) on the complement system expressed as total hemolytic activity¹⁸.

β-Sitosterol (**2**) was observed to have growth inhibitory effects on human breast MCF-7 and MDA-MB-231 adenocarcinoma cells¹⁹. It was shown to be effective for the treatment of benign prostatic hyperplasia²⁰. It was also reported to attenuate β-catenin and PCNA expression, as well as quench the radical *in-vitro*, making it a potential anticancer drug for colon carcinogenesis²¹. It can inhibit the expression of NPC1L1 in the enterocytes to reduce intestinal cholesterol uptake²². It has also been reported to

induce apoptosis mediated by the activation of ERK and the downregulation of Akt in MCA-102 murine fibrosarcoma cells²³.

Triacylglycerols (**4**) from Tuna (1000 mg/kg) have been reported to significantly inhibit the tumor growth in the spleen of mice with intrasplenically implanted Lewis lung carcinoma²⁴. Triacylglycerols exhibited antimicrobial activity against *S. aureus*, *P. aeruginosa*, *B. subtilis*, *C. albicans*, and *T. mentagrophytes*²⁵. Another study reported that triacylglycerols showed a direct relationship between toxicity and increasing unsaturation, which in turn correlated with increasing susceptibility to oxidation²⁶. Linoleic acid belongs to the omega-6 fatty acids. It was reported to be a strong anticarcinogen in a number of animal models. It reduces the risk of colon and breast cancer²⁷ and lowers cardiovascular disease risk and inflammations²⁸. Linolenic and linoleic acids inhibited parasites growth by 70% and 64% respectively, against *P. berghei* using the 4-day suppressive test. The two compounds, when used in combination, inhibited the parasites by 96% on day 4 of treatment²⁹.

Chlorophyll and its various derivatives are used in traditional medicine and for therapeutic purposes³⁰. Natural chlorophyll and its derivatives have been studied for wound healing³¹, anti-inflammatory properties³², control of calcium oxalate crystals³³, utilization as effective agents in photodynamic cancer therapy³⁴⁻³⁶, and chemopreventive effects in humans³⁷⁻³⁸. A review on digestion, absorption and cancer preventive activity of dietary chlorophyll has been provided³⁹.

ACKNOWLEDGEMENT

A research grant from the De La Salle University Science Foundation through the University Research Coordination Office is gratefully acknowledged.

REFERENCES

- Zargari A. 5th ed. Vol. 4. Tehran: University Publication; 1992. Medicinal Plants (in Persian) pp. 233–241.
- Kashiwada Y, Nonaka G, Nishioka I, Yamagishi T. Galloyl and hydroxycinnamoyl glucoses from Rhubarb. *Phytochem* 1988; 27:1473-1477.
- Naemi F, Asghari G, Yousofi H, Yousefi HA. Chemical composition of essential oil and anti trichomonas activity of leaf, stem, and flower of *Rheum ribes* L. extracts. *Avicenna J Phytomed.* 2014; 4(3):191-199.
- Amiri N, Shafaghat A, Salimi F. Screening of the essential oil, hexane extract, chemical composition, antioxidant activity, and antimicrobial activity of the flower *Rheum ribes* L. from Iran. *J Essent Oil Bearing Plants* 2015; (18)(5):1108-1115.
- Nikbakht M-R, Esnaashari S, Afshar FH. Chemical composition and general toxicity of essential oil extracted from the stalks and flowers of *Rheum Ribes* L. growing in Iran, *J Rep Pharm Sci* 2013; 2(2):165-170.

6. Agarwal SK, Singh SS, Lakshmi V, Verma S, Kumar S. Chemistry and pharmacology of Rhubarb (*Rheum species*)- A Review. *J Sci Ind Res* 2001; 60:1-9.
7. Tosun F, Akyüz-Kızılay C. Anthraquinones and flavonoids from *Rheum ribes*. Bitkisinin Antrakinonları ve Flavonoidleri Ankara Ecz Fak Derg 2003; 32(1):31-35.
8. Mericli AH, Tuzlaci E. Constituents of *Rheum ribes*. *Fitoter* 1990; 61(4):375-375.
9. Uyar P, Coruh N, İscan M. Evaluation of *in vitro* antioxidative, cytotoxic and apoptotic activities of *Rheum ribes* ethyl acetate extracts. *J Plant Sci* 2014; 2(6): 339-346.
10. Ragasa CY, Ebajo Jr VD, G. Forst. De Los Reyes MM, Mandia EH, Brkljača R, Urban S. Chemical constituents of *Cordia dichotoma* G. Forst. *J Appl Pharm Sci* 2015; 5(Suppl. 2):16-21.
11. Ragasa CY, Lorena GS, Mandia EH, Raga DD, Shen C-C. Chemical constituents of *Abrus precatorius*. *Amer J Essent Oils Nat Prod* 2013; 1(2):7-10.
12. Ragasa CY, Caro JL, Lirio LG, Shen C-C. *Res J Pharm Biol Chem Sci*. Chemical constituents of *Coix lacrymajobi*. *Res J Pharm Biol Chem Sci* 2014; 5(6):344-348.
13. Ragasa CY, Ng VAS, Agoo EM, Shen C-C. Chemical constituents of *Cycas lacrimans*. *Int J Pharmacog Phytochem Res* 2015; 7(3):616-620.
14. Ragasa CY, de Jesus J. Porphyrins and polyprenols from *Macaranga tanarius*. *Res J Pharm Biol Chem Sci* 2014; 5(3):701-708.
15. Ragasa CY, Ng VAS, Torres OB, Sevilla NSY, Uy KVM, Tan MCS, Noel MG and Shen C-C. Sterols, Triglycerides and essential fatty acid constituents of *Brassica oleracea* varieties, *Brassica juncea* and *Raphanus sativus*. *J Chem Pharm Res* 2013; 5(12):1237-1243.
16. Nguyen AT, Malonne H, Duez P, Vanhaelen-Fastre R, Vanhaelen M, Fontaine J. Cytotoxic constituents from *Plumbago zeylanica*. *Fitoter* 2004; 75:500 -504.
17. Tsai P-W, de Castro-Cruz K, Shen C-C, Ragasa CY. Chemical constituents of *Ficus odorata*. *Pharm Chem J* 2012; 46(4):225-227.
18. Yoon NY, Min BS, Lee HK, Park JC, Choi JS. A potent anticomplementary acylated sterol glucoside from *Orostachys japonicus*. *Arch Pharmacol Res* 2005; 28(8):892-896.
19. Awad AB, Chinnman M, Fink CS, Bradford PG. Beta-sitosterol activates Fas signaling in human breast cancer cells. *Phytomed* 2007; 14:747-754.
20. Jayaprakasha GK, Mandadi KK, Poulouse SM, Jadegoud Y, Gowda GA, Patil BS. Inhibition of colon cancer growth and antioxidant activity of bioactive compounds from *Poncirus trifoliata* (L.) Raf. *Bioorg Med Chem* 2007; 15:4923-4932.
21. Baskar AA, Ignacimuthu S, Paulraj G, Numair K. Chemopreventive potential of β -Sitosterol in experimental colon cancer model - an *in vitro* and *in vivo* study. *BMC Comp Alt Med* 2010; 10:24.
22. Jesch ED, Seo JM, Carr TP, Lee JY. Sitosterol reduces messenger RNA and protein expression levels of Niemann-Pick C1-like1 in FHs 74 Int cells. *Nutr Res* 2009; 29(12):859-66.
23. Moon DO, Kyeong JL, Yung HC, Young KG. Beta-sitosterol induced-apoptosis is mediated by the activation of ERK and the downregulation of Akt in MCA-102 murine fibrosarcoma cells. *Int Immunopharmacol* 2007; 7:1044-1053.
24. Maeda Y, Sumiyoshi M, Kimura Y. J. Effects of tuna oil on tumor growth and metastasis to liver in intrasplenic Lewis lung carcinoma (LLC)-implanted mice. *Traditional Med* 2004; 21(5):215-220.
25. Ragasa CY, Lorena GS, Mandia EH, Raga DD, Shen C-C. Chemical Constituents of *Abrus precatorius*. *Amer J Essent Oils Nat Prod* 2013; 1(2):7-10.
26. Ferruzzi MG, Blakeslee J. Digestion, absorption, and cancer preventative activity of dietary chlorophyll derivatives. *Nutr Res* 2007; 27:1-12.
27. Chan P, Thomas GN, Tomlinson B. Protective effects of trilinolein extracted from *Panax notoginseng* against cardiovascular disease. *Acta Pharmacol Sin* 2002; 23(12):1157-1162.
28. Whelan J. The health implications of changing linoleic acid intakes. *Prostaglandins Leukot Essent Fatty Acids* 2008; 79(3-5):165-167.
29. Melariri P, Campbell W, Etusim P, Smith P. *In vitro* and *in vivo* antimalarial activity of linolenic and linoleic acids and their methyl esters. *Adv Stud Biol* 2012; 4(7):333-349.
30. Edwards BJ. Treatment of chronic leg ulcers with ointment containing soluble chlorophyll. *Physiother* 1954; 40:177-179.
31. Kephart JC. Chlorophyll derivatives- their chemistry, commercial preparation and uses. *Econ Bot* 1955; 9:3-18.
32. Larato DC, Pfao FR. Effects of a water-soluble chlorophyllin ointment on gingival inflammation. *N Y State. Dent J* 1970; 36:291-293.
33. Tawashi R, Cousineau M, Sharkawi M. Effect of sodium copper chlorophyllin on the formation of calcium oxalate crystals in rat kidney. *Invest Urol* 1980; 18:90-92.
34. Sternberg ED, Dolphin D, Bruckner C. Porphyrin-based photosensitizers for use in photodynamic therapy. *Tetrahedron* 1998; 54:4151-4152.
35. Nourse WL, Parkhurst RM, Skinner WA, Jordan RT. Photodynamic toxicity of porphyrins and chlorins for a human tumor cell line: combined light and concentration dose responses for the retained fraction. *Biochem Biophys Res Commun* 1988; 151:506-511.
36. Henderson BW, Bellnier DA, Greco WR, Sharma A, Pandry RK, Vaughan LA. An *in vivo* quantitative structure-activity relationship for a congeneric series of pyropheophorbide derivatives as photosensitizers for photodynamic therapy. *Cancer Res* 1997; 57:4000-4007.
37. Egner PA, Munoz A, Kensler TW. Chemoprevention with chlorophyllin in individuals exposed to dietary aflatoxin. *Mutat Res* 2003; 52(3):209-216.
38. Egner PA, Wang JB, Zhu YR, Zhang BC, Wu Y, Zhang QN. Chlorophyllin intervention reduces aflatoxin-

DNA adducts in individuals at high risk for liver cancer. *Proc Natl Acad Sci* 2001; 98(25):1401-1406.

39. Ferruzzi MG, Blakeslee J. Digestion, absorption, and cancer preventative activity of dietary chlorophyll derivatives. *Nutr Res* 2007; 27:1-12.