

In Vitro Antibacterial Screening of Fatty Acid Fractions from Three Different Microalgae

Saritha S^{1*}, Kala K J², Prashob Peter K J¹, S M Nair¹

¹Department of Chemical Oceanography, School of Marine Sciences, Cochin University of Science and Technology, Cochin, India

²Inter University Center for Development of Marine Biotechnology, Cochin University of Science and Technology, Cochin, India

Received: 23rd Sep, 17; Revised 1st Nov, 17, Accepted: 14th Nov, 17; Available Online: 25th Nov, 17

ABSTRACT

The spectrum of antibacterial activities shown by fatty acid fractions of different microalgae, depending on the unsaturation pattern and chain length are still need to be explored. This study provides information on the fatty acid composition of three microalgae, *Chlorella marina*, *Nannochloropsis oculata* and *Chaetoceros affinis* along with their antibacterial activities against selected clinical and fish pathogens. PUFA content of fatty acid fractions followed the order *N. oculata* > *C. affinis* > *C. marina*. Greater abundance of eicosapentanoic acid found in *Nannochloropsis oculata* ($27.39 \pm 0.15\%$) and *Chaetoceros affinis* ($25.34 \pm 0.23\%$) might be responsible for higher bacterial inhibition shown by these fractions.

Key words: Microalgae, Fatty acid, Eicosapentanoic acid, Antibacterial activity.

INTRODUCTION

Fatty acids represent an important class of metabolites due to their multifarious applications based on the variations in structure and relative abundance^{1,2}. Fatty acid profiling provides a tool for studying chemotaxonomic features in various species of microalgae³. Variations in relative abundance of fatty acids in order with environmental factors described their application as stress responsive biomarkers of microalgae^{4,5}. Microalgae are considered as important producers of poly unsaturated fatty acids (PUFA) because a major portion of PUFA in aquatic organisms, including fish, is accumulated from their microalgal diet. Also fatty acid (FA) profile in microalgae is less susceptible to seasonal variations and chemical contaminations⁶.

Understanding the distribution of FAs in microalgae is essential owing to numerous health benefits they possess⁷. Long chain poly unsaturated FAs (LC-PUFA) were reported to have significant anti-bacterial activity compared to saturated FAs⁸. But, studies exhibiting the spectrum of activities shown by microalgal FA fractions are limited. Therefore, this study focuses on the fatty acid composition of four microalgae and their spectrum of activities against selected clinical and fish pathogens. We have tried to analyse the activity variations of each fraction in connection with their unsaturation patterns.

MATERIALS AND METHODS

Algal growth and experiment design

Microalgae, *Chlorella marina*, *Chaetoceros affinis* and *Nannochloropsis oculata* were isolated from Cochin

Estuary, a tropical ecosystem, South west coast of India. Isolated microalgal strains were cultured in 2L of sterilized water of desirable salinity, enriched with suitable nutrient media. Details of media and respective conditions for each species are presented in Table 1. The microalgae were harvested in the stationary phase.

Extraction of fatty acids

Fatty acid composition of dried algal biomass was determined and expressed as FA methyl esters (FAME)^{11,12}. Lyophilized microalgal samples were ultrasonicated with DCM : Methanol (2:1) and the solvent was removed under reduced pressure. The lipid extract thus obtained was hydrolysed using 6% methanolic KOH to convert fatty acids into their potassium salts. Neutral lipids were separated by solvent extraction. Polar fraction was acidified with 6 N HCl and the resulting free fatty acids were extracted with CH₂Cl₂. Crude fatty acid samples obtained from *Chlorella marina*, *Nannochloropsis oculata* and *Chaetoceros affinis*, were named as FA1, FA2 and FA3 respectively.

Gas chromatographic-mass spectrometric analysis

Dried fatty acid samples were converted into fatty acid methyl esters (FAME) using BF₃/methanol and used for GC-MS analysis. GC-MS analysis was performed on gas chromatography (Perkin Elmymer Clarus 680) coupled with mass spectrometer (Perkin Elmymer Clarus 600). 1 μ L of sample was injected to GC equipped with non-polar HP ultra-double fused silica capillary column. Initially, the temperature was increased from 50 °C to 200 °C at a rate of 2 °C per min and held at 200 °C for 5 min. Then, the temperature was again increased from 200 °C to

Table 1: Growth conditions of selected microalgae.

Scientific name /Class	Media	Temperature (°C)	Salinity(psu.)	Light:Dark (hrs.)	Illustration
<i>Chorella marina</i> (Chlorophyceae)	Walne's ⁹	22	30	12:12	
<i>Chaetoceros affinis</i> (Bacillariophyceae)	F/2 ¹⁰	22	30	12:12	
<i>Nannochloropsis oculata</i> (Eustigmatophyceae)	F/2 ¹⁰	22	30	12:12	

Table 2: Fatty acid composition of selected microalgae.

Fatty acid	Relative %		
	<i>Chlorella marina</i> FA1	<i>Nannochloropsis oculata</i> FA2	<i>Chaetoceros affinis</i> FA3
12:0	0.21	0.21	0.27
14:0	15.16	5.54	5.15
16:0	27.66	30.22	31.47
18:0	1.85	2.14	3.27
14 : 1 n-7	0.13	0.12	0.31
16 :1 n-7	1.33	18.23	18.91
18 :1 n-9	21.66	6.58	5.94
18:1 n-11	0.89	n.d	n.d
18:2 n-6	20.48	3.63	4.48
18:3 n-3	2.45	1.12	2.31
20:4 n-6	0.21	4.9	1.81
20:5 n-3	5.23	27.39	23.54
22:6 n-3	3.63	0.63	0.87
ΣSFA	44.88	38.11	40.16
ΣMUFA	24.01	24.93	24.85
ΣPUFA	32	37.67	33.01
ΣEPA + DHA	8.86	28.02	24.41
Σ3-Hydroxy Fatty acids	n.d	n.d	n.d
U.I. ^A	120.32	195.88	171.21

^AU.I. (unsaturation index) was calculated by multiplying the percentage of each fatty acid by the number of double bonds followed by summing up these contributions.

280 °C at a rate of 10 °C per min and held at 280 °C for 10 min. MS operating parameters were as follows: ionization energy - 70 eV; ion source temperature - 200 °C, solvent delay - 4 min and scan range 40–600 u. Identification of the components was based on direct comparison of mass spectral data with NIST library version 2.1

Assay for antimicrobial effect of fatty acid fractions through disc diffusion

Bacterial culture

Clinical pathogens such as *Enterobacter spp.*, *Escherichia coli*, *Klebsiella pneumonia*, *Vibrio cholera*, *Streptococcus*, *Proteus spp.*, *Pseudomonas aeruginosa*, *Salmonella typhi*, *Staphylococcus aureus*, *Shigella sonnaie* and fish pathogens such as *Vibrio alginolyticus*, *Vibrio parahaemolyticus* and *Aeromonas sp.* were selected

for antibacterial activity studies. Collections were reconstituted in Muller-Hinton's broth and cultured under aerobic conditions at 37 °C and 200 rpm to reach exponential growth. The concentration (10⁷ colony-forming units (CFU)/mL) was routinely estimated by spectrophotometric turbidity measurement at 600 nm on a UV/Vis spectrophotometer (Spectro 200 Plus) and by CFU counts on tryptic soy agar (TSA). Microorganism suspension was smeared thoroughly on the test media before assaying the activity of fatty acid samples.

Disc diffusion assay

Samples were dissolved in DMSO and evaluated for its antimicrobial activity. Bacterial cultures in exponential growth phase (10⁸ cells mL⁻¹) were used for cultivation in Muller- Hinton agar plates. 20 µL each of the fatty acid fractions (250 µg fatty acid/disc), were applied in sterile filter paper discs of 6 mm diameter and then placed in the agar medium of bacterial culture. All the plates were incubated at 37 °C for 24 h. Antibacterial activities of the test fatty acids were observed through zone of inhibition (in mm) on the plates. Assays were performed in duplicate with two repetitions.

RESULTS

Fatty acid profiling

Fatty acid compositions of the microalgal species are shown in Table 2. Palmitic acid (C16:0) was found to be the major component in all the three crude fatty acid fractions. In *Chlorella marina* (FA1), the major fatty acids like palmitic acid (C16:0), oleic acid (C18:1 n9), linoleic acid (C18:2 n6) and myristic acid (C14:0) were estimated to be in the range of 27.66%, 21.66%, 20.48%, 15.16% of total fatty acid content respectively. In *Chlorella marina*, both oleic and linoleic acids were found to have similar abundance. Eicosapentaenoic acid (EPA, C20:5 n3) and docosahexaenoic acid (DHA, C22:6 n3) were also observed in small amounts, 5.23 % and 3.63% of total FA content respectively. α-linolenic (18:3 n3), palmitoleic (16:1 n7) and stearic (18:0) acids were also observed in minor amounts viz. 2.45%, 1.33% and 1.85% of total fatty acid content respectively.

Nannochloropsis oculata (FA2) had high amount of C16:0 (30.22 %), C20:5 n-3 (27.39), C16:1 n-7 (18.23%) and C20:4 n-6 (4.9 %). The relative abundance of fatty acids in this species showed that EPA content is fairly high such that it is near to the most abundant palmitic acid. The major fatty acids in FA3 obtained from the diatom *Chaetoceros affinis*, were C16:0 (31.47%), C20:5 n-3 (23.54) and C16:1 n-7 (18.91%). Small quantities of C18:2 n-6 (4.48%), C14:0 (5.15%), 18:1 n-9 (5.94%) and 18:3 n-3 (2.31%) were also found in FA3 fraction. Higher abundance of EPA was also observed in FA3 fraction but

Table 3: Antimicrobial activity of fatty acid fractions against Clinical and fish pathogens.

S.No	Pathogens	Inhibition zone diameter in mm		
		<i>Chlorella marina</i> FA1	<i>Nannochloropsis</i> <i>occulata</i> FA2	<i>Chaetoceros affinis</i> FA3
1	<i>Enterobacter spp</i>	8.2	14.8	14.4
2	<i>Escherichia coli</i>	9.3	16.4	15.8
3	<i>Klebsiella pneumoniae</i>	9.1	15.7	14.9
4	<i>Vibrio cholera</i>	10.9	21.8	20.9
5	<i>Streptococcus</i>	11.2	20.4	18.7
6	<i>Pseudomonas aeruginosa</i>	12.4	21.5	21.1
7	<i>Salmonella typhi</i>	8.5	14.2	13.6
8	<i>Staphylococcus aureus</i>	9.1	20.8	18.4
9	<i>Shigella sonnei</i>	8.4	16.3	15.7
10	<i>Proteus spp</i>	8.7	14.0	13.7
11	<i>Aeromonas</i>	12.5	21.5	20.6
12	<i>Vibrio alginolyticus</i>	11.6	22.3	21.2
13	<i>Vibrio parahaemolyticus</i>	10.8	23.1	20.7

it less than that of FA2. The mass spectra comprising a unique combination of m/z peaks such as m/z 79 (base peak) along with m/z 93, and 107 gave inference to the poly unsaturated fatty acids, particularly a series of methylene interrupted polyenes (Figure 1)¹³.

Antibacterial assay

Inhibition zone diameter (mm) of fatty acid fractions against selected bacteria are given in Table 3. Highest activity was shown by fatty acid fraction FA2 derived from *Nannochloropsis occulata* against all the pathogens selected as shown in Figure 2. Higher zone inhibitions (21.5-23.1 mm) were detected for the fish pathogens (*Vibrio alginolyticus*, *Vibrio parahaemolyticus* and *Aeromonas* sp). Among the clinical pathogens, greater inhibitions were observed towards the gram negative, *Vibrio cholerae* (21.8 mm) and *Pseudomonas aeruginosa* (21.5 mm), and also towards the gram positive *Streptococcus* sp. (20.4 mm), and *Staphylococcus aureus* (19.8 mm).

Fatty acid fraction FA3 from *Chaetoceros affinis* also showed comparable but slightly lesser inhibition than that of FA2 fraction. FA3 also showed higher zone of inhibition towards the fish pathogens (20.6-21.2 mm). Among clinical pathogens, FA3 showed considerable inhibition towards gram negative, *Vibrio cholerae* (20.9 mm) and *Pseudomonas aeruginosa* (21.1 mm), and the gram positive *Streptococcus* sp. (18.7 mm) and *Staphylococcus aureus* (18.7 mm).

Chlorella marina fatty acid fraction FA1 showed much less inhibition, while comparing antibacterial activities of other two fractions. The inhibition diameters of FA1 against the fish pathogens range from 10.8 to 12.5 mm. FA1 also showed minor activity towards clinical pathogens (8.2-12.4 mm).

DISCUSSION

Extensive studies were carried out for the fatty acid composition of microalgae, but studies exhibiting their spectrum of activities are still limited. It was suggested that mixture of fatty acids act synergistically to inhibit bacterial growth and thus more numbers of bacterial strains might be susceptible to fatty acid mixtures

compared to individual fatty acid¹⁴. Microalgae can be used as potential tool for examining antibacterial potential of mixture of fatty acids as the individual members possess a very unique fatty acid composition depending on taxonomic category and growth conditions¹⁵. In the present study *Nannochloropsis occulata* and *Chaetoceros affinis* were found as eminent producers of EPA. Stationary phase of microalgal cultures were known to be associated with high amount PUFA¹⁶ compared with other growth phases. Therefore, all the fatty acid fractions were taken from stationary phase of algae culture. Hence, percentage abundance of PUFA obtained is assumed as the maximum for those species under the selected growth conditions.

It has been reported that antibacterial and antifungal effect of long-chain saturated fatty acids like 16:0 and 18:0 acids is less than that of long-chain unsaturated fatty acids^{17,18,19}. This might be the reason for highest antibacterial activity shown by FA2 fraction derived from *Nannochloropsis occulata*, which is characterised by highest unsaturation index (195.88) compared to other species under study. The class *Eustigmatophyceae*, in which *Nannochloropsis occulata* belongs, are known as good source of EPA²⁰. The greater percentage of EPA (20:5 n-3) along with other long chain FAs (18:1 n-9, 18:2 n-6, 18:3 n-3) might be responsible for the increased rate of inhibition. n-3 PUFAs especially EPA and DHA are known to possess biological activities such as antibacterial, antiviral and antifungal effects^{21,22}. Higher zone of inhibition shown by FA2 fraction towards the fish pathogens is in accordance with the finding that unsaturated long chain fatty acids possess antibacterial activity against aquaculture bacteria such as *Listonella anguillarum* and other *Vibrio spp.*¹⁴. FA3 fraction from the diatom *Chaetoceros affinis* also showed significant activity which might be due to the high EPA content and unsaturation index of 171.21 in the fatty acid profile. The diatoms *Phaeodactylum tricornutum* and *Chaetoceros calcitrans* are known as eminent producers of EPA^{16,23,24}. *Chlorella marina* fatty acid fraction FA1 showed least activity while comparing the antibacterial inhibition of crude fatty acid fractions. The unsaturation index of FA1




Figure 1: A representative mass fragmentation pattern of polyunsaturated fatty acid, methyl 5,18, 11, 14, 17 eicosapentaenoate, having base peak at $m/z = 79$.




Figure 2: Antibacterial activity of fatty acid fractions from *Chlorella marina* (FA1), *Nannochloropsis oculata* (FA2) and *Chaetoceros affinis* (FA3).

(120.32) fraction is less compared with FA2 and FA3, which is due to the less EPA abundance in that fraction. Linoleic acid (18:2 n6), the major contributor for PUFA content in *Chlorella marina* along with small quantities of EPA (20:5 n3), DHA (22:6 n3) and ALA (18:3 n3) might be responsible for the minor antibacterial activity shown by FA1. Linoleic acid is reported to have activity against a few bacterial pathogens in the free form and also in combinations^{25,26}.

CONCLUSION

In conclusion, the fatty acid fraction separated from *Nannochloropsis oculata*, possessing high concentration of EPA was found to have the highest antibacterial activity and can be suggested as an important source for the development of new antibacterial drugs. Bactericidal activities shown by the fatty acid fractions were

dependent on the abundance of long chain unsaturated fatty acids. Activities may vary depending on the matrix associated with the fatty acids and hence the antibacterial activities of fatty acid fractions from different microalgae under different growth conditions are still need to be explored.

ACKNOWLEDGEMENT

Authors are thank full to Amala Cancer Research and Hospital, Thrissur, Kerala, India for providing clinical pathogens and Department of Marine Biology , Cochin University of Science and Technology, Kochi for fish pathogens. One of the authors is grateful to the financial assistance by CSIR (India). We are also thankful to Government of Kerala for the financial support received to establish a new centre, Inter University Centre for Development of Marine Biotechnology (IUCDMB).

CONFLICT OF INTEREST STATEMENT

We declare that we have no conflict of interest.

REFERENCES

- Mata TM, Martins AA, Caetano NS. Microalgae for biodiesel production and other applications: a review. *Renew Sust Energ Rev* 2010;14(1):217-232.
- Griffiths MJ, van Hille RP, Harrison ST. Lipid productivity, settling potential and fatty acid profile of 11 microalgal species grown under nitrogen replete and limited conditions. *J Appl Phycol* 2012;24(5):989-1001.
- Temina M, Rezankova H, Rezanka T, Dembitsky VM. Diversity of the fatty acids of the *Nostoc* species and their statistical analysis. *Microbiol Res* 2007;162(4):308-321.
- Volkman JK, Johns RB. The geochemical significance of positional isomers of unsaturated acids from an inter tidal zone sediment. *Nature* 1977;267(5613):693-694.
- Lu N, Wei D, Jiang XL, Chen F, Yang ST. Fatty acids profiling and biomarker identification in snow alga *Chlamydomonas nivalis* by NaCl stress using GC/MS and multivariate statistical analysis. *Anal Lett* 2012;45(10):1172-1183.
- Guil-Guerrero JL, Belarbi EH. Purification process for cod liver oil polyunsaturated fatty acids. *J Am Oil Chem Soc* 2001;78(5):477-484.
- Guedes AC, Amaro HM, Barbosa CR, Pereira RD, Malcata FX. Fatty acid composition of several wild microalgae and cyanobacteria, with a focus on eicosapentaenoic, docosahexaenoic and α -linolenic acids for eventual dietary uses. *Food Res Int* 2011;44(9):2721-2729.
- Zheng CJ, Yoo JS, Lee TG, Cho HY, Kim YH, Kim WG. Fatty acid synthesis is a target for antibacterial activity of unsaturated fatty acids. *FEBS Lett* 2005;579(23):5157-5162.
- Walne PR. Studies on the food value of nineteen genera of algae to juvenile bivalves of the genera *Ostrea*, *Crassostrea*, *Mercenaria* and *Mytilus*. *Fish Invest Ser2* 1970;26(5).
- Guillard RR, Ryther JH. Studies of marine planktonic diatoms: I. *Cyclotella nana* Hustedt, and *Detonula confervacea* (Cleve) Gran. *Can J Microbiol* 1962;8(2):229-239.
- Bligh EG, Dyer WJ. A rapid method of total lipid extraction and purification. *Can J Biochem Physiol* 1959;37(8):911-917.
- Harvey HR. Fatty acids and sterols as source markers of organic matter in sediments of the North Carolina continental slope. *Deep Sea Res Part II Top Stud Oceanogr* 1994;41(4-6):783-796.
- Fellenberg AJ, Johnson DW, Poulos A, Sharp P. Simple mass spectrometric differentiation of the n-3, n-6 and n-9 series of methylene interrupted polyenoic acids. *Biomed Environ Mass Spectrom* 1987;14(3):127-129.
- Benkendorff K, Davis AR, Rogers CN, Bremner JB. Free fatty acids and sterols in the benthic spawn of aquatic molluscs, and their associated antimicrobial properties. *J Exp Mar Biol Ecol* 2005;316(1):29-44.
- Tonon T, Harvey D, Larson TR, Graham IA. Long chain polyunsaturated fatty acid production and partitioning to triacylglycerols in four microalgae. *Phytochemistry* 2002;61(1):15-24.
- Reboloso-Fuentes MM, Navarro-Pérez A, Ramos-Miras JJ, Guil-Guerrero JL. Biomass nutrient profiles of the microalga *Phaeodactylum tricorutum*. *J Food Biochem* 2001;25(1):57-76.
- Knapp HR, Melly MA. Bactericidal effects of polyunsaturated fatty acids. *J Infect Dis* 1986;154(1):84-94.
- Sun CQ, O'Connor CJ, Robertson AM. Antibacterial actions of fatty acids and monoglycerides against *Helicobacter pylori*. *FEMS Immunol Med Microbiol* 2003;36(1-2):9-17.
- Zheng CJ, Yoo JS, Lee TG, Cho HY, Kim YH, Kim WG. Fatty acid synthesis is a target for antibacterial activity of unsaturated fatty acids. *FEBS Lett* 2005;579(23):5157-5162.
- Cheng-Wu Z, Zmora O, Kopel R, Richmond A. An industrial-size flat plate glass reactor for mass production of *Nannochloropsis* sp. (Eustigmatophyceae). *Aquaculture* 2001;195(1):35-49.
- Desbois AP. Potential applications of antimicrobial fatty acids in medicine, agriculture and other industries. *Recent Pat Antiinfect Drug Discov* 2012;7(2):111-122.
- Grima EM, Belarbi EH, Fernández FA, Medina AR, Chisti Y. Recovery of microalgal biomass and metabolites: process options and economics. *Biotechnol Adv* 2003;20(7):491-515.
- Delaporte M, Soudant P, Moal J, Lambert C, Quéré C, Miner P, Choquet G, Paillard C, Samain JF. Effect of a mono-specific algal diet on immune functions in two bivalve species-*Crassostrea gigas* and *Ruditapes philippinarum*. *J Exp Biol* 2003;206(17):3053-3064.
- Desbois AP, Lawlor KC. Antibacterial activity of long-chain polyunsaturated fatty acids against *Propionibacterium acnes* and *Staphylococcus aureus*. *Mar Drugs* 2013;11(11):4544-4557.
- Greenway DL, Dyke KG. Mechanism of the inhibitory action of linoleic acid on the growth of *Staphylococcus aureus*. *J Gen Microbiol* 1979;115(1):233-245.
- Dilika F, Bremner PD, Meyer JJ. Antibacterial activity of linoleic and oleic acids isolated from *Helichrysum pedunculatum*: a plant used during circumcision rites. *Fitoterapia* 2000;71(4):450-452.