

GC-MS Analysis of Extract of *Rubia tinctorum* having Anticancer Properties

Raghad D H Abdul Jalill*

College of Science AL- Mustansiriyia University; Baghdad/ Iraq.

Received: 7th Feb, 17; Revised: 2nd March, 17; Accepted: 21st March, 17; Available Online: 25th March, 2017

ABSTRACT

Atomic Absorption Spectrometry determined the percentage of Cu, Zn, Cd, Mn, K, Fe, Co, Ti and N. Their amounts were: (0.002281, 0.004115, 0.000205, 0.007772, 0.833, 0.093952, 0.000332, 0.043 and 1.02) %. There was no Phosphorus. Total Crude protein was 5.8125 %. Thirty-five compounds identified in methanol extract of *R. tinctorum* using Gas Chromatography- Mass Spectrum analysis (GC-MS) in a single run. The major three components present were 9,12-Octadecadienoic acid (Z,Z)- (29.75 %); 9-Octadecenoic acid (Z)-, hexadecyl ester (26.1 %) and 1,3-Propanediol, 2-ethyl-2-(hydroxymethyl)- (10.09 %). The results of cytotoxicity on cancer cell lines showed reduction in cell viability of Human Glioblastoma cell line (AMGM) treating with all concentrations of extract with IC₅₀ reached to 76.44 mg/ml. Inhibition rate ranged between (28.54- 66.1) %. In Ahmed-Mohammed-Nahi-2003 (AMN3) cell line, the lower concentration (50 µg/ml-5 mg/ml), reduced cell viably while the higher concentrations (50-100) mg/ml induced it. IC₅₀ is 21 mg/ml, the best Inhibition rate is 93.693% at 50 µg/ml. The extraction was less effect in cell viability of Human Larynx Epidermoid Carcinoma (HEp-2) cell Line, the maximum Inhibition rat was 22.12 % at 10 mg/ml. All other concentrations were not significant compared with control. Previous results of cytogenetic effects of plant extract on human lymphocytes showed non-significant changing in mitotic index, blast index, cell cycle progression and replication index but the extract induced chromosomal aberration and sister chromatid exchange. These inductions in chromosomal aberration were structural. They were: addition, deletion and ring chromosome. The deletions were the most common abnormalities. There were more than two chromosomal aberrations in the same cell.

Keyword: AMGM; AMN3; cell line; cytogenetic; Hep-2; GC-MS; *Rubia tinctorum*.

INTRODUCTIONS

Rubia tinctorum L. (Family: Rubiaceae), is one of Iraqi plants, its origin is native to southern Europe and Asia. Their common name is Madder. All parts of the plant contained an iridoid, asperuloside. Parts used is roots, they contain anthraquinone and their glycosides, including alizarin, purpurin, purpuroxanthin, pseudopurpurin, rubiadin, ruberythric acid and lucidin primeveroside, nordamnacanthal¹⁻⁴. Other studies found other compound in roots and rhizomes such as: di- and tri-hydroxyanthraquinones, alizarin, purpurin and their derivatives, ruberythric acid (alizarin-primeveroside), pseudopurpurin and lucidin-primeveroside⁵.

The action of roots used for menstrual and urinary disorders and liver diseases¹. Madder is used in amenorrhea and dropsy, and when taken into the stomach imparts a red color to the milk and urine, and to the bones of animals, without sensibly affecting any other tissue⁶.

R. tinctorum L. revealed antimicrobial activity against some Gram (+) and Gram (-) bacteria, yeasts, filamentous fungi and actinomycetes⁷⁻⁸. Aqueous extract of this plant has anti-diarrheal effect⁹.

Some study refers that this plant is toxic, their understanding is not very precise because they mixed between natural extractions of plants and dye that extract

from roots of plant using deferent solvents which has long tradition in dyeing processes of textiles and are toxic due to presence of toxic hydroxyanthraquinones¹⁰, such as: alizarin¹¹, and rubiadin¹²⁻¹³, this is different from natural aquatic or methanol crud extractions of plants which are weak toxicity, the results of Ino and others¹⁴ suggest that dietary exposure of madder root (MR) has no acute or subacute toxic effects on mice. In addition, methanol extract of plant roots exerted here weak toxicity targeting liver, kidney, and possibly RBC and WBC in rats, some renal parameters being apparently affected from the 0.6% dose level¹⁵.

Recently, the treatment with the aqueous extract exhibited a strong decrease in two different metastatic potency human melanoma cell lines in 10⁻⁶-10⁻⁵ M concentration range, which activity was demonstrated already after 48 h. This inhibitory effect was more pronounced in A2058 cell line (10⁻⁶-10⁻⁵ M: 87.4-55.0 %) than in HT168-M1 cells (10⁻⁶-10⁻⁵ M: 87.5-63.7 %) after 72 h. In normal fibroblast cells derived from normal lung tissue (MRC-5 cells) a similar, concentration dependent antiproliferative/cytotoxic effect (10⁻⁶-10⁻⁵ M: 77.2-62.0 %) was observed for this aqueous extract¹⁶.

The aim of the present study was to develop an efficient anticancer drug using aquatic roots' extract of *R. tinctorum*

*Author for Correspondence: raghad.dh.abdul.jalill@gmail.com

and if there are any toxicity in cytogenetic of human lymphocyte.

MATERIALS AND METHODS

Plants extraction

The roots of *Rubia tinctorum* plant obtained from Iraq Medical Herbarium/ Ministry of Health/ / Baghdad / Iraq. Roots dried at 38 °C and ground by a grinder. Fifty grams of powdered material extracted by 250 mL of 70% methanol using a Soxhlet extractor at 40 °C for 3 h.¹⁷. The extracts filtered using Whatman No.1, evaporated and sterilized by 0.22 µ micro filters.

Chemical analysis

Atomic Absorption Spectrometer

Atomic Absorption Spectrometer (Nove AA-350, analytik Jena) was used to determine the concentrations of some element in plants. The elements were: Cu, Zn, Cd, Mn, K, Fe, Co, Ti, P and N. The samples prepared according to Kodama in (1963)¹⁸. The level of Nitrogen and total proteins were recorded by Macro Kjeldahl method¹⁹.

Gas Chromatography- Mass Spectrum analysis (GC-MS) analyses

GC-MS analyses were done using Shimadzu GC-2010 Plus coupled with Shimadzu GCMS-Q2010 Ultra. Capillary column (InertCap 1 MS, 0.25 mm, 30 m, 0.25 µm, Gl Sciences, Japan). Carrier gas was helium. Constant flow rate is 1 ml/min and auto injector is AOC-20i, Shimadzu. Injection volume is 5 µl. Column oven temperature program were in the following order: 100 °C. Oven temperature program was: 100 °C for 3 min., 240 °C for 9 min., 280 °C for 5 min. and 300 °C for 2min. The rates were 15. For identification of components, direct comparison of the retention times and mass spectral data of components had done with those for NIST Standard Reference Database 1A: NIST/EPA/NIH Mass Spectral Library (NIST 08) and NIST Mass Spectral Search Program (Version 2.0f)²⁰.

Cytotoxicity on cancer cell lines

Cell culture

Three cancer cell lines obtained from cell bank unite/ Department of Experimental Therapy / Iraqi Center for Cancer and Medical Genetic Research (ICCMGR). The cell lines were: Human Glioblastoma cell line (AMGM) (passage 69), Ahmed-Mohammed-Nahi-2003 (AMN3), murine mammary adenocarcinoma cell line (passage 203) and Human Larynx Epidermoid Carcinoma (HEp-2) Cell Line (passage 275). They maintained using RPMI 1640 (USbiological - USA) supplemented with 15% calf bovine serum (Gibco, USA), 100 units per ml penicillin, and 100 µg/ml streptomycin as recommended by cell bank unite at ICCMGR. The cells seeded in Flat bottomed 96-well polystyrene and incubated at 37°C under a humidified atmosphere containing 5 % CO₂ for 24 hr. A population was 1.5*10⁴ cells per well.

Cytotoxic activity

Cytotoxicity assay had done according to method of Department of Experimental Therapy / Iraqi Center for Cancer and Medical Genetic Research (ICCMGR) with some modifications. The spent media of the confluence cells, (90-95) % within 24 h. incubation, was removed.

Table 1: The level (%) of some elements and total proteins in *R. tinctorum*.

| Elem. | % |
|-------|----------|
| Cu | 0.002281 |
| Zn | 0.004115 |
| Cd | 0.000205 |
| Mn | 0.007772 |
| K | 0.833 |
| Fe | 0.093952 |
| Co | 0.000332 |
| Ti | 0.043 |
| P | NIL |
| N | 1.02 |
| Prot. | 5.8125 |

Elem.: element, %: Percentage, Cu: Copper, Zn: Zinc, Cd: Cadmium, Mn: Manganese, K: Potassium, Fe: Iron, Co: Cobalt, Ti: Titanium, P: Phosphorus, N: Nitrogen, Prot.: Crude protein.

Cells washed with PBS (0.01 M phosphate buffer, 0.0027 M KCl and 0.137 M NaCl) and 0.2 ml of fresh free serum media (RPMI 1640) was added. Series of dilution of crude plant extract, dissolved in PBS which were diluted with free serum medium (RPMI 1640), were added to plates. There was a negative control (PBS) and six replicates for each tested. Plates incubated at 37°C with 5% CO₂ for (24, 48,72) hr. of AMN3, AMGM and Hep-2 respectively. At the end of exposure time, cells washed twice with PBS. A fifty micro letter of Neutral Red solution (Sigma, U.S.A; 50 µg/ml dissolved in PBS), added to each well and incubated at 37°C for two hr.

The wells washed three times with PBS. Fifty microliter of solution (PBS and absolute ethanol 1:1 V/V) added to each well. The optical density read by a Micro-ELISA. Percentage of cell viability²¹ and inhibiting rate²² calculated. In addition to Compusyn Computer software, (version 2011), which used for IC₅₀ calculation.

Cytogenetic effects on human lymphocytes.

Blood samples (0.2 mL) obtained from nonsmokers, 24-25 years' age with no history of drugs, radiation therapy or viral infection. They planted in Roswell Park Memorial Institute (RPMI) 1640 medium (Sigma/ U.S.A) supplemented with: AB⁺ Plasma, 10 µg/mL bromodeoxyuridine (Serva/ Germany) and 0.1 ml of 10 µg/ml of Phytohemagglutinin (Sigma/ U.S.A). Cells treated with (100, 10 and 1) mg/ml of crude plant extract. There were negative controls and positive control (adding Mytomycin C, Kogyo Company, at 50 ng/ml final concentration. There were triplicates for each treatment. All experiments were carried in the dark.

Lymphocytes were cultured for 72 h at 37 °C. For CA and SCE analysis, 0.06 µg/mL colchicine (Sigma/ U.S.A, dissolved in distilled water) was added to cultures at the last two hours of exposure time. Cells were harvested by centrifugation (216×g, 10 min) and suspended in a hypotonic solution of 0.075 M KCl for 30 min at 37 °C. Cells were centrifuged and fixed by cold methanol acetic acid (3:1) for 20 min. This step repeated three times. Solution dropped in clean slides and air drying. For

Table 2: Composition of methanolic extraction of *R. tinctorum*.

| P | R.T | Area % | Comp. | C.F | M.W |
|----|-------|--------|--|--|-----|
| 1 | 5.23 | 0.45 | 1,2-Cyclopentanedione | C ₅ H ₆ O ₂ | 98 |
| 2 | 8.44 | 0.08 | 4H-Pyran-4-one, 2,3-dihydro-3,5-dihydroxy-6-methyl- | C ₆ H ₈ O ₄ | 144 |
| 3 | 10.24 | 0.44 | N-Methyl-4-(3-nitrobenzamido)phthalimide | C ₁₆ H ₁₁ N ₃ O ₅ | 325 |
| 4 | 10.50 | 0.52 | 2-Methoxy-4-vinylphenol | C ₉ H ₁₀ O ₂ | 150 |
| 5 | 10.82 | 0.59 | Ethylene diacrylate | C ₈ H ₁₀ O ₄ | 170 |
| 6 | 10.96 | 0.51 | 3-Allyl-6-methoxyphenol | C ₁₀ H ₁₂ O ₂ | 164 |
| 7 | 11.27 | 0.46 | 6,7,8,9-Tetrahydro-5H-[1,2,4]triazolo[1,5-a]azepin-2-ylamine | C ₇ H ₁₂ N ₄ | 152 |
| 8 | 11.81 | 10.09 | 1,3-Propanediol, 2-ethyl-2-(hydroxymethyl)- | C ₆ H ₁₄ O ₃ | 134 |
| 9 | 12.94 | 0.46 | Pentadecanoic acid | C ₁₅ H ₃₀ O ₂ | 242 |
| 10 | 13.01 | 1.23 | 3-Deoxy-d-mannonic lactone | C ₆ H ₁₀ O ₅ | 162 |
| 11 | 13.29 | 1.35 | 3-Deoxy-d-mannonic acid | C ₆ H ₁₂ O ₆ | 180 |
| 12 | 13.63 | 0.05 | Carbonic acid, allyl nonyl ester | C ₁₃ H ₂₄ O ₃ | 228 |
| 13 | 13.80 | 0.04 | Dodecanoic acid | C ₁₂ H ₂₄ O ₂ | 200 |
| 14 | 14.14 | 0.04 | Thiocyanic acid, 4.alpha.-methyl-5.alpha.-cholestan-3.alpha.-yl ester | C ₂₉ H ₄₉ NS | 443 |
| 15 | 14.33 | 0.10 | 4-((1E)-3-Hydroxy-1-propenyl)-2-methoxyphenol | C ₁₀ H ₁₂ O ₃ | 180 |
| 16 | 14.63 | 0.16 | Tetradecanoic acid | C ₁₄ H ₂₈ O ₂ | 228 |
| 17 | 16.23 | 0.53 | Oleyl alcohol, trifluoroacetate | C ₂₀ H ₃₅ F ₃ O ₂ | 364 |
| 18 | 16.42 | 6.05 | l-(+)-Ascorbic acid 2,6-dihexadecanoate | C ₃₈ H ₆₈ O ₈ | 652 |
| 19 | 16.58 | 0.49 | 4-Methoxy-2-nitroformanilide | C ₈ H ₈ N ₂ O ₄ | 196 |
| 20 | 17.79 | 0.42 | 9,10-Anthracenedione, 1-hydroxy- | C ₁₄ H ₈ O ₃ | 224 |
| 21 | 17.90 | 1.17 | 9,12-Octadecadienoic acid (Z,Z)-, methyl ester | C ₁₉ H ₃₄ O ₂ | 294 |
| 22 | 18.00 | 0.86 | 9-Octadecenoic acid, methyl ester, (E)- | C ₁₉ H ₃₆ O ₂ | 296 |
| 23 | 18.40 | 29.75 | 9,12-Octadecadienoic acid (Z,Z)- | C ₁₈ H ₃₂ O ₂ | 280 |
| 24 | 18.50 | 26.10 | 9-Octadecenoic acid (Z)-, hexadecyl ester | C ₃₄ H ₆₆ O ₂ | 506 |
| 25 | 18.80 | 5.49 | Octadecanoic acid | C ₁₈ H ₃₆ O ₂ | 284 |
| 26 | 19.27 | 1.05 | Methanone, [4-(methoxymethyl)phenyl]phenyl- | C ₁₅ H ₁₄ O ₂ | 226 |
| 27 | 20.85 | 0.15 | Glycidol stearate | C ₂₁ H ₄₀ O ₃ | 340 |
| 28 | 23.74 | 0.59 | Iron, dicarbonyl(eta.5-2,4-cyclopentadien-1-yl)(2,3,3,4,4,5,5,6,6-nonafluoro-1-cyclohexen-1-yl)- | C ₁₃ H ₅ F ₉ FeO ₂ | 420 |
| 29 | 24.54 | 2.59 | 6,9-Octadecadienoic acid, methyl ester | C ₁₉ H ₃₄ O ₂ | 294 |
| 30 | 24.67 | 1.92 | 9-Octadecenoic acid, 1,2,3-propanetriyl ester, (E,E,E)- | C ₅₇ H ₁₀₄ O ₆ | 884 |
| 31 | 25.78 | 0.67 | Octadecanoic acid, 5,9,13,17-tetramethyl-, methyl ester, [5R-(5R*,9R*,13R*)]- | C ₂₃ H ₄₆ O ₂ | 354 |
| 32 | 26.12 | 3.59 | 9,10-Anthracenedione, 1,8-dihydroxy-3-methyl- | C ₁₅ H ₁₀ O ₄ | 254 |
| 33 | 26.32 | 0.62 | (+)-9-Norartemisinin, 9-n-butyl- | C ₁₈ H ₂₈ O ₅ | 324 |
| 34 | 26.40 | 0.75 | 3.beta.-Fluoro-5-hydroxy-7-keto-17.beta.-acetyloxy-androstane | C ₂₁ H ₃₁ FO ₄ | 366 |
| 35 | 26.77 | 0.64 | Methyl 17-methyl-octadecanoate | C ₂₀ H ₄₀ O ₂ | 312 |
| | | 100. | | | |

P: peak; R.T: retention time; Comp.: compound; C.F: compound formula; M.W: molecular weight; M.S: Mass spectra.


chromosome aberrations CA, they stained with Giemsa according to²³.

For SCE, other slides stained with Giemsa according to the Fluorescence - plus giemsa (FPG), (Tawn and Holdsworth 1992)²⁴ which was modified by current study. Default method is described as following: Hoechst stain was prepared at a final concentration 0.5 mg / ml by dissolving 5 mg of dye powder in 10 ml of water distilled, when there were need it, a 5µg / ml was prepared by PBS as a solvent, keeping the solution away from light under the temperature (-18) C°. Slides Stained with Hoechst dye (5µg / ml) for 25 minutes, then washed with a solution of the BPS (BDH company) and left to dry, slides immersed in PBS solution in a suitable container and exposed to a source of ultraviolet radiation (UV light) a wavelength of (365) nm

for (2.5-2 hours) at a distance of (10) cm, washed with PBS, and left to dry, stained with Giemsa prepared (1ml dye: 9ml Gurr's buffer) for (1.5 -1) minutes. This method has been modified as follows:

Hoechst stain prepared in the same way at a final concentration of 5 µg/ ml. Slides immersed in a Hoechst stain, which dilute (1 dye: 2 PBS) and exposed to ultraviolet radiation (UV light) at a wavelength (365) nm for (2.5-2 hours) at a distance of (10) cm distance, slides washed with distilled water and left to dry, the slide either: Staining directly by Giemsa (which prepared above) for (1.5-1) a minute.

or immersed into PBS solution (PH= 6.8) and exposed to direct sun light for 25 minutes, staining by Giemsa (which prepared above) for (1.5-1) hours, left them to dry and

Figure 1: Chromatogram of methanolic extraction of *R. tinctorum*.Table 3: percentage of mitotic index, blast index, chromosomal aberrations, sister chromatid exchange, cell cycle progression and replication index in human lymphocyte cultures exposed to *R. tinctorum* crude extracts.

| Con. (mg/ml) | M.I | B.I | C.A/cell | SCE/cell | CCP | | | R.I |
|--------------|--------|--------|----------|----------|--------|--------|-------|--------|
| | | | | | M1 | M2 | M3 | |
| 100 | 2.6 | 37.5 | 0.6 * | 5 * | 22.35 | 49.05 | 28.6 | 2.0625 |
| 10 | 2.562 | 39.915 | 0.52 * | 4 | 21.35 | 50.05 | 28.6 | 2.074 |
| 1 | 2.552 | 49.2 | 0.38 * | 4.584 * | 24.975 | 39.275 | 35.75 | 2.11 |
| Ct- | 3.1687 | 40.577 | 0.16 | 4.167 | 46.65 | 40.02 | 13.33 | 1.668 |
| Ct+ | 1.01 * | 10.9* | 3.1 * | 12.24* | 60.5 | 33.25 | 6.25 | 1.4 |

Data shows Means; Con.: concentration (mg/ml); M.I: mitotic index, B.I: blast index, C.A: chromosomal aberrations, SCE: sister chromatid exchange which appear by first method of modifying staining, CCP: cell cycle progression, R.I: replication index, Ct-: control negative (without treatment), Ct+: control positive (Mitomycin c with 50 ng/ml final concentrations, *: ($P < 0.05$).

washed with a Sorensen solution (Na_2HPO_4 with 9.08 g of KH_2PO_4 in a liter of distilled water).

Slides were examined under light microscope. Chromosomal aberrations were scored from 50 metaphases. It was classified according to the International System for Human Cytogenetic Nomenclature (ISCN). The number of SCE's was scored from 50 cells under second metaphases. Mitotic index and blast index were determined by scoring 1000 cells from each donor. One hundred cells were scored for the determination of the cell cycle progression and replication index (RI)²⁵⁻²⁷.

Analysis of variance (ANOVA) and the least significant difference (LSD) were used for the statistical analysis of the results using SPSS, version 10, P-values at levels ($P \leq 0.05$).

RESULT AND DISCUSSION

Chemical analysis

Atomic Absorption Spectrometer

Atomic Absorption Spectrometry determined the percentage of Cu, Zn, Cd, Mn, K, Fe, Co, Ti and N. Their amounts showed in Table (1). There were no phosphorus. The level of total proteins was 5.8125.

GC-MS analysis

Thirty-five compounds identified in methanol extract of *R. tinctorum* using GC-MS analysis. The major three components present were 9,12-Octadecadienoic acid

(Z,Z)- (29.75 %); 9-Octadecenoic acid (Z)-, hexadecyl ester (26.1 %) and 1,3-Propanediol, 2-ethyl-2-(hydroxymethyl)- (10.09 %). Mass spectra from full scan analysis of components were showed in (Table 2). Figure (1) showed chromatogram of methanolic extraction of *R. tinctorum*.

Cytotoxicity on cancer cell lines

In AMN3 cell line, the lower concentration (50 $\mu\text{g/ml}$ -5 mg/ml), reduced cell viably at level ($P \leq 0.01$), while the higher concentrations (50-100) mg/ml induced it. IC₅₀ is 21 mg/ml; the best Inhibition rate is 93.693% at 50 $\mu\text{g/ml}$. (Figure 2 A). The results of Figure (2) showed reduction, ($P \leq 0.05$), in cell viability of AMGM cell lines treating with all concentrations of extract with IC₅₀ reached to 76.44 mg/ml. Inhibition rate ranged between (28.54- 66.1) %, (Figure 2 B).

The extraction was less effect in cell viability of Hep-2 cell line, the maximum Inhibition rat was 22.12 % at 10 mg/ml, ($P \leq 0.05$). All other concentrations were not significant compared with control. There was no IC₅₀ in this cell line, (Figure 2 C).

Cytotoxicity on human lymphocytes.

Two methods of staining were succeeding to appear chromosomes banding. The changing in mitotic index, blast index, cell cycle progression and replication index were not significant. But the extract induced chromosomal aberration and sister chromatid exchange. These




Figure 2: Cytotoxicity activity of *R. tinctorum* crude extract on (A): AMN3 cell line, 24 hr., IC₅₀= 21 mg/ml, (B): AMGM cell line, 48 hr., IC₅₀= 76.44 mg/ml, (C): Hep-2 cell line, 72 hr. *: (P < 0.05), **: (P < 0.01).

inductions in chromosomal aberration were structural. They were: addition, deletion and ring chromosome. The deletions were the most common abnormalities. There were more than two chromosomal aberrations in the same cell, (Table 3). These changing were less effect in treatment of Mitomycin C which induced CA and SCE

with reduction in MI, BI and R.I compared with *R. tinctorum* crude extracts and negative control.

Madder is used in medicine, as its components are reported to exhibit various pharmacological and biological activities, including anticancer. Current results found that main components of madder roots extract are: 9,12-

Octadecadienoic acid (Z,Z)- (29.75 %); 9-Octadecenoic acid (Z)-, hexadecyl ester (26.1 %) and 1,3-Propanediol, 2-ethyl-2-(hydroxymethyl)- (10.09 %), their common names are Linoleic; Oleic acid, hexadecyl ester and Etriol respectively. So, the anticancer activity of crud extract, which found in this study, may due to these compounds. Linoleic has good anticancer activity against mouse myeloma XS 63-5 cells²⁸, colon cancer cell- mediated by the CDK inhibitor²⁹, and Fights and prevention of cancer³⁰. Different linoleic acid isomers inhibited the growth of mammary, colon, colorectal, gastric prostate, and hepatoma cell lines, mediated induction the expression of apoptotic genes³¹. The anticancer activity of madder may be due to oleic acid components rather than to Linoleic. Lot of studies approved the inhibition in cell proliferation induced by oleic acid in different cancer cell lines. Oleic acid could suppress the over-expression of HER₂ (erbB-2), a well-characterized oncogene, invasive progression and metastasis in different human cancers. Oleic acid could play a role in intracellular calcium signalling pathways linked to the proliferation event. Concerning to cell death, oleic acid has been shown to stimulate apoptosis in carcinoma cells. The mechanisms of it could be due to an increase in intracellular ROS production or caspase 3 activities³². In addition, Oleic acid is responsible for the prevention of breast cancer, precipitated a controversy³³. Many anti-cancer plants have similar components, in current results there are: Tetradecanoic acid and hexadecyl ester in small quantities, they may be responsible for anticancer of *R. tinctorum*. Tetradecanoic acid was found in *Microcosmus exasperates* which has good anticancer activity against Dalton's Lymphoma Ascites (DLA) cells³⁴. hexadecyl ester found in anticancer plant *Andrographis paniculata* leaf extract³⁵.

CONCLUSIONS

There were some elements in crud extract: Cu, Zn, Cd, Mn, K, Fe, Co, Ti and N but Phosphorus was absent. Thirty-five compounds identified extract using (GC-MS). The major three components present were 9,12-Octadecadienoic acid (Z,Z)- ; 9-Octadecenoic acid (Z)-, hexadecyl ester and 1,3-Propanediol, 2-ethyl-2-(hydroxymethyl)-. The results of cytotoxicity of crud extract on cancer cell showed reduction in cell viability of (AMGM) cell line treating with all concentrations of extract. In (AMN3) cell line, the lower concentrations, reduced cell viably while the higher concentrations (50-100) mg/ml induced it. The extraction was less effect in cell viability of (HEp-2) cell Line. The toxicity human lymphocytes showed important changing in mitotic index, blast index, cell cycle progression and replication index but the extract induced chromosomal aberration and sister chromatid exchange. More study about toxicity of major three components that present in this plant will be benefit.

ACKNOWLEDGMENTS

The author gratitude Prof. Dr. Nahi Yousef Yaseen and Assist. Prof. Ahmed Majeed Al-Shammari, The Iraqi Centre For Cancer and Medical Genetic Research from

AL-Mustansiriyah University, Baghdad/ Iraq for their supports and for providing the necessary facilities. The authors would, also, like to thank Al-Mustansiriyah University (www.uomustansiriyah.edu.iq) Baghdad Iraq for its support in the present work.

REFERENCES

1. Khare, C.P. 2007. Indian medicinal plants, An illustrated dictionary. Siproinger 836p.
2. Boldizs'ar, I.; Sz'ucs, Z.; F'uzfai, Zs. and Moln'ar-Perl, I. (2006). Identification and quantification of the constituents of madder root by gas chromatography and high-performance liquid chromatography. Journal of Chromatography A, 1133: 259–274.
3. Ba'nyai, P.; Kuzovkina, I. N.; Kursinszki, L. and Sz'oke, E'. (2006a). HPLC Analysis of Alizarin and Purpurin Produced by *Rubia tinctorum* L. Hairy Root Cultures. Chromatographia Supplement. 63: S111-S114.
4. Nartop, N.; Akay, S. and G'urel, A. (2013). Immobilization of *Rubia tinctorum* L. suspension cultures and its effects on alizarin and purpurin accumulation and biomass production. Plant Cell Tiss. Organ Cult (2013) 112:123–128. DOI 10.1007/s11240-012-0212-z.
5. Ba'nyai, P.; Kuzovkina, I. N.; Kursinszki, L. and Szoke, E. (2006b). HPLC analysis of alizarin and purpurin produced by *Rubia tinctorum* L. Hairy root cultures. Chromatographia Supplement. 63: 66–72.
6. Remington, J. P. and Horatio, C. (1918). Woods and others. The Dispensatory of the United States of America, - Botanicals Only- The Southwest School of Botanical Medicine, 12^{Ed}, 1586 p.
7. Kalyoncu, F.; Cetin, B. and Saglam, H. (2006). Antimicrobial Activity of Common Madder (*Rubia tinctorum* L. *Phytother. Res.* 20, 490–492.
8. Aboud, A. S. (2010). HPLC Analysis of *Rubia tinctorum* and its effect of Methanol aqueous extract on bacteria isolated from burns infection. Journal of Al-Nahrain University. (4): 166-175.
9. Karim, A.; Mekhfi, H.; Ziyayat, A.; Legssyer, A.; Bnouham, M.; Amrani, S.; Atmani, F.; Melhaoui, A. and Aziz, M. (2010). Anti-diarrhoeal activity of crude aqueous extract of *Rubia tinctorum* L. roots in rodents. J Smooth Muscle Res., 46(2):119-23.
10. Kawasaki, Y.; Goda, Y.; and Yoshihira, K. (1992). The Mutagenic Constituents of *Rubia tinctorum*. Chemical and Pharmaceutical Bulletin. 40(6): 1504-1509.
11. Mouri, C. and Laursen, R. (2012). Identification of anthraquinone markers for distinguishing *Rubia* species in madder-dyed textiles by HPLC. Microchim Acta, 179:105–113.
12. Inoue, K.; Shibutani, M.; Masutomi, N.; Toyoda, K.; Takagi, H.; Uneyama, C.; Nishikawa, A. and Hirose, M. (2008). A 13-week subchronic toxicity study of madder color in F344 rats. Food and Chemical Toxicology 46: 241–252.
13. Maier, M.S.; G'omez, B.A.; Parera, S.D.; Elkin, D.; Rosa, H. De, Ciarlo, N.C. and Svoboda, H. (2010). Characterization of cultural remains associated to a

- human skeleton found at the site HMS Swift (1770). *Journal of Molecular Structure*, 978: 191–194.
14. Ino, N.; Tanaka, T.; Okumura, A.; Morishita, Y.; Makita, H.; Kato, Y.; Nakamura, M. and Mori, H. (1995). Acute and subacute toxicity tests of madder root, natural colorant extracted from madder (*Rubia tinctorum*), in (C57BL/6 X C3H)1 mice. *Toxicol Ind. Health*. 11(4):449-58.
 15. Inoue, K.; Yoshida, M.; Takahashi, M.; Fujimoto, H.; Shibutani, M.; Hirose, M. and Nishikawa, A. (2009). Carcinogenic potential of alizarin and rubiadin, components of madder color, in a rat medium-term multi-organ bioassay. *Cancer Sci.*, 100(12): 2261–2267.
 16. Lajkó, E.; Bányai, P.; Zámbo, Z.; Kursinszki, L.; Szóke, É. and L. Kóhidai. (2015). Targeted tumor therapy by *Rubia tinctorum* L.: analytical characterization of hydroxyanthraquinones and investigation of their selective cytotoxic, adhesion and migration modulator effects on melanoma cell lines (A2058 and HT168-M1). *Cancer Cell International*. 15:119, 1-15.
 17. Harborn, J. B. (1984). *Phytochemical methods: A guide to modern techniques of plant analysis*. Chapman and Hall, New York. 1-4p.
 18. Kodama, K. (1963). *Methods of quantitative inorganic analysis*. John Wiley and Sons. New York, London.
 19. Egli, H. (2008). *Kjeldahl Guide*. BUCHI Labortechnik AG, CH-9230 Flawil, Switzerland 58p.
 20. Abdul Jalill R. DH. (2014). GC-MS analysis of *Calendula officinalis* and cytotoxic effects of its flower crude extract on human epidermoid larynx carcinoma (Hep-2). *World Journal of Pharmacy and Pharmaceutical Sciences*. 3(4): 237-275.
 21. Kamuhabwa, A.; Nshimo, C. and Witte, P. D. (2000). Cytotoxicity of some medicinal plant extracts used in Tanzanian traditional medicine. *J. Ethno-pharma.*, 70: 143-149.
 22. Gao, S.; Yu, B.P.; Li, Y.; Dong, W.G. and Luo, H.S. (2003). Antiproliferative effect of octreotide on gastric cancer cells mediated by inhibition of Akt/PKB and telomerase. *W. J. G.*, 9(10): 2362-2365.
 23. Verma, R.S.; Babu, A. 1989. *Human chromosomes manual of basic techniques*. Pergamon. Press., pp. 240.
 24. Tawn, E. J.; Holdsworth, D. (1992). Mutagen induced chromosome damage in human lymphocyte. 191p. In: Rooney, D.E. and Czepulkowski, B.H. *Human Cytogenetics*. Vol.II: Malignancy and Acquired Abnormalities. A practical approach. IRI Press at Oxford University Press, Oxford, New York and Tokyo.
 25. Stites, D. (1979). Laboratory methods of detection cellular immune function. In : *Basic and clinical immunology*, Fundberg, H. H; Stiter, D.; Coldwell, J. (ed), LAGE, Medical publication, California, pp. 318-322.
 26. Schnieder, E.; Lewiss, J. (1982). Comparison of *in vivo* and *in vitro* sister chromatid exchange induction. *Mutat. Res.*, (114): 183-187.
 27. King, M.; Wild, D.; Gocke, E. and Eckhardt, K. (1982). 5- Bromo. deoxy uridine tablets with improved depot effect for analysis *in vivo* of SCE, in bone marrow and spermatogonial cells. *Mut. Res*. 97: 7-9.
 28. Holley, R.W.; Baldwin, J.H. and Kiernan, J.A. (1974). Control of Growth of a Tumor Cell by Linoleic Acid. *Proc. Nat. Acad. Sci. USA*. 71(10): 3976-3978.
 29. Lim, D. Y.; Tyner, A. L.; Park, J.; Lee, J.; Choi, Y. H. and Park, J. H. Y. (2005). Inhibition of colon cancer cell proliferation by the dietary compound conjugated linoleic acid is mediated by the CDK inhibitor p21CIP1/WAF1. *Journal of Cellular Physiology*. 205(1): 107–113.
 30. Lee, K.W.; Lee, H.J.; Cho, H.Y. and Kim, Y.J. (2005). Role of the conjugated linoleic acid in the prevention of cancer. *Crit Rev Food Sci Nutr*. 45(2):135-44.
 31. Kelley, N.S.; Hubbard, N.E.; Erickson, K.L. (2007). Conjugated linoleic acid isomers and cancer. *J Nutr*. 137(12):2599-607.
 32. Carrillo, C.; Cavia, M. M. and Alonso-Torre, S. R. (2012). Antitumor effect of oleic acid; mechanisms of action: A review. *Nutr Hosp*. 27(6):1860-1865.
 33. Win, D. T. (2005). Oleic Acid – The Anti-Breast Cancer Component in Olive Oil. *AU J.T*. 9(2): 75-78.
 34. Meenakshi, V.K.; Senthamarai, S.; ParipooranaSelvi, M.; Gomathy, S.; Sankaravadivu, S. and Chamundeswari, K.P. (2013). *In vitro* and *in vivo* antitumor and immunomodulatory studies of *Microcosmus exasperatus* against DLA bearing mice. Chamundeswari1, *European Journal of Applied Engineering*. 2(3): 18-25.
 35. Wei, L. S.; Wendy Wee, Siong, J. Y. F. and Syamsumir, D. F. (2011). Characterization of antibacterial, antioxidant, anticancer properties and chemical composition of Malaysia *Adrographis pauculata* leaf extract. *Pharmacologyonline* 2: 996-1002.