

Study of Polyphenol Content and Anti- Oxidative Potential of *Tribulus terrestris* Dry Fruit Extract

Durgawale P P^{1*}, Datkhile K D²

Research Officer, Molecular and Genetic Laboratory, Krishna Institute of Medical Sciences Deemed University (KIMSUDU), Karad, Maharashtra, India.

Received: 2nd March, 17; Revised 1st May, 17, Accepted: 15th May, 17; Available Online: 25th May, 2017

ABSTARCT

The dry fruit extract of *Tribulus terrestris* has been used in traditional medicine for various therapeutic applications. This study was intended to investigate its anti- oxidant potential using *in-vitro* assays. Initially, the total polyphenolic, flavonoid, and tannin content of the ethanolic extract of its dry fruit were tested since these compounds confer majority of the reducing power to plant extracts. The anti- oxidant potential of the extract was investigated using radical scavenging assays such as DPPH and ABTS, superoxide radical scavenging assay, and ferric reducing anti-oxidant power assay. The polyphenol content was found to be 6.65 ± 0.64 (Mean \pm S.E.M.; n = 3) mg GAE/ g of extract, flavonoid content was 0.40 ± 0.02 (Mean \pm S.E.M.; n = 3) mg RE/ g of extract, and tannin content was 4.09 ± 0.67 (Mean \pm S.E.M.; n = 3) mg TAE/ g of extract. The results for anti- oxidant assay obtained were found to be comparable to ascorbic acid and butylated hydroxytoluene standards. The results indicate presence of polyphenol compounds which confer majority of anti- oxidant properties to plant extracts. Further *in-vitro* and *in-vivo* investigations would reveal the efficacy of the extract as a dietary source of anti- oxidants.

Keywords: *Tribulus terrestris*, polyphenols, radical scavenging.

INTRODUCTION

The cellular metabolic pathways are complex and well-regulated. Free radicals are continuously being produced in the human body as a result of the metabolic activities. Such free radicals are extremely damaging and interact immediately with biological molecules causing DNA strand modifications, lipid peroxidation, and protein structure modifications¹. The body's innate defense mechanism against these free radicals includes glutathione redox reaction, activation of enzymes such as superoxide dismutase, catalase, reductase among others. In certain disorders, the defense mechanism fails to keep up with the rate of free radical production and such excess free radicals prove deleterious causing cardiovascular disorders, diabetes, and also cancer¹⁻³.

The food we consume provides anti- oxidants in a form which can be readily utilized by the cells to neutralize the excess free radicals before they cause much damage⁴. Fruits and vegetables contain polyphenolic compounds mostly as their secondary metabolites⁴⁻⁷. Various reports state that there is a direct correlation in the amount of polyphenolic compounds in plants and their measured anti-oxidant activities. As such, polyphenolic compounds contribute majorly to the exhibited anti- oxidant activity. Including more polyphenol- rich food in our diet is thus the most accessible way to provide anti- oxidants to the body to protect it from the ill- effects of free radicals^{8,9}.

The traditional systems of medicine have relied on the use of a large number of plant species for their medicinal

properties^{10- 13}. Recent studies indicate that these plant sources are, at the same time, rich sources of anti-oxidants^{14- 18}. One of such medicinal plants is *Tribulus terrestris* which is an annual plant with thorny fruits belonging to Zygophyllaceae family growing mostly in warm conditions. Traditionally, the plant is used as an aphrodisiac, as a tonic, analgesic, diuretic and for treatment of urinary tract infections¹⁹. In the present study, we attempted to investigate the polyphenolic content of the ethanolic extract of the dry fruit of *T. terrestris* and study its anti- oxidant potential using *in-vitro* techniques.

MATERIALS AND METHODS

The dry fruits of *Tribulus terrestris* were purchased from a local Ayurvedic medicine shop. The required reagents such as sodium carbonate, sodium hydroxide, ethanol, L- ascorbic acid (ascorbic acid), butylated hydroxytoluene (BHT), 2, 2- diphenyl-1- Picrylhydrazyl (DPPH), 2, 2'-Azino- bis (3- ethylbenzothiazoline-6- sulfonic acid) Diammonium salt (ABTS), potassium persulfate, potassium dihydrogen phosphate, di- potassium hydrogen phosphate, potassium fericyanide, trichloroacetic acid, ferric chloride, nitro blue tetrazolium chloride (NBT), β - nicotinamide adenine dinucleotide phosphate reduced tetrasodium salt (β - NADPH), Tris(hydromethyl)amonimethane (Tris) were purchased from Sisco Research Laboratories Pvt. Ltd. Shimadzu spectrophotometer (UV- 1800) present in the Molecular and Genetic Laboratory was used for all spectrometric

*Author for Correspondence: pratikpdurgawale@gmail.com

measurements. Phenazine methosulfate (PMS) was purchased from MP Biologics. Folin-Ciocalteu reagent, rutin, tannic acid, sodium nitrite and aluminium chloride were provided by the Department of Pharmacology; while gallic acid was gifted by Department of Biochemistry.

Extraction of plant materials

The dried fruits of *T. terrestris* were powdered using mortar pestle. 10 gram (g) of the powdered fruit was extracted in 200 millilitres (ml) of ethanol using a Soxhlet apparatus. After four extraction cycles, the powdered fruit appeared colorless and the extract was collected. The dried extract was stored in -20 degree Celsius (° C) freezer. During experiments, the extract was re-suspended in ethanol and utilized.

Total phenolic content estimation

Gallic acid was used as the standard for determining the total phenolic content using the Folin-Ciocalteu method. The method was modified as follows²⁰. 12.5 microlitres (µl) of standard solution in ethanol of varying concentration was added to 900 µl water with 60 µl Folin-Ciocalteu reagent and vortexed. After 5 minutes, 170 µl of 7.5 % sodium carbonate solution was added, mixed by vortexing and incubated at room temperature for 90 minutes. Absorbance was measured using a spectrophotometer at 750 nanometer (nm) wavelength. The blank tube contained 12.5 µl of ethanol. The standard curve was constructed using gallic acid concentrations from 40 microgram/ ml (µg/ ml) to 200 µg/ ml in ethanol. The extract solution was diluted in ethanol to obtain absorbance within the standard curve values. Each solution was prepared in triplicate in three separate experimental repeats. The total phenolic content was calculated as mg gallic acid equivalent per gram of extract (mg GAE/ g of extract).

Estimation of tannin content

For the estimation of amount of tannins in the extract, tannic acid was used as the standard for the Folin-Ciocalteu method²⁰. The tannic acid dilutions of 20 µg/ ml to 100 µg/ ml were prepared in ethanol and 10 µl of each solution was added to 740 µl water. 50 µl Folin-Ciocalteu reagent, 200 µl of 17.5 % sodium carbonate solution was added, vortexed, and incubated at room temperature for 30 minutes after which absorbance was measured at 725 nm. The blank tube contained 10 µl ethanol instead of standard solution, followed by similar additions as the standard dilutions as mentioned. The standard curve was plotted with concentration of the standard against their absorbance. The extract solution was diluted in ethanol accordingly and the tannin content was estimated as mg of tannic acid equivalent/ g of extract (mg TAE/ g of extract). Each solution was prepared in triplicate for three separate experimental repeats.

Estimation of flavonoid content

The aluminum chloride assay was used for determining the quantity of flavonoids in the extract using rutin as the standard²⁰. Briefly, rutin dilutions ranging from 20 µg/ ml to 100 µg/ ml were prepared in ethanol. 100 µl of each dilution was added to a microfuge tube containing 400 µl water and 30 µl of 5 % sodium nitrite. After 5 minutes, 30 µl of 10 % aluminium chloride was added and mixed by

vortexing. 200 µl of 1 molar (M) sodium hydroxide and 240 µl water was added, mixed and absorbance was measured at 510 nm using spectrophotometer. The blank tube contained 100 µl ethanol instead of standard solution. The extract solution was diluted in ethanol appropriately and flavonoid content was estimated from the standard curve plotted using concentration and absorbance values of rutin. The flavonoid content was represented as mg of rutin equivalent/ g of extract (mg of RE/ g of extract). Each solution was prepared in triplicate for three independent experimental repeats.

DPPH radical scavenging assay

The DPPH free radical scavenging assay was performed to study the free radical scavenging activity of the extract²¹. As standards, ascorbic acid (range 5 µM to 30 µM) and BHT (range 10 to 500 µM) dilutions were prepared in ethanol in a final volume of 250 µl. The extract dilutions from 0.1 µg/ ml to 40 µg/ ml were prepared in ethanol in a final volume of 250 µl. To each tube 750 µl of 200 µM DPPH solution in ethanol was added, mixed by vortexing and incubated at room temperature in the dark for 30 minutes following which absorbance was measured at 517 nm. The control tube contained 250 µl of ethanol instead of the samples and treated the same way as the standards and samples. Each solution was prepared in triplicate for two independent experimental repeats. The percentage inhibition or scavenging (% scavenging) of the DPPH radical was calculated as follows:

$$\% \text{ Inhibition} = \frac{[(\text{Absorbance of control} - \text{Absorbance of sample}) / (\text{absorbance of control})] \times 100}{1}$$

ABTS radical scavenging assay

The ABTS radical scavenging assay was performed according to earlier published methods with some modifications²². Ascorbic acid (dilution range 2 – 32 µg/ ml) and BHT (dilution range 5 – 80 µg/ ml) solutions in ethanol were used as standards. Extract dilutions (dilution range 4- 40 mg/ ml) were prepared in ethanol in a final volume of 0.2 ml. For production of ABTS^{•+} radical, 7 mM ABTS solution with 2.4 mM potassium persulfate was incubated at room temperature in dark for 16 hours. The absorbance of the ABTS^{•+} solution at 734 nm was adjusted to 0.7 and 1 ml of this solution was added to 0.2 ml of standard and sample dilutions in ethanol. The solution was mixed by vortexing and incubated at room temperature for 30 minutes, following which absorbance was measured at 734 nm. The control tube contained 0.2 ml ethanol instead of sample or standard and was treated in a similar way. Each solution was prepared in triplicate for two independent experimental repeats. The percentage inhibition or scavenging (% scavenging) of the ABTS^{•+} radical was calculated as follows:

$$\% \text{ Inhibition} = \frac{[(\text{Absorbance of control} - \text{Absorbance of sample}) / (\text{absorbance of control})] \times 100}{1}$$

Superoxide radical scavenging assay

For the generation of superoxide free radicals, the NADPH- PMS method was used for the reduction of NBT²³. Ascorbic acid (dilution range 20 – 100 µg/ ml) and BHT (dilution range 0.1 – 0.5 mg/ ml) solutions in ethanol were used as standards. Extract dilutions (dilution range




Figure 1: Calibration curve for determination of total phenolic content, tannin content, and flavonoid content. (a) Graph of gallic acid concentration ($\mu\text{g/ml}$) against the measured absorbance at 550 nm. (b) Graph of tannic acid concentration ($\mu\text{g/ml}$) against the measured absorbance at 725 nm. (c) Graph of rutin concentration ($\mu\text{g/ml}$) against the measured absorbance at 510 nm.

0.5 - 5 mg/ml) were prepared in ethanol in a final volume of 0.1 ml. Briefly, 0.1 ml of 1.13 mM NADPH, 0.25 ml of 500 μM NBT, 0.35 ml Tris-HCl (pH 8.0) were added to a tube containing 0.1 ml of standard or sample dilutions. The solution was mixed by vortexing and the reaction was initiated by adding 0.2 ml of 40.3 μM PMS. The tube contents were mixed and incubated at room temperature for 5 minutes following which absorbance was measured at 560 nm. The control sample contained 0.1 ml ethanol instead of sample or standard. Each solution was prepared in triplicate for two independent experimental repeats. The superoxide radical scavenging activity was expressed as percentage inhibition as follows:

$$\% \text{ Inhibition} = \frac{[\text{Absorbance of control} - \text{Absorbance of sample}]}{(\text{absorbance of control})} \times 100$$

Ferric reducing antioxidant power

To investigate the total reducing power of the extract, the ferric reducing antioxidant power assay was performed²⁴. Ascorbic acid (dilution range 2 - 32 $\mu\text{g/ml}$) and BHT (dilution range 5 - 80 $\mu\text{g/ml}$) solutions in ethanol were used as standards. Extract dilutions (dilution range 1 - 20 mg/ml) were prepared in ethanol in a final volume of 0.1 ml. 0.25 ml of phosphate buffer (pH 6.6), 0.25 ml of 1% potassium ferricyanide were added to a tube containing 0.1 ml of standard or sample in ethanol. The mixture was incubated at 50 $^{\circ}\text{C}$ for 20 minutes and after cooling 0.25 ml of 10% trichloroacetic acid was added. The samples were then centrifuged at 5000 rpm for 10 minutes and the upper layer was added to a separate tube containing 0.25 ml water

and 0.05 ml 0.1% ferric chloride. The mixture was incubated at room temperature for 10 minutes following which absorbance was measured at 734 nm. The control tube contained 0.1 ml ethanol instead of standard or sample. The reducing power was estimated by plotting the concentration of samples against the measured absorbance at 734 nm. Each solution was prepared in triplicate for two independent experimental repeats.

RESULTS AND DISCUSSION

The polyphenolic compounds found in plants are mostly secondary metabolites which confer certain characteristics to the plant. When such polyphenolics containing food is consumed, these compounds have been proven to be beneficial in many ways including protection against oxidative stress by scavenging the harmful free radicals produced in the body or by inducing expression of factors in the cell which help in abating the oxidative damage caused by such free radicals⁵. The content of polyphenolics in plant extract samples has been positively co-related to its observed anti-oxidant activity measured *in-vitro*^{7, 14-18}. As such, studying the content of polyphenolic compounds in a given plant extract sample can prove to be informative while studying its anti-oxidant activities. The term polyphenol compound encompasses large groups of compounds such as flavanoids, tannins, anthocyanins, catechins, lignans, etc. having diverse structures⁵. Flavanols and flavones found in pomegranate have anti-cancer properties²⁵⁻²⁷. Tannins and flavonoids have anti-




Figure 2: Graph of anti-oxidant activity of extract against the respective concentration studied using different anti-oxidant assays such as (a) DPPH radical scavenging assay; (b) ABTS radical scavenging assay; (c) Superoxide radical scavenging assay; (d) Ferric reducing anti-oxidant power assay.

oxidant, anti-inflammatory, anti-fungal and healing properties²⁸. Catechins found in tea have cytoprotective activity against oxidative stress induced toxicity and are known to have anti-angiogenic and anti-tumour properties²⁹.

The total phenolic content analysis was performed using the Folin – Ciocalteu method. The Folin- Ciocalteu reagent is made up of sodium molybdate and sodium tungstate which form complex with polyphenols in alkaline conditions. Higher the amount of reacting polyphenolic groups under given conditions higher is the amount of complex formed and higher the absorbance. For measuring the total phenolic content, gallic acid was used as the standard to plot the calibration curve and the phenolic content of the extract was plotted accordingly. The obtained total phenolic content was expressed as gallic acid equivalent per gram dry weight of the extract. The total phenolic content of the extract was determined to be 6.65 ± 0.64 (Mean \pm S.E.M.; $n = 3$) mg GAE/ g of extract Fig. 1-a. The tannin content of the extract was also determined using the Folin- Ciocalteu method under slightly modified conditions and tannic acid was used to plot the standard curve. The calculated tannin content of the extract was expressed as tannic acid equivalent per gram dry weight of the extract. The tannin content was found to be 4.09 ± 0.67 (Mean \pm S.E.M.; $n = 3$) mg TAE/ g of extract Fig. 1-b.

The flavonoid content of the extract was determined using the aluminum chloride method. Varying concentrations of rutin were used to plot the calibration curve and the

flavonoid content of extract was calculated as mg rutin equivalent per g dry weight of the extract. The flavonoid content was found to be 0.40 ± 0.02 (Mean \pm S.E.M.; $n = 3$) mg RE/ g of extract Fig. 1-c.

After studying the polyphenolic contents, the anti-oxidant potential of the extract was studied using various free radical scavenging assays. One of the methods employed the use of stable free radical, DPPH, with maximum absorbance at 517 nm. The polyphenolic compounds in the extract neutralize the stable free radical DPPH resulting in discoloration of the solution proportional to the amount of radicals neutralized²¹. Ascorbic acid and BHT were used as the standards to compare the anti-oxidant potential of the extract. Ascorbic acid exhibited 90 % inhibition at 30 µM; while BHT inhibited 88 % of the DPPH radical at 350 µM concentration. The extract solution showed increasing inhibition corresponding to increase in its concentration with maximum inhibition of 84.6 % at 30 µg/ ml concentration Fig. 2- a.

The ABTS radical scavenging assay relies on the production of $ABTS^{\cdot+}$ cation by reacting ABTS with potassium persulfate resulting in a blue/ green solution. The subsequent reduction of the radical $ABTS^{\cdot+}$ cation by anti-oxidants results in discoloration of the solution which can be measured as decrease in absorbance at 734 nm²². Ascorbic acid and BHT solutions exhibited complete inhibition at 32 µg/ ml and 80 µg/ ml, respectively. The extract solution was observed to have a proportional increase in inhibition with increasing concentration, although at higher concentration compared to standards,

with complete inhibition at 40 mg/ ml concentration ^{Fig. 2- b}.

In the superoxide radical scavenging assay, superoxide radical produced by using the NADPH/ PMS system reduces the tetrazolium salt NBT to insoluble blue colored formazan crystals. The anti-oxidants prevent the reduction of NBT by scavenging the superoxide radicals and thus resulting in decrease in absorbance measured at 560 nm²³. Ascorbic acid and BHT used as standards showed 30.9 % inhibition at 100 µg/ ml and 12.5 % at 500 µg/ ml, respectively. The extract solution showed increasing scavenging activity with corresponding increase in concentration from 6.6 % inhibition up to a maximum of 15.3 % inhibition at 5 mg/ ml ^{Fig. 2- c}. It was observed that above 5 mg/ ml concentrations of the extract, the solution turned turbid which interfered with the absorbance measurements.

The ferric reducing antioxidant assay is an indirect measure of the samples' anti-oxidant potential indicated by reduction of ferrous (Fe³⁺) ions in the solution to ferric (Fe²⁺) ions which can be measured by a change in absorbance²⁴. Ascorbic acid and BHT standard solutions exhibited a proportional linear increase in absorbance with increasing concentration across the entire concentration range. Similarly, the sample also showed increased absorbance with corresponding increase in concentration from 5 mg/ml to 40 mg/ ml. Extract solutions with concentration greater than 40 mg/ ml were turbid and hence could not be tested ^{Fig. 2- d}.

CONCLUSION

Based on the results obtained from the above described experiments, it can be seen that the ethanolic extract of the dry fruit of *T. terrestris* contains measurable polyphenolics comprising of flavonoids, and tannins among others. As mentioned, there is a positive relation between the proportion of polyphenolic content of plant extracts and its anti- oxidant potential^{7, 14- 18}. This anti- oxidant potential was successfully studied using the above methods which indicate that the extract has anti- oxidant potential comparable to the standards used. Further *in-vitro* and *in-vivo* experiments with a detailed phytochemical screening might shed light on the use of this extract as a source of anti- oxidants for human use.

ACKNOWLEDGEMENT

The authors would like to extend their gratitude towards the Honorable Vice Chancellor Dr. Mrs. Neelima Malik and the Directorate of Research, KIMSDU for supporting and guiding this project. The authors would also like to thank Mr. Anup Hendre, Department of Biochemistry, KIMSDU and Mrs. Trupti Durgawale, Department of Pharmacology, KIMSDU for providing reagents as gift samples.

REFERENCES

- Halliwell B. Free radicals, antioxidants, and human disease: curiosity, cause, or consequence? *The Lancet* 1994; 344:8924.
- Halliwell B. Free radicals and antioxidants: a personal view. *Nutrition reviews* 1994; 52(8): 253-265.
- Lawrence MJ and Bendich A. Free radical tissue damage: protective role of antioxidant nutrients. *The FASEB Journal* 1987; 1(6): 441-445.
- Halliwell B. Free radicals and antioxidants in food and in vivo: what they do and how they work. *Critical Reviews in Food Science & Nutrition* 1995; 35(1-2): 7-20.
- Pandey KB and Rizvi SI. Plant polyphenols as dietary antioxidants in human health and disease. *Oxidative medicine and cellular longevity* 2009; 2(5): 270-278.
- Devasagayam TPA. Free radicals and antioxidants in human health: current status and future prospects. *Japi* 2004;52(10): 794-804.
- Pulido R, Bravo L, Saura-Calixto F. Antioxidant activity of dietary polyphenols as determined by a modified ferric reducing/antioxidant power assay. *Journal of agricultural and food chemistry* 2000;48(8): 3396-3402.
- Bravo L. Polyphenols: chemistry, dietary sources, metabolism, and nutritional significance. *Nutrition reviews* 1998;56(11): 317-333.
- Scalbert A and Williamson G. Dietary intake and bioavailability of polyphenols. *The Journal of nutrition* 2000;130(8): 2073-2085.
- Iqbal A, Zafar M, Faiz M. Screening of some Indian medicinal plants for their antimicrobial properties. *Journal of ethnopharmacology* 1998;62(2): 183-193.
- Fabricant DS and Farnsworth NR. The value of plants used in traditional medicine for drug discovery. *Environmental health perspectives* 2001;109 (1): 69.
- Tahraoui A. Ethnopharmacological survey of plants used in the traditional treatment of hypertension and diabetes in south-eastern Morocco (Errachidia province). *Journal of ethnopharmacology* 2007;110(1): 105-117.
- Lin J. Preliminary screening of some traditional Zulu medicinal plants for anti-inflammatory and anti-microbial activities. *Journal of ethnopharmacology* 1999;68(1): 267-274.
- Cai, Yizhong. Antioxidant activity and phenolic compounds of 112 traditional Chinese medicinal plants associated with anticancer. *Life sciences* 2004; 74(17): 2157-2184.
- Auddy B. Screening of antioxidant activity of three Indian medicinal plants, traditionally used for the management of neurodegenerative diseases. *Journal of Ethnopharmacology* 2003;84 (2): 131-138.
- Gacche RN, Dhole A. Antioxidant and possible anti-inflammatory potential of selected medicinal plants prescribed in the Indian traditional system of medicine. *Pharmaceutical biology* 2006;44(5): 389-395.
- López-Alarcón C, Denicola A. Evaluating the antioxidant capacity of natural products: A review on chemical and cellular-based assays. *Analytica chimica acta* 2013;763: 1-10.
- Katalinic V. Screening of 70 medicinal plant extracts for antioxidant capacity and total phenols. *Food chemistry* 2006; 94(4): 550-557.

19. Ody P. The Complete Guide Medicinal Herbal, Dorling Kindersley 2000.
20. Singleton VL, Orthofer R, Lamuela-Raventós RM. Analysis of total phenols and other oxidation substrates and antioxidants by means of folin-ciocalteu reagent. *Methods in enzymology* 1999;299: 152-178.
21. Mensor LL. Screening of Brazilian plant extracts for antioxidant activity by the use of DPPH free radical method. *Phytotherapy research* 2001;15(2): 127-130.
22. Roberta R. Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free radical biology and medicine* 1999;26(9): 1231-1237.
23. Okamoto, Genichi, Fumitaka H, Kato H. Scavenging of active oxygen species by glycated proteins. *Bioscience, biotechnology, and biochemistry* 1992;56(6): 928-931.
24. Gülçin İ. Radical scavenging and antioxidant activity of tannic acid. *Arabian Journal of Chemistry* 2010;3(1): 43-53.
25. Kowalski J. Effect of apigenin, kaempferol and resveratrol on the expression of interleukin-1beta and tumor necrosis factor-alpha genes in J774. 2 macrophages. *Pharmacological reports: PR* 2004;57(3): 390-394.
26. Mano H. Luteolin induces apoptosis via death receptor 5 upregulation in human malignant tumor cells. *Oncogene* 2005;24(48): 7180-7189.
27. Lansky EP, Newman RA. *Punica granatum* (pomegranate) and its potential for prevention and treatment of inflammation and cancer. *Journal of ethnopharmacology* 2007;109(2): 177-206.
28. Araújo DS, Antônio T. A new approach to study medicinal plants with tannins and flavonoids contents from the local knowledge. *Journal of Ethnopharmacology* 2008;120(1): 72-80.
29. Singh BN, Sharmila S, Srivastava RK. Green tea catechin, epigallocatechin-3-gallate (EGCG): mechanisms, perspectives and clinical applications. *Biochemical pharmacology* 2011;82(12): 1807-1821.