

In Vitro Evaluation of Antioxidant and Antimicrobial Activity of Carrot Peel

Sheila John^{1*}, Priyadarshini S¹, Sarah Jane Monica¹, Sivaraj C², Arumugam P²

¹Department of Home Science, Women's Christian College, Chennai – 600 006.

²Armats Biotek Research and Training Institute Chennai – 600 032.

Received 29th May, 17; Revised 27th June, 17, Accepted 13th July, 17; Available Online 25th July, 2017

ABSTRACT

The objectives of the study were to assess the total phenolic content and flavonoid content, and to investigate the antioxidant and antimicrobial potential of carrot peel extracted with methanol and acetone. Total Phenolic content and Flavonoid content was higher in carrot peel acetone extract (AE) i.e. 106.36 µg/mg of GAE and 27.83 µg/mg of QE respectively than methanol extract (ME). Carrot peel acetone extract showed greater antioxidant capacity for DPPH radical scavenging activity whereas methanol extract showed better result for FRAP reducing power and Phosphomolybdenum assay. Both AE and ME could inhibit the growth of *Shigella flexneri*, *E coli*, *Staphylococcus aureus* and *Klebsiella pneumonia*. This study substantiates the potential of carrot peel to be used as a novel functional food ingredient, natural antioxidant and antimicrobial agent.

Keywords: antimicrobial potential, carrot peel, Flavonoid, Phenolic content.

INTRODUCTION

Vegetables and some fruits yield between 25% and 30% of nonedible products, which include skins and seeds that normally have no further usage and are commonly wasted or discarded^{1,2}. The byproducts of plant food processing represent a major disposal problem for the industry concerned, but they are also promising sources of compounds which may be used because of their favourable technological or nutritional properties³. The by-products may still contain many valuable substances, such as, pigments, sugars, organic acids, flavours and bioactive compounds with antioxidant and antimicrobial activities, as well as being valuable sources of dietetic fibre⁴. Growing body of research evidence proves that the content of phytochemical compounds is higher in peel and seeds with respect to the edible tissue of fruits and vegetables. For eg. the total phenolic compounds in the peels of lemons, oranges, and grapefruits were 15% higher than that of the pulp of these fruits⁵.

Carrot (*Daucus carota*) is classified as vitaminized food as it is rich in β-carotene, ascorbic acid and tocopherol⁶. Carrot is also a significant source of phenolic compounds such as hydroxycinnamic acids and derivatives⁷, para-hydroxybenzoic acids⁸ and polyacetylenes^{9,10}. Due to appreciable level of variety of different compounds present, carrots are considered as a functional food with significant health promoting properties¹¹. Carrot processing residues e.g. peels, pomace, are usually discarded or used as animal feed which are high in phenolics and antioxidant properties, could be considered for value-added utilization. The studies on evaluating the phenol content of carrot revealed the peel had higher

content of phenolics than the flesh. Kähkönen et al.,¹² found that carrot peel and flesh contained 6.6 and 0.6 mg gallic acid equivalent/g dry weight respectively. Phenolic content in different tissues decreased from peel, phloem to xylem, similarly antioxidant and radical scavenging activities in different tissues decreased in same order as the phenolic content and correlated well with total phenolic contents⁷.

Although studies have been reported on the antioxidant and antimicrobial properties of the carrot peel, it should be noted that result may vary according to the geographical location, soil in which the plant is grown, season, varieties and method of extraction. Therefore the current study was undertaken to elucidate antioxidant and antimicrobial properties of the carrot peel so as to validate its use as nutraceutical and as food preservative.

MATERIALS AND METHODS

Preparation of carrot peel extracts

Healthy, disease free and mature carrots were purchased from the local market. Cleaned and washed carrots were manually peeled using a sterilized peeler and the peels were then shade dried at room temperature for 5 days. The shade dried carrot peels were powdered in a laboratory blender and was kept in airtight bottles until further use. The powdered carrot peel was soaked in methanol and acetone for 72 h by maceration technique. The supernatants were filtered through Whatman No.1 filter paper and concentrated using rotary evaporator. The dry residue was preserved at 5°C in airtight bottles until further use.

Estimation of total phenolic content

Figure 1: Carrot (*Daucus carota*) and powdered carrot peel.

Table 1: Total Phenolic and flavonoid content of carrot peel extract.

Extracts	Total content	Phenolic content	Total Flavonoid content
Methanol	19.94 µg/g of GAE	5.39 µg/g of QE	
Acetone	106.36 µg/g of GAE	27.83 µg/g of QE	

Table 2 a: DPPH activity of carrot peel acetone extract.

S.No	Concentration (µg/µL)	Acetone extract % of Inhibition
1	50	32.82
2	100	51.90
3	150	72.51
4	200	83.97
5	250	95.38
6	300	96.95

Table 2 b: DPPH activity of carrot peel acetone.

S.No	Concentration (µg/µL)	Methanol extract % of Inhibition
1	100	7.45
2	200	10.78
3	300	20.28
4	400	21.69
5	500	27.98
6	600	34.91

Table 3: FRAP reducing power assay of carrot peel acetone and methanol extract.

S.No	Concentration (µg/µL)	Absorbance at 700 nm	
		Acetone extract	Methanol extract
1	20	0.09	0.14
2	40	0.46	0.52
3	60	0.66	0.67
4	80	0.70	0.80
5	100	0.85	4.52
6	120	0.86	5.45

Total phenolic content of the carrot peel extracts (Methanol and Acetone) were estimated by the Folin-Ciocalteu reagent method¹³. Hundred µL of extracts (1 mg/mL) were made up to 1 mL by methanol and mixed with 1 mL of Folin Ciocalteu reagent (1:10 diluted with

distilled water) and after 5 min one mL of 20% sodium carbonate (Na_2CO_3) was added. The mixture was incubated at room temperature for 30 minutes and the absorbance was measured at 760 nm by spectrophotometer. The total phenolic content was expressed in terms of Gallic acid equivalent (µg/mg of extract), which is a common reference standard.

Estimation of total Flavonoid content

The total flavonoid content of the carrot peel AE and ME was determined by the Aluminium Chloride reagent method¹⁴. The extract (500 µg/mL) was made up to one mL by methanol and were mixed with 0.5 mL of 5% sodium nitrate (NaNO_2) solution and allowed to stand for 5 min. Then 0.3 mL of 10% Aluminium Chloride (AlCl_3) solution was added and the mixture was allowed to stand for further 5 min. Finally, one mL of 1 M Sodium hydroxide (NaOH) solution was added, and the final volume of the mixture was brought to 5 mL with distilled water. The mixture was incubated for 15 min at room temperature and absorbance was measured at 510 nm. The total flavonoid content was expressed as quercetin equivalent (µg/mg of extract), which is a common reference standard.

Antioxidant activity

DPPH (1,1-diphenyl-2-picryl-hydrazyl) radical scavenging assay

The DPPH radical scavenging activity was carried out according to the method described by Blois¹⁵, where one mL of the carrot peel extracts (ME and AE) were taken in various concentrations and mixed with one mL of 0.1 mM of DPPH solution in methanol. The setup was left at dark in room temperature and the absorption was monitored after 30 minutes. Absorbance was read at 517 nm in spectrophotometer. The ability of the test sample to scavenge DPPH radical was calculated by the following formula:

$$\frac{\text{Absorbance in control} - \text{Absorbance in sample}}{\text{Absorbance in control}} \times 100$$

Absorbance control was the absorbance of DPPH and methanol

Absorbance sample was the absorbance of DPPH radical and the test sample.

Ferric Reducing Antioxidant Power assay

The Fe^{3+} reducing power assay was done according to the method described by Yen and Chen¹⁶. Different

Table 4 a: Phosphomolybdenum assay of carrot peel acetone extract.

S.No	Concentration ($\mu\text{g}/\mu\text{L}$)	Absorbance at 695 nm
1	50	0.29
2	100	0.63
3	150	0.69
4	200	0.72
5	250	0.73
6	300	0.74

Table 4 b: Phosphomolybdenum assay of carrot peel methanol extract.

S.No	Concentration ($\mu\text{g}/\mu\text{L}$)	Absorbance at 695 nm
1	20	0.29
2	40	0.74
3	60	0.84
4	80	0.85
5	100	0.86
6	120	0.89

concentrations of the carrot peel AE and ME (each 10 mg) were mixed with one mL of phosphate buffer (0.2 M, pH 6.6) and one mL of potassium ferricyanide (1%). The mixture was incubated at 50°C for 20 min. Then one mL of 10% trichloroacetic acid was added to the mixture. Followed by addition of one mL of 0.1% of freshly prepared ferric chloride and the absorbance of the resultant solution was measured at 700 nm.

Phosphomolybdenum reduction assay

The antioxidant activity was evaluated by reduction assay method¹⁷ by the formation of green phosphomolybdenum complex. The carrot peel extracts ME and AE (each 1 mL) of various concentrations were combined with 1 mL of reagent solution (0.6 M sulphuric acid, 28 mM sodium phosphate and 4 mM ammonium molybdate). The tubes were capped and incubated in a water bath at 95°C for 90 min. The samples were cooled to room temperature and the absorbance of the mixture was measured at 695 nm against blank.

Antibacterial activity

The antibacterial activity of the carrot peel was analysed by well diffusion method¹⁸. Muller Hinton agar was prepared according to the standard procedure and 25 mL was poured into the plates and was allowed to solidify. The standard inoculum suspension was streaked over the surface of the media using sterile cotton swab to ensure the confluent growth of the organism and the plates were allowed to dry for 5 minutes. After drying, the different concentrations (50, 75 and 100 $\mu\text{g}/\text{mL}$) of the methanol and acetone extract of carrot peel were poured into the wells. Tetracycline was used as a standard (1 $\mu\text{g}/\text{mL}$) and methanol was used as control. Finally the inoculated plates were incubated for 24 hours at 37°C. The zone of inhibition was measured and noted.

RESULTS AND DISCUSSION

Total Phenolic and flavonoid content

The total phenolic and flavonoid content of carrot peel extracts (ME and AE) is presented in Table.1. Total Phenolic content and Flavonoid content was higher in carrot peel acetone extract (AE) i.e. 106.36 $\mu\text{g}/\text{g}$ of GAE and 27.83 $\mu\text{g}/\text{g}$ of QE respectively than methanol extract (ME). Phenolic content can be used as an important indicator of antioxidant capacity and can be used as a preliminary screen for any product when intended to be used as a natural source of antioxidants in functional foods¹⁹. Phenolic compounds are dietary antioxidants found in plants that are shown to inhibit LDL oxidation, inhibit platelet aggregation and adhesion, decrease total and LDL cholesterol and induce endothelium- dependent vaso-relaxation^{20,21,22}. Flavonoids are the major phenolic compounds in plants²³. They are considered as potential antioxidants exerting their antioxidant activity by the mechanisms of radical scavenging and metal ion chelation to inhibit lipid peroxidation²⁴.

Antioxidant activity

The antioxidant activity of the carrot peel was evaluated using three *in vitro* assays namely DPPH Assay, FRAP Assay, Phosphomolybdenum Assay.

DPPH Assay

DPPH assay is widely used to test the ability of compounds to act as free radical scavengers or hydrogen donors and to evaluate antioxidant activity²⁵. The free radical scavenging activity was high in acetone extract compared to methanol extract. The result of the DPPH assay of carrot peel is presented in Table 2 a and Table 2 b The AE showed highest activity (96.95 %) at 300 μL concentration while the highest activity for ME i.e. 34.91% was at 600 μL . The IC₅₀ carrot peel AE and ME are 76 $\mu\text{g}/\mu\text{L}$ and 859 $\mu\text{g}/\mu\text{L}$ respectively. The IC₅₀ value of ascorbic acid was 4.89 $\mu\text{g}/\mu\text{L}$.

FRAP Assay

The antioxidant capacity of the carrot peel extracts is determined by the ability of the antioxidants in these extracts to reduce ferric to ferrous iron in FRAP reagent. The ferric reducing power of the carrot peel AE and ME at various concentrations is presented in Table 3. From the Table 3, it is evident that the Methanol extract showed greater ferric reducing capacity than acetone extract.

Nguyen et al.,²⁶ studied the antioxidant activity of the carrot peel extracted with four solvents (methanol, ethanol, water and hexane). The ferric reducing antioxidant power of carrot peel was 15.31, 8.88, 4.82 and 0.31 mg TE/g dry weight for methanol, ethanol, water and hexane, respectively.

Phosphomolybdenum Assay

The results for phosphomolybdenum assay are presented in Table 4 a and Table 4.2. From the Table 4a and Table 4 b, it can be inferred that Methanol extract had greater activity (0.89) for phosphomolybdenum assay than acetone extract of carrot peel.

The peels of vegetable and fruit could be used as natural antioxidant to preserve foods. Addition of other peels with the carrot peel could enhance its antioxidant properties. In a study carried out by Singh & Immanuel²⁷, the potential

Table 5: Antimicrobial activity of carrot peel acetone and methanol extract.

S.NO	Extract	Organisms	Standard	Zone of Inhibition		
				50 μ L	75 μ L	100 μ L
1	Acetone	<i>Shigella flexneri</i>	25 mm	11 mm	12 mm	13 mm
2		<i>E coli</i>	22 mm	12 mm	14 mm	16 mm
3		<i>Staphylococcus aureus</i>	23 mm	12 mm	13 mm	16 mm
4		<i>Klebsiella pneumonia</i>	26 mm	-	13 mm	14 mm
1	Methanol	<i>Shigella flexneri</i>	27 mm	14 mm	15 mm	16 mm
2		<i>E coli</i>	25 mm	13 mm	14 mm	16 mm
3		<i>Staphylococcus aureus</i>	29 mm	12 mm	13 mm	17 mm
4		<i>Klebsiella pneumonia</i>	21 mm	12 mm	14 mm	15 mm

of pomegranate, lemon and orange peels as natural antioxidants in paneer was studied. Among the peel used Pomegranate showed greater antioxidant activity by preventing peroxide formation and extending shelf life, followed by lemon peel and orange peel.

Antimicrobial activity

Antimicrobial potential of carrot peel extracts against four pathogens were assessed in terms of zone of inhibition of bacterial growth. The results of the antimicrobial activity are presented in Table 5.

The antimicrobial activity of the extracts increased linearly with increase in concentration of the extracts (μ g/mL). The growth inhibition zone measured for acetone extracts for various bacteria ranged from 11-16 mm. The growth inhibition zone measured for methanol extracts were in the range 12-17 mm. The antibacterial potential was dose-dependent against *Shigella flexneri*, *E coli*, *Staphylococcus aureus* and *Klebsiella pneumonia*. Carrot peel inhibited the growth of four food borne disease causing pathogens. There was little difference in antimicrobial activity in both the extracts. In a study carried out by Al-Baarri²⁸ showed that addition of carrot extract with Hypthiocyanite enhanced its antimicrobial activity against *Staphylococcus aureus* and *E coli*.

CONCLUSION

This is a preliminary study to evaluate the antioxidant and antimicrobial potential of carrot peel. Further studies in carrot peel are warranted to validate its functional and physiological benefits. Increasing awareness among consumers about the side effects of synthetic compounds in food and medicine are prompting food industries, scientists and pharmacists to search for newer and natural ingredients for human consumption which are safe and health promoting.

REFERENCES

- Ajila CM, Bhat SG, Rao UJSP. Valuable components of raw and ripe peels from two Indian mango varieties. *Food Chem* 2007; 102(4):1006–11.
- Ajila CM, Aalami M, Leelavathi K, Rao UJSP. Mango peel powder: a potential source of antioxidant and dietary fiber in macaroni preparations. *Innov Food Sci Emerg* 2010; 11(1):219-24.
- Schieber A, Stintzing FC, Carle R. By-products of plant food processing as a source of functional compounds-recent developments. *Trends Food Sci Technol* 2001; 12(11):401-13.
- Fernández-López J, Sendra E, Sayas-Barberá E, Navarro C, Pérez-Alvarez JA. Physico-chemical and microbiological profiles of “salchichón” (Spanish dry-fermented sausage) enriched with orange fiber. *Meat Science* 2008; 80(2):410-417.
- Gorinstein S, Martin-Belloso O, Park YS, Haruenkit R, Lojek A, Ciz M, Caspi A, Libman I, Trakhtenberg S. Comparison of some biochemical characteristics of different citrus fruits. *Food Chem* 2001; 74(3):309-16.
- Hashimoto T, Nagayama T. Chemical composition of ready-to eat fresh carrot. *J Food Hyg Soc Japan* 2004; 39:324-328.
- Zhang D, Hamaazee Y. Phenolic compounds and their antioxidant properties in different tissues of carrots (*Daucus carota* L.). *Food Agric Environ* 2004; 2:95-101.
- Babic I, Amiot MJ, Ngugen-The C, Aubert S. Changes in phenolic content in fresh, ready-to-use and shredded carrots during storage. *J Food Sci* 1993; 58:351-356.
- Hansen SL, Purup S, Christensen LP. Bioactivity of falcariol and the influence of processing and storage on its content in carrots (*Daucus carota* L.). *J Sci Food Agric* 2003; 83:1010-1017.
- Kidmose U, Hansen SL, Christensen LP, Edelenbos M, Larsen M, Norback R. Effects of genotypes, root size, storage and processing on bioactive compounds in organically grown carrots (*Daucus carota* L.). *J Food Sci* 2004; 69:388–394.
- Hager TJ, Howard LR. Processing effects on carrot phytonutrients. *Hortic Sci* 2006; 41:74-79.
- Kahkonen MP, Hopia AI, Vuorela HJ, Rauha JP, Pihlaja K, Kujala TS, Heinonen M. Antioxidant activity of plant extracts containing phenolic compounds. *J Agric Food Chem* 1999; 47:3954-3962.
- Singleton VL, Orthofer R, Lamuela-Raventos RM. Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. *Methods Enzymol* 1999; 299:152-178.
- Chang CC, Yang MH, Wen HM, Chern JC. Estimation of total flavonoid content in propolis by two complementary colorimetric methods. *Journal of Food and Drug Analysis* 2002; 10:178-182.
- Blois MS. Antioxidant determinations by the use of a stable free radical. *Nature* 1958; 29:1199-1200.
- Yen GC, Chen HY. Antioxidant activity of various tea extracts in relation to their antimutagenicity. *J. Agr. Food Chem.* 1995; 43:27-32.

17. Prieto P, Pineda M, Aguilar M. Spectrophotometric quantitation of antioxidant capacity through the formation of a phosphomolybdenum complex: Specific application to the determination of vitamin E. *Analytical Biochemistry* 1999; 269:337-41.
18. Langfield RD, Scarano FJ, Heitzman ME, Kondo M, Hammond GB, Neto CC. Use of a modified microplate bioassay method to investigate antibacterial activity in the Peruvian medicinal plant *Peperomia galiodes*. *J. Ethnopharmacol* 2004; 94(2-3): 279-281.
19. Viuda-Martos M, Ruiz-Navajas Y, Fernández-López J, Sendra E, Sayas-Barberá E, Pérez-Álvarez JA. Antioxidant properties of pomegranate (*Punica granatum* L.) bagasses obtained as co-product in the juice extraction. *Food Research International* 2011; 44: 1217-1223.
20. Lagiou P, Samoli E, Lagiou A, Skalkidis Y, Katsouyanni K, Petridou E, et al. Flavonoid classes and risk of peripheral arterial occlusive disease: a case control study in Greece. *European Journal of Clinical Nutrition* 2006; 60:214-219.
21. Lapointe A, Couillard C, Lemieux S: Effects of dietary factors on oxidation of low-density lipoprotein particles. *J Nutritional Biochemistry* 2006, 17:645-658.
22. Vita JA. Polyphenols and cardiovascular disease: effects on endothelial and platelet function. *Am J Clin Nutr* 2005; 81:292S-297S.
23. Walter M, Marchesan E. Phenolic compounds and antioxidant activity of rice. *Braz. Arch. Biol. Technol.* 2011; 54:371-377.
24. Fernando, Soysa. Total phenolic, flavonoid contents, in-vitro antioxidant activities and hepatoprotective effect of aqueous leaf extract of *Atalantia ceylanica*. *BMC Complementary and Alternative Medicine* 2014; 14:395.
25. Kirtikar KR, Basu BD. *Indian medicinal plants*. International book distributors, Dehradun, 2006, 993-994.
26. Singh S, Immanuel G. Extraction of Antioxidants from Fruit Peels and its Utilization in Paneer. *J Food Process Technol* 2014; 5:349.
27. Nguyen VT, Scarlett CJ. Mass Proportion, Bioactive Compounds and Antioxidant Capacity of Carrot Peel as Affected by Various Solvents. *Technologies* 2016; 4:3-6.
28. Al-Baarri AN, Legowo AM, Hayakawa S, Ogawa M. Enhancement Antimicrobial Activity of Hypthiocyanite using Carrot Against *Staphylococcus aureus* and *Escherichia coli*. *Procedia Food Science*. 2015 ; 3:473 – 478.