

# Fluorescence Detection of Phytoconstituents in Hydroalcoholic Extract of *Ipomoea aquatica* Forssk. by Thin Layer Chromatographic Analysis

Sasikala M\*, Sundaraganapathy R

Faculty of Pharmacy, Karpagam University, Karpagam Academy of Higher Education, Coimbatore – 641 021, Tamil Nadu, India

Received 2<sup>nd</sup> June, 17; Revised 23<sup>rd</sup> June, 17, Accepted 14<sup>th</sup> July, 17; Available Online 25<sup>th</sup> July, 2017

## ABSTRACT

**Introduction:** The medicinal plant water spinach is applicable for many therapeutic activities like anti cancer, anti oxidant, etc. the plant is used for purification of metals and waters of different categories. The benefit of hydroalcoholic extract in this study is non toxic, preservative and high yield of extract from the herbal powder. Thin layer chromatographic analysis is one of the best analytical methods for the separation and identification of phytoconstituents from the crude extract. The spots separated on the stationary phase could be detected by using the non destructive fluorescence method. The detection will be done at shorter and longer wavelength in UV cabinet. **Aim and Objective:** The aim was to separate and identify the various active metabolites present in the crude extract of the herb of interest using  $R_f$  value by thin layer chromatographic analysis. **Method:** The thin layer chromatographic parameters were set by trial and error method. Ethyl acetate: Methanol: Water (10.0:1.35:1.0 v/v/v) for alkaloids, Ethyl acetate: Pyridine: Water (5.0:1.0:4.0 v/v/v) for glycosides, Toluene: Acetone: Formic acid (4.5:4.5:1.0 v/v/v) for tannins and Toluene: Ethyl acetate (9.3:0.7 v/v) for triterpenoids were the mobile phases for the individual constituents. The constituents were identified at 365nm. **Results and Discussion:** Nearly 24 various constituents of alkaloids, glycosides, flavonoids, triterpenoids and tannins were separated using respective mobile phases through  $R_f$  value. **Conclusion:** The mentioned TLC profile for the plant of interest will hopefully help the researchers in herbal field to develop more herbal medicines requiring the particular constituent or its extract.

**Keywords:** TLC,  $R_f$ , alkaloids, glycosides, flavonoids, tannins, triterpenoids and UV light.

## INTRODUCTION

Extraction methods are widely used to get phytoconstituents from the plant. This is the basic method to detect the compounds present in each and every plant. The application of this process for an important medicinal plant *Ipomoea aquatica* helps in revealing its phytochemicals. These are formed in the primary and secondary metabolism of the herb. The metabolic products are extracted into suitable solvents in which those are soluble. The solvents for extraction are available with various polarities. Mostly therapeutically active secondary metabolites of the plant will come into polar solvents like alcohol, water, etc. Thereby extraction is found as the most important basic method to recognize the phytochemical potential of a natural source<sup>1-6</sup>.

Phytochemical Analysis is devoted to the publication of original articles concerning the development, improvement, validation and/or extension of application of analytical methodology in the plant sciences. The spectrum of coverage is broad, encompassing methods and techniques relevant to the detection (including bio-screening), extraction, separation, purification, identification and quantification of compounds in plant biochemistry, plant cellular and molecular biology, plant biotechnology, the food sciences, agriculture and horticulture<sup>7-17</sup>.

Phytoconstituents of the herbals are important source for herbal drug development. Hence, the study on detection about those is playing majorly in the pharmaceutical, botanical, agricultural and other life sciences fields. These chemicals are divided into two categories based on its metabolism such as primary and secondary metabolites. Among this, the later one acts as ailments in treating many diseases in human and are known as traditional herbal medicine<sup>18-26</sup>.

*Ipomoea aquatica* belonging to the family Convolvulaceae is a semi-amphibious, equatorial plant ripened as a vegetable for its tender boughs and leaves. It is begun all over the tropical and subtropical domains of the planet, although it is not acknowledged where it was emerged<sup>27-38</sup>.

The medicinal plant water spinach is applicable for many therapeutic activities like anti cancer, anti oxidant, etc. the plant is used for purification of metals and waters of different categories. The benefit of hydroalcoholic extract in this study is non toxic, preservative and high yield of extract from the herbal powder. Thin layer chromatographic analysis is one of the best analytical methods for the separation and identification of phytoconstituents from the crude extract. The spots separated on the stationary phase could be detected by using the non destructive fluorescence method. The

Table 1: Plant details.

S. No.	Parameters	Subject
1.	Plant Name	Water Spinach
2.	Botanical Name	<i>Ipomoea aquatica</i> FORSSK.
3.	Family	Convolvulaceae
4.	Location	Parambikulam – Aliyar Riverine, Pollachi
5.	Part of the plant	Whole plant
6.	Authentication No.	BSI/SRC/5/23/2017/Tech./3 269
7.	Place of Authentication	BSI, Coimbatore-641003, Tamil Nadu, India

Table 2: Instruments used.

S. No.	Name of the Instrument	Model Name
1.	Precision Balance	Wensar
2.	Hot plate	Cintex
3.	Ultra Sonicator	Labman
4.	Electrical Water bath	Technico
5.	UV cabinet	CAMAG and Deep Vision

Table 3: Chemicals/Reagents used.

S.No.	Name of the Reagent	Company	Location
1.	Petroleum Benzine boiling range 60.0°C-80.0°C GR (Petroleum ether)	Merck Specialities Private Limited	Mumbai – 400 018
2.	Pyridine GR		
3.	Ethanol 99.9% AR	Jiangsu Huaxi International Trade Co., Ltd.	China
4.	Distilled water		
5.	Toluene (Sulphur free)	Reachem Laboratory Chemicals	Chennai – 600 098
6.	Formic acid LR	Private Limited	
7.	Acetic acid Glacial LR		
8.	Acetone LR		
9.	Methanol LR	S d Fine Chemicals Limited	Mumbai – 400 030
10.	Ethyl acetate LR		
11.	Silica gel G for TCL	Loba Chemie Private Limited	Mumbai – 400 005

detection will be done at shorter and longer wavelength in UV cabinet<sup>39-55</sup>.

## MATERIALS AND METHODS

### Materials

The intended plant contents are given in the table 01. The instruments, chemicals/reagents and glass wares/apparatus effective for the research are described in the table 02, table 03 and table 04 respectively.

### Miscellaneous

Aluminium foil, Muslin cloth, Filter paper, Tripod stand, Test tube holders and test tube stands and butter paper.

### Methods

#### Plant Collection, drying and powdering

The plant was collected from Parambikulam – Aliyar Riverine in Pollachi. The collected portions of the plant were washed with distilled water three times. They were allowed to dry under shade kept over the news paper. Then the half dried portions were cut into small pieces using stainless steel knife and kept under shade only for drying completely. It took 22 days for complete drying. The dried material was pulverized into coarse powder by means of manual blender. The powdered plant material was stored in air tight containers at 4.0°C for further use. 350.0g of coarse powder of drug was weighed and was taken in a 5000.0ml Round bottomed flask. Petroleum ether was added to remove the fatty matters associated with the powder. The solvent retained was evaporated at room temperature after rinsing for few minutes. Then the dried defatted powder was immersed in 2000.0ml of solvents which comprises 1000.0ml of distilled water and 1000.0ml of ethanol (50:50 v/v). After 7 days, the content of extraction was strained through a muslin cloth. The marc was separated from the menstruum. The extract was kept at 40.0°C for concentration and evaporation at the same temperature. Then the completely dried extract was cooled to room temperature and weighed<sup>56-66</sup>.

#### Preliminary Phytochemicals Evaluation

The experimental procedures were followed for detecting metabolic products of the plant. From the analysis, primary metabolites such as carbohydrates, proteins & aminoacids and secondary metabolites such as “alkaloids, glycosides, flavonoids, tannins, steroids & triterpenoids” were found positive for the tests carried out<sup>67-75</sup>.

#### Thin Layer Chromatographic Analysis

The existing constituents were separated by using proper mobile phase which was selected based on trial and error method<sup>76-77</sup>.

#### Chromatographic parameters

##### Stationary phase selection

Principle: Adsorption

Support material: Glass plate

Dimension of the plate: 20.0×10.0cm

Adsorbent: Silica gel G

Method of thin layer preparation: Pouring method

Layer thickness: 1.0mm

Plate activation temperature: at 105.0°C for one hour

##### Mobile phase selection

Chamber: Twin trough mobile phase chamber

Chamber dimension: 20.0×10.0cm

Mobile phase selected for separation:

##### Alkaloids

Ethyl acetate: Methanol: Water (10.0:1.35:1.0 v/v/v).

##### Glycosides




Figure 1: TLC for mobile phase selection (Trial and error method).


Figure 2: TLC Profile of Hydroalcoholic extract of *Ipomoea aquatic*.

Table 4: Glass wares/Apparatus used.

S. No.	Name of the Glassware	Capacity	Brand Name
1.	Round bottomed flask	1000.0ml	Riviera
2.	Funnel	Medium Size	Sh Borosilicate Glass
3.	Beaker	1000.0ml	Borosilicate Glass
4.	Measuring cylinder	10.0ml	Riviera
5.	Measuring cylinder	50.0ml	Sh Borosilicate Glass
6.	China dish	Big & Small size	Chinese Porcelain
7.	Stirrer	Small size	Sh Borosilicate Glass
8.	Conical flask	250.0ml	Borosilicate Glass
9.	Test tubes	10.0ml	Borosilicate Glass
10.	Pipettes	5.0ml	Borosilicate Glass
11.	Mobile phase chamber (Twin trough)	20×10cm	CAMAG
12.	Beaker	250.0ml	Borosilicate Glass
13.	Petridish lid	Medium size	Borosil S - Line

Ethyl acetate: Pyridine: Water (5.0:1.0:4.0 v/v/v).

*Tannins and Flavonoids*

Toluene: Acetone: Formic acid (4.5:4.5:1.0 v/v/v).

*Steroids and Triterpenoids*

Toluene: Ethyl acetate (9.3:0.7 v/v).

Chamber saturation time: 45.0 minutes

Fluorescence detection: at 365.0nm (Longer wavelength)

Calculation of  $R_f$  value:

$$R_f = \frac{\text{Distance travelled by the solute from the sample application position}}{\text{Distance travelled by the solvent from the sample application position}}$$

Ideal  $R_f$  value: 0.1-0.9

## RESULTS AND DISCUSSION

The TLC mechanisms are interpreted in the figure 01, figure 02 and table 05.

The existence of separated compounds on the stationary phase adsorbent was noticed with different colour of fluorescence at the longer wave length. It was pointed out that the presence of more than ten variable metabolites. These could be quantified, isolated and characterized by using suitable pharmaceutical analytical techniques.

## CONCLUSION

Table 5: Thin Layer Chromatographic Analysis.

S. No.	Sample interest	of Mobile phase	Develop-ment time (min)	Fluorescence spot colour at 365nm	Distance travelled by the solute (cm)	Solvent (cm)	Front	R <sub>f</sub> value
1.	Alkaloids	Ethyl acetate: Methanol: Water (10.0: 1.35: 1.0 v/v/v)	33	Green	0.9	13.6		0.07
				Blue	1.8			0.13
				Green	3.0			0.22
				Blue	4.7			0.35
				Green	6.5			0.48
				Blue	7.5			0.55
				Dark Blue	10.6			0.78
				Green	11.9			0.88
				Pink	12.8			0.94
				Blue	5.2			0.42
2.	Glycosides	Ethyl acetate: Pyridine: Water (5.0: 1.0: 4.0 v/v/v)	107	Green	6.9	12.3		0.56
				Green	8.4			0.68
				Blue	10.3			0.84
				Blue	10.8			0.88
				Pink	11.8			0.96
3.	Tannins and Flavonoids	Toluene: Acetone: Formic acid (4.5: 4.5: 1.0 v/v/v)	30	Brown	3.0	13.2		0.23
				Blue	10.3			0.78
				Pink	10.9			0.83
				Pink	12.5			0.95
4.	Steroids and Tri-terpenoids	Toluene: Ethyl acetate (9.3: 0.7 v/v)	25	Blue	0.7	12.8		0.06
				Green	1.4			0.11
				Pink	2.9			0.23
				Pink	4.6			0.36
				Pink	7.5			0.59

TLC is the best sensitive analytical method for the separation and identification of natural products. The non destructive method of detection supports the separated constituents for isolation by means of scrapping method. The various colour of fluorescence indicates that the presence of different type of similar compounds. After isolation, the particular compound could be characterized using spectroscopic analytical techniques. The mentioned TLC profile for the plant of interest will hopefully help the researchers in herbal field to develop more herbal medicines requiring the particular constituent or its extract.

#### ACKNOWLEDGEMENT

This study was inspired by the Management and Principal of Karpagam Educational Institutions, Coimbatore. We obligate our colleagues from Karpagam College of Pharmacy and Faculty of Pharmacy, Karpagam University, Karpagam Academy of Higher Education, Coimbatore who provided insight and expertise that immensely assisted the analysis.

#### REFERENCES

- [https://en.wikipedia.org/wiki/Ipomoea\\_aquatica](https://en.wikipedia.org/wiki/Ipomoea_aquatica).
- "Kangkong / Ipomoea aquatica Forsk./ POTATO VINE / Herbal Medicinal Therapies / Philippine Alternative Medicine". [Stuartschange.org](http://Stuartschange.org). Retrieved 2014-01-13.
- Chitrajit Malakar and Pinak Pani Nath Choudhury. Pharmacological Potentiality and Medicinal Uses of Ipomoea Aquatica Forsk: A Review. *Asian J Pharm Clin Res.* 2015; 8(2):60-63.
- Manjulika Yadav, Sanjukta Chatterji, Sharad Kumar Gupta and Geeta Watal. Preliminary Phytochemical Screening of Six Medicinal Plants used in Traditional Medicine. *International Journal of Pharmacy and Pharmaceutical Sciences.* 2014;6(5):539-542.
- Mamta Saxena, Jyoti Saxena, Rajeev Nema, Dharmendra Singh and Abhishek Gupta. Phytochemistry of Medicinal Plants. *Journal of Pharmacognosy and Phytochemistry.* 2013;1(6):168-182.
- Selvaraj K, Ranjana Chowdhury and Chiranjib Bhattacharjee. Isolation and structural elucidation of flavonoids from aquatic fern *Azolla microphylla* and evaluation of free radical scavenging activity. *International Journal of Pharmacy and Pharmaceutical Sciences.* 2013;5(3):743-749.
- Salna KP, Sreejith K, Uthiralingam M, Mithu A Prince, John Milton MC and Albin T Fleming. A Comparative Study of Phytochemicals Investigation of *Andrographis paniculata* and *Murraya koenigii*. *International Journal of Pharmacy and Pharmaceutical Sciences.* 2011;3(3):291-292.
- <https://www.medicinalplants-pharmacognosy.com/pharmacognosy-s-topics/extraction-methods/>
- Khadabadi SS, Deore SL and Baviskar BA. Experimental Phytopharmacognosy – A

- Comprehensive guide. Nirali Prakashan, Pune, Maharashtra. Second edition, 2013:p.3.1-3.13, A 1.1-2.8.
10. Ranjith Kumar D, Mohan S., Vanathi P., Prakash G., Sasikala M., Karpakavalli M. Development and Validation of UV Spectroscopic Method for Estimation of Milrinone in Formulation. *World Journal of Pharmaceutical Research*. 2016;5(4):1501-1508.
  11. Sasikala M., Karpakavalli M., Ranjithkumar D., Prakash G., Vanathi P., Mohan S. In-vitro free radical scavenging activity studies of extracts and isolated compounds of *Eugenia jambolana* lam. Seeds. *World Journal of Pharmaceutical Research*. 2016;5(6): 2021-2030.
  12. Sasikala M., Karpakavalli M., Vanathi P., Ranjithkumar D., Prakash G., Mohan S. Validated UV spectrophotometric method development for evaluation and stability analysis of Tolvaptan in bulk drug and its formulation. *World Journal of Pharmacy and Pharmaceutical Sciences*. 2016; 5(6):2048-2063.
  13. Karpakavalli M, Sangilimuthu AY, Sasikala M, Komala M, Ranjithkumar D and Mohan S. A Predominant Evaluation of Viable Templates for Asthma Therapy Exerting an In silico Analog Method, Virtual Library Screening. *Journal of Chemical and Pharmaceutical Research*. 2017; 9(1):43-53.
  14. Sasikala M, Sundara Ganapathy R, Karpakavalli M and Mohan S. Chemotaxonomical and pharmacological review on medicinal plants in temperate region. *International Journal of Pharmacy & Technology*. 2017;9(1): 29165-29172.
  15. Wallis T.E. Text Book of Pharmacognosy. 3rd edn. CBS Publishers and Distributors, Delhi. 1985.
  16. Evans WC. Trease and Evan's Pharmacognosy. 14th Edition. W. B. Saunders Company Limited, London. 1998; p. 119.
  17. Sambamurthy K. Pharmaceutical Engineering. Reprint, New Age International (P.) Ltd., New Delhi. 2002:p. 173-194.
  18. Khandelwal KR. Practical Pharmacognosy – Techniques and Experiments. 9th Edn. Nirali Parkashan, Pune. 2002:82.
  19. Quality control methods for medicinal plant material by WHO; 1998.
  20. Anonymous. Quality control methods for medicinal plant materials. WHO, Geneva. 1998.
  21. Anonymous, the Ayurvedic Pharmacopoeia of India, Vol-I, the Controller of Publication, New Delhi, 2001, 321-322.
  22. Anonymous, The Wealth of India: A Dictionary of Indian Raw Materials and Industrial Products. Vol-2 (B). CSIR, New Delhi. 1998:49-52.
  23. WHO. WHO traditional medicine strategy, [cited May 9]; Available from:[http://www.who.int/medicines/publications/traditional/trm\\_strategy14\\_23/en2014](http://www.who.int/medicines/publications/traditional/trm_strategy14_23/en2014). 2014-23p.
  24. WHO. Quality control methods for medicinal plant materials, [cited May 9]; Available from:<http://apps.who.int/medicinedocs/en/d/Jh1791e2014>.
  25. WHO. General guidelines for methodologies on research and evaluation of traditional medicine, [cited May 9]; Available from:<http://www.who.int/medicines/areas/traditional/definitions/en2014>.
  26. [https://en.wikipedia.org/wiki/Plant\\_secondary\\_metabolism](https://en.wikipedia.org/wiki/Plant_secondary_metabolism).
  27. Kokate CK, Purohit AP and Gokhale SB. Pharmacognosy, 51<sup>st</sup> edition, Nirali Prakashan, Pune, Maharashtra. 2015: A.23-A.27.
  28. Junli Hu, Fuyong Wu, Shengchun Wu, Cheung Lung Lam, Xiangui Lin and Ming Hung Wong. Biochar and *Glomus caledonium* Influence Cd Accumulation of Upland Kangkong (*Ipomoea aquatica* Forsk.) Intercropped with Alfred Stonecrop (*Sedum alfredii* Hance). *Scientific Reports*. 2014;4: 4671. DOI: 10.1038/srep04671.
  29. Dhore Mukund Manohar and Lachure Paresh Shyaam. Survey of Aquatic Macrophyte diversity in Yavatmal District, Maharashtra, India. *Int. J. of Life Sciences*. 2014;2(3):273-275.
  30. Parimala K and Sukumar D. Phytochemical Studies on the Flowers of *Ipomoea Aquatica*. *International Journal of Scientific Research*. 2013;2(6):53-55.
  31. Elija Khatiwora, Vaishali B. Adsul, Manik Kulkarni, N.R. Deshpande and R.V. Kashalkar. Antibacterial activity of Dibutyl Phthalate: A secondary metabolite isolated from *Ipomoea carnea* stem. *Journal of Pharmacy Research*. 2012;5(1):150-152.
  32. Shirin Akhtar, Chandan Karak, Priyanka Biswas, Arup Chattopadhyay and Pranab Hazra. Indigenous Leafy Vegetables: A Potential Source of  $\beta$ -Carotene and Ascorbic Acid. *International Journal of Vegetable Science*. 2012;18:370–375.
  33. Elija Khatiwora, Vaishali B. Adsula, Anjali D. Ruikar, Tushar Gadkari, Deshpande NR and Kashalkar RV. Isolation and Characterization of Dibutyl phthalate from *Ipomoea carnea* stem and its Quantification by HPTLC. *Journal of Pharmacy Research*. 2011;4(10):3264-3265.
  34. Hongfei Fu, Bijun Xie, Shaojun Ma, Xinrong Zhu, Gang Fan and Siyi Pan. Evaluation of antioxidant activities of principal carotenoids available in water spinach (*Ipomoea aquatica*). *Journal of Food Composition and Analysis*. 2011;24:288–297.
  35. Kanchan Lata Vishwakarma and Veenapani Dubey. Nutritional analysis of indigenous wild edible herbs used in eastern Chhattisgarh, India. *Emir. J. Food Agric*. 2011;23(6):554-560.
  36. Mital N. Manvar. Pharmacognostical Investigations on *Ipomoea Aquatica* Forsk. *International Journal of Pharmaceutical Sciences and Research*. 2011;2(11):2812-2815.
  37. Ratnatilaka A, Yakandawala D, Rupasinghe N and Ratnayake K; Poisoning of 'binthamburu' (*Ipomoea asarifolia*) due to misidentification as 'kankun' (*Ipomoea aquatica*); *The Ceylon Medical Journal*. 2010;55(2):54-56.
  38. Maria Greger and Beata Dabrowska. Influence of nutrient level on methyl mercury content in water

- spinach; Environmental Toxicology and chemistry. 2010;29(8):1735–1739.
39. Vimal Kumar, Gogoi BJ, Meghvansi MK, Lokendra Singh, Srivastava RB, Deka DC. Determining The Antioxidant Activity of Certain Medicinal Plants of Sonitpur, (Assam), India Using DPPH Assay. Journal of Phytology. 2009;1(1):49–56.
  40. Shanta IS, Anisuzzaman, Mohanta UK, Farjana T and Mondal MMH. Efficacy of some Indigenous Plants in Controlling Vector Snails of Trematode Parasites of Medical and Veterinary Importance. Bangladesh Journal of Veterinary Medicine. 2008;6(1):107–114.
  41. Aruni Weerasinghe, Sanduni Ariyawansa and Rohan Weerasooriya. Phyto-remediation potential of *Ipomoea aquatica* for Cr(VI) mitigation. Chemosphere. 2008;70(3):521–524.
  42. Wang BL and Xie ZM. Effects of phosphorus application on translocation of lead, zinc and cadmium in the soil-plant system. Huan Jing ke Xue= Huanjing Kexue; 2008;29(11):3225-3229.
  43. Balasuriya BMGK and Dharmaratne HRW. Cytotoxicity and antioxidant activity of green leafy vegetables consumed in Sri Lanka. Journal of National Science Foundation of Sri Lanka. 2007;35(4):255-258.
  44. Daniel F. Austin. Water Spinach (*Ipomoea aquatica*, Convolvulaceae) A food gone wild. Ethnobotany Research & Applications. 2007;5:123-146.
  45. Miao Li, Yue-Jin Wu, Zeng-Liang Yu, Guo-Ping Sheng and Han-Qing Yu. Nitrogen removal from eutrophic water by floating-bed-grown water spinach (*Ipomoea aquatica* Forsk.) with ion implantation. Water Research. 2007;41(14):3152–3158.
  46. Yan L, Wang SH, Luo WG, Huang J and Zhong QS. Study on the oxygen condition in subsurface flow wetlands in operation. Huan Jing ke Xue= Huanjing Kexue. 2006;27(10):2009-2013.
  47. Zhou X, Wang J, Xue L, Xu X and Yang L. N and P removal characters of eutrophic water body under planted float. The Journal of Applied Ecology. 2005;16(11):2199-2203.
  48. Lu Jindeng, Shuai Fangmin, Zhao Liya, Kang Qun and Li Zhaohua. Choosing plants for treating eutrophic water by artificial floating rafts. Journal of Hubei University (Natural Science). 2005;27(4):402-404.
  49. Bhat SC, Goswami S and Ghosh UC. Removal of trace chromium (VI) from contaminated water: bio-sorption by *Ipomea aquatic*. Journal of Environmental Science & Engineering. 2005;47(4):316-321.
  50. Li Xiuling, Li Wei, Li Yeguang, Gui Jianping, Geng Yahong and Hu Hongjun. A primary study on the removal of eutrophication and restraint of algae by some plants. Wuhan Botanical Research. 2005;23(1):53-57.
  51. Sun Zhaohai, Zheng Chunrong, Zhou Dongmei, Yang Linzhang and Chen Huaiman. Effects of soil Pb pollution on growth of *Brassica chinensis* and *Ipomoea aquatica* and urease activities. Rural Eco-Environment. 2005;21(1):38-43.
  52. Qin Houguo, Ye Zhengxiang, Huang Shuijin, Ding Jian and Luo Renhua. The correlation of the different host plants with preference level, life duration and survival rate of *Spodoptera litura* Fabricius. Chinese Journal of Eco-Agriculture. 2004;12(2):40-42.
  53. Cai Q, Mo C, Zeng Q, Li Y, Xiao K, Li H, Xu G, Wang B and Wu Q. Accumulation of di-(2-ethylhexyl) phthalate in various genotype *Ipomoea aquatica*-paddy soil system. The Journal of Applied Ecology. 2004;15(8):1455-1458.
  54. Lin Dongjiao, Tang Shujun, He Jiawen, Luo Jian, Li Yan, Zhang Jian and Liu Shizhe. Purification of piggery wastewater by *Ipomoea aquatica* and *Eichhornia crassipes* in a floating culture system. Journal of South China Agricultural University. 2004;25(3):14-17.
  55. Cai Q, Mo C, Zhu X, Wu Q, Wang B, Jiang C and Li H. Effect of municipal sludge and chemical fertilizers on phthalic acid esters (PAEs) contents in *Ipomoea aquatica* grown on paddy soils. The Journal of Applied Ecology. 2003;14(11):2001-2005.
  56. Zhao Xiaoying, Liu Mingyue, Xiao Jie, Song Yong and Liu Xuanming. Effects of Nitrogen, Kalium, Molybdenum on the Nitrate Accumulation and Nitrate Reductase Activity of Water Spinach. Journal of Hunan Agricultural University. 2003;29(3):239-242.
  57. Agneta Gothberg, Maria Greger and Bengt-Erik Bengtsson. Accumulation of heavy metals in water spinach (*Ipomoea aquatica*) cultivated in the Bangkok region, Thailand. Environmental Toxicology and Chemistry. 2002;21(9):1934-1939.
  58. Hu Shiyi and Yuan Zongfei. Distribution of the cytoplasmic DNA in male cells of the Convolvulaceae and its systematic significance. Acta Phytotaxonomica Sinica. 2002;40(6):511-516.
  59. Koo Hui Mian and Suhaila Mohamed. Flavonoid (Myricetin, Quercetin, Kaempferol, Luteolin, and Apigenin) Content of Edible Tropical Plants. J. Agricultural and Food Chem. 2001;49(6):3106–3112.
  60. Kanungo VK, Sinha S and Naik ML. Net primary productivity of some aquatic macrophytes in sewage-sullage mixture. Journal of Environmental Biology. 2001;22(3):219-223.
  61. Sugandhika Malalavidhane T, Nalinie Wickramasinghe SMD and Jansz ER. Oral hypoglycaemic activity of *Ipomoea aquatica*; Journal of Ethnopharmacology. 2000;72:293 – 298.
  62. En-Jang Sun and Fen-Yi Wu. Wu-Nickel indicator symptoms on water spinach along-vein necrosis as indicator symptom on water spinach caused by nickel in water culture. Bot. Bull. Acad. Sin. 1998;39:255-259.
  63. Zhenbo Yang. The potential of using *Eichhornia crassipes* and *Ipomoea aquatica* as biomonitors in assessing heavy metal pollution in the Mae Kha Canal and the Mae Ping river in Chiang Mai, Northern Thailand. Journal of Social Sciences, Thailand. 1997;23:371-380.
  64. Will RBH and Azhari Rangga. Determination of carotenoids in Chinese Vegetables. Food Chemistry. 1996;56(4):451-455.

65. Tang Shaohu, Sun Min and Li Kunpei. Studies on artificial seed of *Ipomoea aquatica* Forsk. Acta Horticulturae Sinica. 1994;21(1):71-75.
66. Chen BH, Chuang JR, Lin JH and Chiu CP. Quantification of Provitamin A Compounds in Chinese Vegetables by High-Performance Liquid Chromatography. Journal of Food Protection. 1993;56(1):51-54.
67. Chen BH. Determination of carotenoids and chlorophylls in water convolvulus (*Ipomoea aquatica*) by liquid chromatography. Food Chemistry. 1992;45(2):129-134.
68. Hiromu Kameoka, Kanji Kubo and Mitsuo Miyazawa. Essential Oil Components of Water-Convolvulus (*Ipomoea aquatica* Forsk.). Journal of Essential Oil Research. 1992;4(3):219-222.
69. Rahul Kasar, Ashish Gogia, Kartik Shah, Vetrivelan and Chinmay Anand. Critical Factor That Governs a Successful TLC/HPTLC Analysis of Herbal Medicinal Products (HMPS). Research and Reviews: Journal of Pharmaceutical Analysis. 2013; 2(4):1-7.
70. Daya L. Chothani, M. B. Patel and S. H. Mishra. HPTLC Fingerprint Profile and Isolation of Marker Compound of *Ruellia tuberosa*. Chromatography Research International. 2012; 2012:1-6.
71. Peach and Tracey, M.V., Modern Methods of plant Analysis, Springer & Verlag, Berlin, Vol.III, 321-322(1955).
72. Peach, K. and Tracy, M.V., "Modern Methods of Plant Analysis", Vol. III, Springer-Verlag, Berlin, 64 & 464 (1955).
73. Singh J, Bagchi GD, Khanuja SPS (2003). Manufacturing and quality control of Ayurvedic and herbal preparations, In: Verpoorte, R. and Mukherjee, P. K.(Eds), GMP for Botanicals, Regulatory and Quality Issues on Phytomedicine (1st Edition), Business Horizons, New Delhi, p. 201-230.
74. E. Stahl, "Thin layer chromatography," in *A Laboratory Hand Book*, Springer, Berlin, Germany, 2nd edition, 1965.
75. Sethi P.D. Quantitative Analysis of Pharmaceutical Formulations, High Performance Thin Layer Chromatography. CBS Publishers and Distributors, New Delhi, India, 1996, 1-30.
76. Snyder, L.R., Kirkland, J.J., Glajch, J.I., Practical HPLC Method Development. 2nd ed. 13. 1997, 21-173.
77. <http://www.separationsnow.com/details/journal/sepspec1715journal/PhytochemicalAnalysis.html?tzcheck=1>.