

Chemical Composition and Larvicidal Activity of *Mentha pulegium* Essential Oil from Central Morocco Against Larvae of Mosquito *Culex pipiens* (Diptera: Culicidae)

Fouad El-Akhal¹, Yassine Ezzoubi², Taghzouti Khalid³, Abdelhakim El Ouali Lalami^{1,4}

¹Regional Diagnostic Laboratory of Epidemiological and Environmental Health, Regional Health Directorate, EL Ghassani Hospital, 30000 Fez, Morocco.

²Laboratory of Applied Organic Chemistry, Faculty of Sciences and Technology, Sidi Mohamed Ben Abdellah University, B.P. 2202 – Route d'Imouzzer, Fez, Morocco.

³Laboratory of Animal Physiology, Department of Biology, Faculty of Science, University Mohammed V, Rabat, Morocco.

⁴Institute of Nursing Professions and Health Techniques, EL Ghassani Hospital, 30000 Fez, Morocco.

Received 18th Feb, 17 ; Revised 22nd June, 17, Accepted 14th July, 17; Available Online 25th July, 2017

ABSTRACT

Background: In Morocco, the species *Culex pipiens* (Diptera: Culicidae) has developed resistance to the synthetic insecticide. Therefore, there is an urgent need to find alternatives to the insecticides, as natural herbal biocides. Objective: The essential oil of *Mentha pulegium* grown in the North center of Morocco has been analyzed and their larvicidal activity on *Culex pipiens* was determined. Methods: The analysis and the identification of the various constituents of essential oil obtained by hydro-distillation of the aerial part were carried out by Gas Chromatography coupled with Mass Spectrometry (GC/MS). Biological tests were realized according to a methodology inspired from standard World Health Organization protocol. The larvicidal activity of medicinal plant extracts were tested against early fourth-instar larvae of *Culex pipiens*. The repellent efficacy was determined against mosquito species (*Culex pipiens*) at six concentrations (5, 10, 20, 40, 80 and 160 ppm) under laboratory conditions. The mortality was observed 24h after treatment; data were subjected to probit analysis to determine the Lethal Concentrations (LC₅₀ and LC₉₀) to kill 50 and 90 percent of the treated larvae of tested species. Results: The major constituents of *Mentha pulegium* essential oil were identified as pulegone (53.23%), α -pinene (12.1%), piperitone (9.62%), menthone (9.26) and piperitenone (6.6%). The LC₅₀ and LC₉₀ were estimated at 25.45 ppm and 98.66 ppm respectively. Conclusion: This investigation indicates that the *Mentha pulegium* essential oil could serve as a potential larvicidal and could be used as an ideal ecofriendly approach for the control of the *Culex pipiens*.

Keywords: *Mentha pulegium*, Essential oil, Chemical composition, *Culex pipiens*, Larvicidal activity, Fez, Morocco.

INTRODUCTION

Vector-borne diseases remain a public health problem. Mosquitoes constitute an important group of arthropods for public health¹. They transmit a wide range of human diseases such as Filariasis, Malaria, Dengue, Yellow fever and Japanese encephalitis, causing millions of deaths worldwide each year^{2,3}.

The continuous application of synthetic insecticides causes development of resistance in vector species, contamination by toxic substances through the food chain and adverse effects on environmental quality and non target organisms including human health⁴.

It was found that *Culex* (Diptera: Culicidae) populations could develop resistance on account of the widespread use of insecticides⁵⁻⁹. Considering the above, there is an urgent need to develop new insecticides which are biodegradable and environmentally safe¹⁰. Natural products such as extracts and essential oils of plants could be suitable alternatives to use of synthetic insecticides in

the fight against the vectors of diseases in particular versus the *Culex pipiens* species¹¹⁻¹³.

Several plants were chosen for their traditional use as insect repellents¹⁴. Among these plants, *Mentha pulegium* (*M. pulegium*) belonging to the genus of *Mentha* (Lamiaceae), is one of the most popular essential oils crops¹⁵. Essential oil obtained from plants of the genus *Mentha* is a potent medicinal herb with many activities in food, flavours, cosmetics and for pharmaceutical purposes¹⁶. Besides medicinal applications, insecticidal properties of several *Mentha* spp essential oils are reported against ants, mosquitoes, wasps, hornets and cockroaches¹⁷.

This is in order to make our contribution to this part of the research we have harvested, extracted and analyzed the chemical composition of essential oil of *M. pulegium* as well as determined their larvicidal activity on *Culex pipiens* (*C. pipiens*), suspected to be the vector of West Nile in Morocco in 1996^{18,19}, in 2003²⁰. The insecticidal



Figure 1: Photo of plant *Mentha pulegium*.

activity of *M. pulegium* plant against *C. pipiens* has never been studied before in the North center of Morocco.

MATERIALS AND METHODS

Plant material and authentication.

A sample of plant *M. pulegium* (leaves, stems and roots) (Figure 1) was collected in February and March 2014 from natural habitat in province of Taounate (North east of Morocco).

This plant has been identified and a specimen was deposited in the herbarium of The National Institute of Medicinal and Aromatic Plants of Taounate city falling within of the Sidi Mohamed Ben Abdellah University, Fez, Morocco. Samples of 200g of the fresh aerial parts of *M. pulegium* were subjected to hydrodistillation for 2 hours using a Clevenger apparatus; the obtained essential oil was stored at 4°C so that can be used in the upcoming experiments. Essential oil yield was calculated as the percentage of essential oil volume obtained by dry weight of plant used.

Chemical characterization of essential oil.

The chromatographic analysis of essential oil was conducted in "Centre Universitaire Régionale d'Interface" (CURI) in Fez city. The Gas Chromatography (GC) analysis were performed using a Hewlett Packard Gas Chromatographer (HP 6890) with electronic pressure control, equipped with a HP-5MS capillary column (30 m x 0.25 mm, film thickness 0.25 µm), a FID detector set at 300°C and using a H₂/Air mixture, and a *split-splitless* injector set at 220 °C. The injection mode was split (split ratio: 1/50, flow rate: 1.4 ml/min) and the injected volume was about 1 µl. Nitrogen was used as carrier gas with a flow rate of 1.7 ml/min. The column temperature was programmed from 40 to 300 °C at a heating rate of 4°C/min, during 5 min. The apparatus was controlled by a "Chemstation" computer system.

The Gas Chromatography/Mass Spectrometry (GC/MS) analysis were performed by a Hewlett-Packard Gas Chromatographer (HP 6890) coupled with a mass spectrometer (HP 5973). Fragmentation was performed by electron impact at 70 eV. The carrier gas is helium whose flow is fixed at 1.5 ml/min. The injection mode

was split (split ratio: 1/70). The apparatus was controlled by a "Chemstation" computer system.

The identification of the components is based on the comparison of their mass spectra (GC/MS), respective with spectra of the library (NIST 98), of the bibliography²⁰, Kovats index for each compound on OPTIMA-5 column was calculated in reference to n-alkanes.

Characteristic of breeding site.

The collection of larvae of *C. pipiens* was performed in a breeding site located in the urban area of the city of Fez, named "Hafat My Driss" (395 m altitude; 34°04'00.76" latitude and 4°59'28.96" longitude). This site corresponded to very high density of *C. pipiens* larvae.

Collecting C. pipiens larvae.

Larvae were collected using a rectangular plastic in May 2014. This collection of larvae were maintained in breeding at an average temperature of at 23 ± 2°C, 70 ± 10% relative humidity in the Entomology Unit at the Regional Diagnostic Laboratory Epidemiological and Environmental Hygiene falling within Regional Health Directorate of Fez.

Morphological identification of larvae.

Larvae were determined using a Moroccan Culicidae identification key²² and the Mediterranean Africa mosquito's identification software²³.

Protocol for larvicidal efficacy of plant extracts on C. pipiens.

Biological tests were inspired from standard World Health Organization protocol²⁴. A stock solution (1000 ppm) and dilution series for *M. pelugium* extract, was prepared. Preliminary experiments were used to select a range of concentrations to be tested. About 1 ml of each solution prepared was placed in beakers containing 99 ml of distilled water in contact with 20 larvae of stadium 3 (L3) and 4 (L4)²⁴, and the same number of larvae was placed in a beaker containing 99 ml indicator of distilled water plus 1 ml of ethanol. Three replicates were carried for each dilution and for the control.

After 24 h contact, living and dead larvae were counted. The results of susceptibility testing sites were expressed in percentage of mortality versus concentrations of essential oils used. If the % mortality in controls is greater than 5%, the % mortality in larvae exposed to the essential oil is corrected by using Abbott's formula²⁵.

% Mortality Corrected = [(% Mortality Observed - % Mortality Control) / (100 - % Mortality Control)] × 100.

If the control mortality exceeds 20%, the test is invalid and must be repeated.

Data processing.

Probit analysis of concentration-mortality data (Windl version 2.0) software developed by CIRAD-CA/MABIS was conducted to estimate the LC₅₀ values²⁶ and associated 95% confidence limits for each treatment²⁷. The analysis of the averages and standard deviation was also performed by using the test of analysis of variance ANOVA. Mean and standard deviation (± SD) were determined from at least three independent experiments.

RESULTS

Table 1: Percent yield and physical characters of *M. pulegium* from center of Morocco.

Botanical name	Part used	Physical characteristics		Density (g/ml)	Yield(%)
		Color	Odor		
<i>M. pulegium</i>	Areal part	Pale Yellow	Persistent, fragrant Aromatic mint-like	0.934	1.65±0.15

Table 2: Chemical composition of *M. pulegium* essential oil analyzed by GC-MS.

Number	RT	Compound	% Area
1	773	4-isopropyl-1,3-cyclohexanedione	0.6
2	929	α -pinene	12.1
3	913	d-limonene	1.3
4	920	caryophyllene	0.7
5	929	α -humulene	0.2
6	940	3-octanol	1.4
7	963	cyclohexene, 4-methyl-1-(1-methylethenyl)	0.5
8	1021	1.8-cineole	0.2
9	1163	menthone	9.26
10	1078	2-cyclohexen-2-one	0.7
11	1189	α -terpineol	0.9
12	1241	pulegone	53.23
13	1247	piperitone	9.62
14	1323	menthylacetate	0.1
15	1347	piperitenone	6.6
16	1410	β -caryophyllene	0.8
Total		98.21 %	

Yield of essential oil

The yield of essential oil of *M. pulegium* was founded of the order of 1.65%±0.15. This essential oil yield was calculated on the basis of the dry matter.

Chemical composition of essential oil

The composition of essential oil was analyzed by GC and GC/MS showed that the major components of *M. pulegium* are pulegone (53.23%), α -pinene (12.1%), piperitone (9.62%), menthone (9.26) and piperitenone (6.6%) (Table 2).

Variation of mortality

After exposure to different concentrations of essential oil of *M. pulegium* for 24 h, the mortality rate of larvae of *C. pipiens* at stage 3 and 4 varied from 10% to 100% (Figure 2).

The lowest concentration necessary to achieve 100% mortality of larvae of *C. pipiens* was evaluated at 160ppm (Figure 2).

The LC_{50} and LC_{90} Lethal Concentrations

According to the table below (Table 3), LC_{50} and LC_{90} show the larvicidal effect of the essential oils tested. The essential oil of *M. pulegium* has the lowest LC_{50} of 25.45 ppm and an LC_{90} = 98.66 ppm (Equation of the regression line: $Y = -3.06179 + 2.17811 * X$; calculated Chi²: 9.835). According to the shown LC_{50} and LC_{90} values (Table 3), there are no significant differences between the tested essential oils (CI_{95} overlap).

DISCUSSION

Average yield of *M. pulegium* essential oil was 1.65%±0.15. Which is in agreement with that reported by (Derwich et al., 2010)²⁸ which has found a yield of order of 1.54%, from the species *M. pulegium* grown at Skoura near Boulmane, 90 km in the south east of Fez City, central of Morocco. Similar study realized by (Foganholi, et al. 2015)²⁹, demonstrated that the yield of *M. pulegium* essential oil was between 0.17 % and 0.23 %. Oliveira et al., (2011)³⁰ also reported that the yields of *M. pulegium* essential oils grown in Brazil varied between 0.2 % and 0.09 % in the spring and winter, respectively. The differences in the yield of Mentha essential oils with respect to geographical regions were reported by Aziz and Craker (2009)³¹.

In our research, the major constituents of *M. pulegium* essential oil found are: pulegone, α -pinene, piperitone, menthone and piperitenone. In the study realized by (Derwich et al., 2010)²⁸, twenty eight compounds were identified in leaves oil of *M. pulegium* collected from Skoura near Boulemane City (central of Morocco) and the major component was piperitone (35.56%) and other predominant constituents were: piperitenone (21.18%), α -terpineol (10.89%), pulegone (6.452%), pipéritone oxide (4.02%), menthol (3.28%), menthone (3.09%), neomenthol (2.80%), menthofuran (2.15%), isomenthone (1.56%), carvone (1.13%), geranylacétate (1.06%), germacrène D (1.03%) and limonène (1.02%). Daniel et al., (2002)³², studied the composition of *M. Pulegium* oil collected from the Uruguay, they reported that the composition is characterized by a high content of pulegone (73.4%) and isomenthone (12.9%). While, the *M. pulegium* essential oil from Egypt was characterized by pulegone (43.5%) and piperitone (12.2%)³³; and that from Tunisia was composed of: pulegone (41.8%) and isomenthone (11.3%)³⁴. However, the pulegone was found as the main constituent in samples collected in the spring (61.43 %) and in the winter (28.40 %) by (Oliveira et al., 2011)³⁰, followed by trans-caryophyllene (18.68 % and 10.20 %), respectively. Derwich et al., (2010)³⁵ also reported that the piperitone (35.56 %) and piperitenone (21.18 %) as the major constituents of the essential oil of *M. pulegium* from the region of Boulmane (Skoura) of Morocco.

In our study, the insecticidal activity of the oil of *M. pulegium* can be considered very important. Indeed, more than 90% of *C. pipiens* larvae were killed using 98.66 ppm of the essential oil (LC_{50} : 25.45 ppm). Pavela et al. (2014)³⁶, conducted a study on the activities of essential oils of Mentha species on *Culex quinquefasciatus* (Diptera: Culicidae) and also he found a significantly different larvicidal efficacy. In the literature, it has been shown that Mentha species have insecticidal activity against various insects such as *Aedes aegypti* (LC_{50} =




Figure 2: Mortality (%) of larvae of *C. pipiens* depending on the plant species *M. pulegium* essential oil concentration after 24 h of exposure.

Table 3: Concentrations LC_{50} and LC_{90} lethal larvae of *C. pipiens* after 24 h of exposure.

Plant species	LC_{50} (ppm) (LI-UI)*	LC_{90} (ppm) (LI-UI)*
<i>M. pulegium</i>	25.45 (13.86- 44.34)	98.66 (58.50- 130.44)

* LI-UI: Lower limit-Upper limit.

114.33 ppm) and *Culex quinquefasciatus* (LC_{50} = 112.18 ppm)³⁷.

Also, the literature reports that the Mentha oil species, from India, recorded the LC_{50} value of 42.25 ppm and LC_{90} value of 132.41 ppm against the larvae of *Culex quinquefasciatus* at 24 h exposure³⁸. Koliopoulos et al., (2010)³⁹ reported that the LC_{50} values of Mentha species, from Central Greece, ranged from 47.88 to 74.28 ppm and LC_{90} values ranged from 64.34 to 107.45 ppm against *Culex pipiens* after 48h exposure. The important larvicidal activity observed among the essential oil of *M. pulegium* might be explained by the action or the effect of the major components. Inhere results are the first in Morocco; this might allow developing an effective low cost and eco-friendly larvicides.

As a perspective of this study and as part of our researchs on bio-insecticide conducted in the laboratory^{40,41}, we plan to conduct studies on the larvicidal activity of *M. pulegium* essential oil against other species of the Culicidae family, including the species *Anopheles labranchaie*, responsible for indigenous malaria in Morocco.

CONCLUSION

In the present work, realized for the first time in Morocco, we were interested to examine the chemical composition and to evaluate the effect of *Mentha pulegium* essential oil on the toxicity of mosquito larvae (*Culex pipiens*). This study demonstrated that the GC/MS analysis of *Mentha puegium* showed that the major components were: pulegone (53.23%), α -pinene (12.1%), piperitone (9.62%), menthone (9.26) and piperitenone (6.6%). The essential oil of *M. pulegium* showed a

larvicidal activity against *Culex pipiens* larvae in stages 3 and 4, with respective values of LC_{50} and LC_{90} about 25.45 ppm and 98.66 ppm. Therefore, this essential oil could be a promising alternative against mosquitoes.

ACKNOWLEDGMENT

We thank everyone who contributed to this work.

CONFLICT OF INTERESTS

The authors declare that they have no conflict of interests.

REFERENCES

1. Eliningaya JK. Tamires CL. Chrian MM. Damião PS. Larvicidal efficacy of monoterpenes against the larvae of *Anopheles gambiae*. Asian Pac J Trop Biomed 2016; 6(4): 290–294.
2. Das NG. Goswami D. Rabha B. Preliminary evaluation of mosquito larvicidal efficacy of plant extracts. J Vector Borne Dis 2007; 44: 145-8.
3. Pugazhvendan SR. Elumali K. Larvicidal activity of selected plant essential oil against important vector mosquitoes: dengue vector, *Aedes aegypti* (L.), malarial vector, *Anopheles stephensi* (Liston) and filarial vector, *Culex quinquefasciatus* (Say) (Diptera: Culicidae). Middle East J Sci Res 2013; 18(1): 91-5.
4. Ghosh A. Chowdhury N. Chandra G. Plant extracts as potential mosquito larvicides. Indian J Med Res 2012; 135: 581-598.
5. Faraj C. El Kohli M. El Rhazi M. Laqraa M. Lyagoubi M. Niveau actuel de la résistance du moustique *Culex pipiens* aux insecticides au Maroc. Sc Lett 2002; 4: 4.
6. Jones CM. Machim C. Mohammed K. Majambere S. Ali SA. Khatib BO. Mcha J. Ranson H. Kelly-Hope

- LA. Insecticide resistance in *Culex quinquefasciatus* from Zanzibar: implications for vector control. *Parasites & Vectors* 2012; 5: 78.
7. El Ouali Lalami A. El-Akhal F. El Amri N. Maniar S. Faraj C. State resistance of the Mosquito *Culex pipiens* towards Temephos central Morocco. *Bull. Soc. Pathol. Exot* 2014; 107: 194-198.
 8. El Joubari M. Faraj C. Louah A. Himmi O. Susceptibility of *Anopheles labranchiae*, *Culex pipiens*, *Ochlerotatus detritus* and *Ochlerotatus caspius* mosquitoes in the Smir region (northwest Morocco) to organophosphates used in public health. *Environ Risque Sante* 2015; 14(1): 72-79.
 9. EL-Akhal F. Maniar S. El Bachir A. Chafiq F. Badoc A. El Ouali Lalami A. Resistance of *Culex pipiens* (Diptera: Culicidae) to Organophosphate Insecticides in Central Morocco. *International Journal of Toxicological and Pharmacological Research* 2016; 8(4): 263-568.
 10. Pavela R. Vrchotová N. Tríska J. Mosquitocidal activities of thyme oils (*Thymus vulgaris* L.) against *Culex quinquefasciatus* (Diptera: Culicidae). *Parasitology Research* 2009; 105: 1365-1370.
 11. El Ouali Lalami A., El-Akhal F., Ez Zoubi Y., Taghzouti K. Study of Phytochemical Screening and Larvicidal Efficacy of Ethanolic Extract of *Salvia officinalis* (Lamiaceae) from North Center of Morocco Against *Culex pipiens* (Diptera: Culicidae) Vector of Serious Human Diseases. *International Journal of Pharmacognosy and Phytochemical Research* 2016; 8(10); 1663-1668.
 12. Ez zoubi Y., El-Akhal F., Farah A., El Ouali Lalami A. Phytochemical Screening and Larvicidal Activity of Moroccan *Ammi visnaga* Against Larvae West Nile Vector Mosquito *Culex pipiens* (Diptera: Culicidae). *International Journal of Pharmacognosy and Phytochemical Research* 2016; 8(10); 1684-1688.
 13. El-Akhal F. El Ouali Lalami A. Ez Zoubi Y. Greche H. Guemmouh R. Chemical composition and larvicidal activity of essential oil of *Origanum majorana* (Lamiaceae) cultivated in Morocco against *Culex pipiens* (Diptera: Culicidae). *Asian Pac J Trop Biomed* 2014; 4(9): 746-750.
 14. Tisgratog R. Sanguanpong U. Grieco JP. Ngoen-Kluan R. Plants traditionally used as mosquito repellents and the implication for their use in vector control. *Acta Tropica* 2016; 157: 136-144.
 15. Padalia RC. Verma RS. Chauhan A. Sundaresan V. Chanotiya CS. Essential oil composition of sixteen elite cultivars of *Mentha* from western Himalayan region, India. *Maejo International Journal of Science and Technology* 2013; 7(1): 83-93.
 16. Al-Rawashdeh IM. Molecular taxonomy among *Mentha spicata*, *Mentha longifolia* and *Ziziphora tenuior* populations using the RAPD technique. *Jordan Journal of Biological Science* 2011; 4: 63-70.
 17. Conceição C. Matos O. Barbosa A. Mexia A. Potential of plant products As protectants of stored maize against *Sitophilus zeamais* Motschulsky (Coleoptera: Curculionidae). In: *Proceedings 20th IWCSP2010*: 425: 618-624.
 18. Harrack ME. Guenno LEB. Gounno LEP. Isolement du virus West Nile au Maroc. *Virologie* 1997; 1(6): 248-249.
 19. Murgue B. Murri S. Triki H. Deubel V. Zeller HG. West Nile in the Mediterranean basin: 1950-2000. *Ann N Y Acad Sci* 2001; 951: 117-126.
 20. Schuffenecker I. Peyrefitte CN. Harrak MEL. Murri S. Lebond A. Zeller HG. West Nile virus in Morocco, 2003. *Emerg Infect. Diseases* 2005; 11: 306-309.
 21. Adams RP. Identification of Essential Oil Components by Gas Chromatography/Mass Spectrometry, 4th Edition. Allured Publishing Corporation 2007; p 1-804.
 22. Himmi O. Dakki M. Trari B. El Agbani MA. The culicidae of Morocco: identification keys with biological and ecological data (work of the Scientific Institute), Rabat: Rabat institute Scientific 1995: 44: 1-51.
 23. Brunhes J. Rhaim A. Geoffroy B. Hervy JP. Mosquitoes of the Mediterranean Africa: software identification and education, Paris (FRA); Tunis: IRD; IPT, 2000, 1 CD-ROM (Didactiques). ISBN2-7099-1446-8.
 24. WHO. Guidelines for laboratory and fieldtesting of mosquito larvicides, Tech. Rep. Who/cds/whopes/gcdpp/2005.13, 2005.
 25. Abbott WS. A methode of computing the effectiveness of an insecticide. *J. Econ. Entomol* 1925; 18(2): 265-267.
 26. Giner M, Vassal JM, Vassal C, Chiroleu F, Kouaik Z. WinDL Software version 2.0, CIRAD-CA. URBI/MABIS, Montpellier, France., 1999.
 27. Finney DJ. Probit Analysis (3rd edition). Cambridge University Press, Cambridge, UK., 1971.
 28. Derwich E. Benziane Z. Taouil R. Senhaji O. Touzani M. Comparative Essential oil Composition of Leaves of *Mentha rotundifolia* and *Mentha pulegium* a Traditional Herbal Medicine in Morocco. *American-Eurasian Journal of Sustainable Agriculture* 2010; 4(1): 47-54.
 29. Foganholi APAM. Daniel JFS. Santiago DC. Orives JR. Pereira JP. Faria TJ. Chemical composition and antifungal activity of pennyroyal essential oil in different stages of development. *Semina: Ciências Agrárias, Londrina* 2015; 36(5): 3091-3100.
 30. Oliveira RA. SA ICG. Duarte LP. Oliveira FF. Constituintes voláteis de *Mentha pulegium* L. e *Plectranthus amboinicus* (Lour.) Spreng. *Revista Brasileira Plantas Medicinai, Campinas* 2011; 13(2): 165-169.
 31. Aziz EE. Craker LE. Essential oil constituents of peppermint, pennyroyal, and apple mint grown in a desert agrosystem. *Journal of Herbs, Spices & Medicinal Plants, Philadelphia* 2009; 15 (4): 361-367.
 32. Daniel L. Daniel P. Eduardo D. Philip D. Roser V. Salvador C. Essential oils of *Mentha pulegium* and *Mentha rotundifolia* From Uruguay. *Brazi. Archi. Biol and Technol* 2002; 45(4): 519-524.

33. El-Ghorab AH. The chemical composition of *Mentha pulegium* L. essential oil from Egypt and its antioxidant activity. *Journal of Essential Oil Bearing Plant* 2006; 9: 183-195.
34. Mkaddem M. Bousaid M. Ben Fadhel N. Variability of volatiles in Tunisian *Mentha pulegium* L. (Lamiaceae). *Journal Essent Oil Res* 2007; 19: 211-215.
35. Derwich E. Benziane Z. BOUKIR A. GC/ MS analysis and antibacterial activity of the essential oil of *Mentha pulegium* grown in Morocco. *Research Journal of Agriculture and Biological Sciences, Faisalabad* 2010; 6(3): 191-198.
36. Pavela R. Kaffková K. Kumšta M. Chemical composition and larvicidal activity of essential oils from different *Mentha* L. and *Pulegium* species against *Culex quinquefasciatus* Say (Diptera: Culicidae). *Plant Protect. Sci* 2014; 50: 36-42.
37. Manzoor F. Samreen KB. Parveen Z. Larvicidal activity of essential oils against *Aedes aegypti* and *Culex quinquefasciatus* larvae (Diptera: Culicidae). *The Journal of Animal & Plant Sciences* 2013; 23(2):420-424.
38. Manimaran A. Mary JeeJee Cruz M. Muthu C. Vincent S. Ignacimuthu S. Larvicidal and knockdown effects of some essential oils against *Culex quinquefasciatus* Say, *Aedes aegypti* (L.) and *Anopheles stephensi* (Liston). *Advances in Bioscience and Biotechnology* 2012; 3: 855-862.
39. Koliopoulos G. Pitarokili D. Kioulos E. Michaelakis A. Tzakou O. Chemical composition and larvicidal evaluation of *Mentha*, *Salvia*, and *Melissa* essential oils against the West Nile virus mosquito *Culex pipiens*. *Parasitology Research* 2010;107: 327-335.
40. El Ouali Lalami A., EL-Akhal F., Maniar S., Ez zoubi Y., Taghzouti K. Chemical Constituents and Larvicidal Activity of Essential Oil of *Lavandula Stoechas* (Lamiaceae) From Morocco Against the Malaria Vector *Anopheles Labranchiae* (Diptera: Culicidae). *International Journal of Pharmacognosy and Phytochemical Research* 2016; 8(3); 505-511.
41. El-Akhal F. El Ouali Lalami A. Guemmouh R. Larvicidal activity of essential oils of *Citrus sinensis* and *Citrus aurantium* (Rutaceae) cultivated in Morocco against the malaria vector *Anopheles labranchiae* (Diptera: Culicidae). *Asian Pac J Trop Dis* 2015; 5(6): 930-934.