

## An Observational Assessment of Endoscopic Transcortical Transventricular Management of Cystic Craniopharyngioma

Dhiraj Kumar<sup>1</sup>, Mukesh Kumar<sup>2</sup>, Anil Kumar Peethambaran<sup>3</sup>

<sup>1</sup>Senior resident, Department of Neurosurgery, Medical College Trivandrum Thiruvananthapuram, Kerala, India

<sup>2</sup>Senior resident, Department of Neurosurgery, Medical College Trivandrum Thiruvananthapuram, Kerala, India

<sup>3</sup>HOD, Department of Neurosurgery, Medical College Trivandrum Thiruvananthapuram, Kerala, India

Received: 18-09-2023 / Revised: 12-10-2023 / Accepted: 20-11-2023

Corresponding Author: Dr. Dhiraj Kumar

Conflict of interest: Nil

### Abstract

**Aim:** The aim of the present study was to assess the endoscopic transcortical transventricular management of cystic craniopharyngioma.

**Methods:** The present study was conducted in the Department of Neurosurgery, Medical College Trivandrum Thiruvananthapuram, Kerala, India and data of patients who underwent transcortical transventricular endoscopic approach and reservoir placement for cystic craniopharyngioma were reviewed retrospectively from August 2016 to August 2018. 10 patients were included in the study.

**Results:** The mean age was 22.6 years (range 2–49 years) with a slight female preponderance. Headache (75%) was the most common presentation followed by vomiting (55%) and vision disturbances (45%). An equal number of patients presented with motor weakness and memory disturbances (12%). 5 patients presented with altered behavior and 5 patients with polyuria. Most of the patients had a single cyst. All patients had hydrocephalus. The presurgery hormone evaluation showed that 10% of patients had hypocortisolemia and 6% of patients had marginally increased prolactin levels. The follow-up duration ranged from 3 to 72 months with a median of 18 months. 38 (38%) patients had improvement in vision. 20 (20%) patients required thyroid hormone replacement and 18 patients (18%) steroid replacement. 42 (42%) patients received radiotherapy. 28 (28%) cases had a cyst recurrence and required repeat intervention.

**Conclusion:** Transventricular endoscopy is a versatile procedure that plays an important part in the management of craniopharyngioma, especially for those lesions with a predominant cystic component growing within the third ventricle. It has been used to fenestrate cystic tumors into the ventricles or subarachnoid space (ventriculocystostomy or cystocisternostomy), to establish CSF flow in case of hydrocephalus, for delivery of intracavitary chemotherapy, and to achieve variable extent of resection as a first step of combined procedure.

**Keywords:** Endoscopy, Neuroendoscopy, Hydrocephalus, Craniopharyngioma, Reservoir.

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

### Introduction

Craniopharyngiomas are benign sellar and suprasellar tumors, deemed to originate from the cellular residue of Rathke's pouch. [1,2] Half of all cases occur in adulthood with a peak incidence at the age of 40–44 years and another peak at the sixth decade. They have a variable worldwide incidence ranging from 2% to 5% of all primary intracranial tumors. [2,3] The malignant course is noted in long-term management due to compression of the pituitary gland and stalk, the tendency to repeated cystic recurrence with the potential invasion of the hypothalamus and third ventricle, which further complicates the clinical status due to hydrocephalus. Hydrocephalus is present in 20–

30% of adult craniopharyngioma at the time of presentation. [4-6]

Cystic craniopharyngiomas present a unique challenge because of the possibility of recurrence of cyst irrespective of the modality of initial treatment. [7] Almost 90% of craniopharyngiomas have a cystic component. [8] The presence of a cyst in craniopharyngioma can account for major total bulk of the tumor, and a predominantly cystic tumor is seen in 60% of craniopharyngiomas. [9] The traditional approach to cystic craniopharyngioma is a variable extent of microsurgical resection. Ommaya reservoir is often

placed in the residual cyst for repeated aspiration or intracavitary therapy. [10] However, the malposition of reservoir is not uncommon. [11,12]

Microscopic surgical resection and endoscopic cyst fenestration along with intracystic catheter placement of the Ommaya reservoir are the commonly performed surgical techniques. [13,14] Transcortical transventricular endoscopic surgery for cystic craniopharyngioma has gained acceptance as a safe minimally invasive alternative to resection. [15,16] The traditional approach to cystic craniopharyngioma is a variable extent of microsurgical resection. Ommaya reservoir is often placed in the residual cyst for repeated aspiration or intracavitary therapy. [10] However, malposition of the reservoir is not uncommon. [11,12] In such circumstances a neuroendoscopic approach provides a less invasive and more accurate technique to deal with cystic craniopharyngiomas. [11,17]

The aim of the present study was to assess the endoscopic transcortical transventricular management of cystic craniopharyngioma.

#### Materials and Methods

The present study was conducted in the Department of Neurosurgery, Medical College Trivandrum Thiruvananthapuram, Kerala, India and data of patients who underwent transcortical transventricular endoscopic approach and reservoir placement for cystic craniopharyngioma were reviewed retrospectively from August 2016 to August 2018. 10 patients were included in the study. The patients were identified and none of the patients' personal information is included in the article so that none of these patients can be identified or contacted, considering this, retrospective studies are waived off from review by our institute's ethics committee. The clinical features, imaging findings, surgical steps, complications, and outcomes were recorded. All patients underwent a neurological and neuro-ophthalmological examination before surgery and at follow-up. Basal serum levels of anterior pituitary hormones were assessed for all patients before surgery and at follow-up. All patients underwent preoperative magnetic resonance imaging of the brain, and computerized tomography scan of the head immediately after surgery and at follow-up.

#### Technique

The technique of transcortical transventricular endoscopic approach and Ommaya reservoir placement for cystic craniopharyngioma was similar to any intraventricular cystic tumors.

The patient was positioned supine with the head in neutral position. A semilunar scalp flap based on coronal suture was raised. A precoronal burr hole, 3 cm from the midline, was made. The ventricles were entered with trocar and endoscope sheath. Gaab 0° telescope (Karl Storz, Tuttlingen, Germany), was introduced in the ventricle and the cyst was identified at the foramen of Monro. The 0° scope was removed and the operating scope was introduced.

The cyst wall was coagulated, and a piece of wall was taken for biopsy. On puncturing the cyst olive green (machine oil) colored fluid, typical of craniopharyngioma was aspirated. Continuous irrigation with suction was done to remove all the fluid, till the contents of the cyst became clear. The inner wall of the cyst was inspected and a biopsy was taken from the wall of the cyst and solid portion. No attempt was made to resect the solid portion of the tumor. A further fenestration at the deeper portion of the cyst for communication to the basal cistern was attempted only if the cyst wall was transparent and the basilar artery could be easily visualized. Neuronavigation was used for multiloculated cysts to ensure communication between cysts. At the end of irrigation, a ventricular catheter was passed through the operating channel of scope under direct vision till it touched the deepest wall of the cyst. There was good three-dimensional visualization and orientation. The operating scope along with the sheath was slowly withdrawn under vision leaving behind the ventricular catheter. The catheter was connected to the reservoir in the subgaleal space. The reservoir was punctured to aspirate fluid and confirm patency. The histopathology of the cyst wall confirmed the diagnosis of craniopharyngioma in all cases. All patients received perioperative antibiotics, steroids, and antiepileptic drugs. Adults and children over 7 years of age were referred for radiotherapy.

#### Results

**Table 1: Demographic details**

Parameter (N=50)	
Age	Mean 22.6 y (range 1.8–49)
Gender	Male 45 (45%)
	Female 55 (55%)
Clinical features	
Headache	75 (75%)
Vision disturbances	45 (45%)

Polyuria	5 (5%)
Vomiting	55 (55%)
Motor weakness	12 (12%)
Memory disturbances	12 (12%)
Altered behavior	5 (5%)
Radiology	
Single cyst	75 (75%)
Multiloculated	25 (25%)
Endocrine disturbances	
Hypocortisolism	10 (10%)
Marginally increased prolactin (stalk effect)	6 (6%)

The mean age was 22.6 years (range 2–49 years) with a slight female preponderance. Headache (75%) was the most common presentation followed by vomiting (55%) and vision disturbances (45%). An equal number of patients presented with motor weakness and memory disturbances (12%). 5

patients presented with altered behavior and 5 patients with polyuria. Most of the patients had a single cyst. All patients had hydrocephalus. The presurgery hormone evaluation showed that 10% of patients had hypocortisolemia and 6% of patients had marginally increased prolactin levels.

**Table 2: Surgical details and follow-up data**

Parameter (n=100)	
Surgery duration	Median 90 min
Hospital stay	Mean 3.4 d
Follow-up (n¼ 29)	
Duration	Median 18 mo (3–72 mo)
Cyst recurrence	28 (28%)
Reaspiration	18 (18%)
Craniotomy	8 (8%)
Shunt	4 (4%)
Median recurrence-free survival	43 mo
Vision improved	38 (38%)
Radiotherapy	42 (42%)
Hormone replacement	
Thyroid	20 (20%)
Steroid	18 (18%)
Desmopressin	4 (4%)

The follow-up duration ranged from 3 to 72 months with a median of 18 months. 38 (38%) patients had improvement in vision. 20 (20%) patients required thyroid hormone replacement and 18 patients (18%) steroid replacement. 42 (42%) patients received radiotherapy. 28 (28%) cases had a cyst recurrence and required repeat intervention.

### Discussion

Surgery for craniopharyngioma has a major negative impact on long-term quality of life (QoL). In a recent literature review on quality-adjusted life years (QALYs) of four surgical approaches to craniopharyngiomas: aggressive tumor removal (attempt at gross total resection [GTR]), planned subtotal removal plus radiotherapy, biopsy plus radiotherapy, and transnasal endoscopic resections were reviewed. [18] Cystic craniopharyngiomas present a unique challenge because of the possibility of recurrence of cyst irrespective of the modality of initial treatment. Almost 90% of craniopharyngiomas have a cystic component. [19]

Endoscopy has some advantages over other drainage techniques. Through a minimally invasive approach it is possible to perforate the capsule under direct visual control as compared with repeated attempts of perforation and the risk of deviation from the tract due to the consistency of the cystic wall in stereotactic procedures. Endoscopy offers more radical cyst drainage, partial resection of the capsule and sampling of tissue fragments under direct vision for diagnostic purposes avoiding the risks of closed biopsy, and wide marsupialization into cerebrospinal fluid (CSF) fluid. The most common procedure performed through the transventricular endoscopic approach for craniopharyngiomas is fenestration and aspiration of the cyst with catheter placement inside the cyst connected to a subgaleal reservoir. Instead of placement of reservoir Delitala et al [9] positioned a transcystic multiholed catheter with ends reaching into the cistern (caudally) and lateral ventricle (rostrally). In such stenting, ventricle acts as reservoir in which the cyst fluid collects and gets

absorbed along with CSF. This stenting was devised in an attempt to promote continuous CSF circulation and prevent the cyst's reclosure. The mean age was 22.6 years (range 2–49 years) with a slight female preponderance. Headache (75%) was the most common presentation followed by vomiting (55%) and vision disturbances (45%). An equal number of patients presented with motor weakness and memory disturbances (12%). 5 patients presented with altered behavior and 5 patients with polyuria. Most of the patients had a single cyst. All patients had hydrocephalus. The presurgery hormone evaluation showed that 10% of patients had hypocortisolemia and 6% of patients had marginally increased prolactin levels.

Nakamizo et al [16] had an early single case report of trans ventricular endoscopic craniopharyngioma drainage and Ommaya catheter and reservoir placement. Nakahara et al [13] reported three adult cases of trans-ventricular endoscopic management for craniopharyngioma with mixed solid/cystic components where they could evacuate the cyst contents and obtain a biopsy. The presurgery hormone evaluation showed that 10% of patients had hypocortisolemia and 6% of patients had marginally increased prolactin levels. The follow-up duration ranged from 3 to 72 months with a median of 18 months. 38 (38%) patients had improvement in vision. 20 (20%) patients required thyroid hormone replacement and 18 patients (18%) steroid replacement. 42 (42%) patients received radiotherapy. 28 (28%) cases had a cyst recurrence and required repeat intervention.

Previously, complex cases of cystic craniopharyngiomas that invade the third ventricle and induce obstructive hydrocephalus were treated at two steps. Initially, a transcranial trans-ventricular endoscopic fenestration of the cyst wall with placing a catheter into the cyst to release the increased intracranial pressure. This is followed by another elective intervention within the next days by an endonasal trans-sphenoidal procedure for excision of the tumour. [20] Furthermore, Park et al [21] described vision loss as the most common presenting symptom (83%), followed by headache (48%) and endocrinopathy (33%) in a multicentre study. Further, Cavallo et al [22] found visual disturbances (76.7%) and endocrinopathy (56.6%) in affected adult patients.

### Conclusion

Transventricular endoscopy is a versatile procedure that plays an important part in the management of craniopharyngioma, especially for those lesions with a predominant cystic component growing within the third ventricle. It has been used to fenestrate cystic tumors into the ventricles or subarachnoid space (ventriculocystostomy or cystocisternostomy), to establish CSF flow in case

of hydrocephalus, for delivery of intracavitary chemotherapy, and to achieve variable extent of resection as a first step of combined procedure. Among all the endoscopic treatment options the preferred treatment is fenestration, aspiration of cyst, and placement of catheter reservoir followed by radiation. The transventricular endoscopic approach is not shown to be particularly effective for improvement of vision in patients with complete vision loss before surgery. Residual tumor with persistent symptoms would require resection for relief of symptoms of neural compression.

### References

1. Balde NM, Diallo MM, Poirier JY, Sow MS, Brassier G, Lorcy Y. Long-term outcome of the adult onset craniopharyngiomas. In *Annales D'endocrinologie* 2007 May 25 (Vol. 68, No. 2-3, pp. 186-190).
2. Clark AJ, Cage TA, Aranda D, Parsa AT, Sun PP, Auguste KI, Gupta N. A systematic review of the results of surgery and radiotherapy on tumor control for pediatric craniopharyngioma. *Child's Nervous System*. 2013 Feb; 29:231-8.
3. Dandurand C, Sepehry AA, Lari MH, Akagami R, Gooderham P. Adult craniopharyngioma: case series, systematic review, and meta-analysis. *Neurosurgery*. 2018 Oct 1; 83(4):631-41.
4. Daubenbüchel AM, Hoffmann A, Gebhardt U, Warmuth-Metz M, Sterkenburg AS, Müller HL. Hydrocephalus and hypothalamic involvement in pediatric patients with craniopharyngioma or cysts of Rathke's pouch: impact on long-term prognosis. *European journal of endocrinology*. 2015 May; 172(5):561-9.
5. Karavitaki N, Brufani C, Warner JT, Adams CB, Richards P, Ansorge O, Shine B, Turner HE, Wass JA. Craniopharyngiomas in children and adults: systematic analysis of 121 cases with long-term follow-up. *Clinical endocrinology*. 2005 Apr; 62(4):397-409.
6. Van Effenterre R, Boch AL. Craniopharyngioma in adults and children: a study of 122 surgical cases. *Journal of neurosurgery*. 2002 Jul 1; 97(1):3-11.
7. Muller HL. Childhood craniopharyngioma. Recent advances in diagnosis, treatment and follow-up. *Hormone Research*. 2008 Jan 21; 69(4):193-202.
8. Barajas MA, Ramírez-Guzmán G, Rodríguez-Vázquez C, Toledo-Buenrostro V, Velásquez-Santana H, del Valle Robles R, Cuevas-Solórzano A, Rodríguez-Hernández G. Multimodal management of craniopharyngiomas: neuroendoscopy, microsurgery, and radiosurgery. *Journal of neurosurgery*. 2002 Dec 1; 97 (Supplement 5):607-9.

9. Delitala A, Brunori A, Chiappetta F. Purely neuroendoscopic transventricular management of cystic craniopharyngiomas. *Child's Nervous System*. 2004 Nov; 20:858-62.
10. Gutin PH, Klemme WM, Lagger RL, MacKay AR, Pitts LH, Hosobuchi Y. Management of the unresectable cystic craniopharyngioma by aspiration through an Ommaya reservoir drainage system. *J Neurosurg* 1980;52(1):36-40
11. Mori R, Joki T, Nonaka Y, Ikeuchi S, Abe T. Parallel insertion endoscopic technique for precise catheter placement in cystic craniopharyngiomas. *J Neurol Surg A Cent Eur Neurosurg* 2014;75(6):442-446
12. Zanon N, Cavalheiro S, da Silva MC. Does the choice of surgical approach to insert an intratumoral catheter influence the results of intratumoral cystic treatment? *Surgical neurology*. 2008 Jul 1;70(1):66-9.
13. Nakahara Y, Koga H, Maeda K, Takagi M, Tabuchi K. Neuroendoscopic transventricular surgery for suprasellar cystic mass lesions such as cystic craniopharyngioma and Rathke cleft cyst. *Neurologia medico-chirurgica*. 2004;44(8):408-15.
14. DiPatri AJ Jr., Prabhu V. A history of the treatment of craniopharyngiomas. *Childs Nerv Syst* 2005; 21:606-21.
15. Joki T, Oi S, Babapour B, Kaito N, Ohashi K, Ebara M, Kato M, Abe T. Neuroendoscopic placement of Ommaya reservoir into a cystic craniopharyngioma. *Child's Nervous System*. 2002 Nov; 18:629-33.
16. Nakamizo A, Inamura T, Nishio S, Inoha S, Ishibashi H, Fukui M. Neuroendoscopic treatment of cystic craniopharyngioma in the third ventricle. *min-Minimally Invasive Neurosurgery*. 2001 Jun;44(02):85-7.
17. Pettorini BL, Tamburrini G, Massimi L, Caldarelli M, Di Rocco C. Endoscopic transventricular positioning of intracystic catheter for treatment of craniopharyngioma. Technical note. *J Neurosurg Pediatr* 2009;4(03):245-248.
18. Ali ZS, Bailey RL, Daniels LB, Vakhshori V, Lewis DJ, Hossain AT, Sitterley KY, Lee JY, Storm PB, Heuer GG, Stein SC. Comparative effectiveness of treatment options for pediatric craniopharyngiomas. *Journal of Neurosurgery: Pediatrics*. 2014 Feb 1;13(2):178-88.
19. Barajas MA, Ramírez-Guzmán G, Rodríguez-Vázquez C, Toledo-Buenrostro V, Velásquez-Santana H, del Valle Robles R, Cuevas-Solórzano A, Rodríguez-Hernández G. Multimodal management of craniopharyngiomas: neuroendoscopy, microsurgery, and radiosurgery. *Journal of neurosurgery*. 2002 Dec 1;97(Supplement 5):607-9.
20. Deopujari CE, Karmarkar VS, Shah N, Vashu R, Patil R, Mohanty C, Shaikh S. Combined endoscopic approach in the management of suprasellar craniopharyngioma. *Child's Nervous System*. 2018 May; 34:871-6.
21. Park HR, Kshetry VR, Farrell CJ, Lee JM, Kim YH, Won TB, Han DH, Do H, Nyquist G, Rosen M, Kim DG. Clinical outcome after extended endoscopic endonasal resection of craniopharyngiomas: two-institution experience. *World neurosurgery*. 2017 Jul 1; 103:465-74.
22. Cavallo LM, Frank G, Cappabianca P, Solari D, Mazzatenta D, Villa A, Zoli M, D'Enza AI, Esposito F, Pasquini E. The endoscopic endonasal approach for the management of craniopharyngiomas: a series of 103 patients. *Journal of neurosurgery*. 2014 Jul 1;121(1):100-13.