

**Nerve Conduction Study in Vitamin B 12 Deficiency Patients**Shaik Meera Sharief<sup>1</sup>, Arjuman Parveen Shaik<sup>2</sup>, Shaik Auliya Parveen<sup>3</sup><sup>1</sup>Associate Professor, Department of Neurology, Rangaraya Medical College, Kakinada, Andhra Pradesh<sup>2</sup>Assistant Professor, Department of Physiology, Rangaraya Medical College, Kakinada, Andhra Pradesh<sup>3</sup>Assistant Professor, Department of Physiology, Guntur Medical College, Guntur, Andhra Pradesh

Received: 08-11-2023 / Revised: 14-11-2023 / Accepted: 16-12-2023

Corresponding Author: Shaik Meera Sharief

Conflict of interest: Nil

**Abstract**

Vitamin B12 deficiency is clinically recognized to be associated with neurological disorders, such as dementia, cognitive impairment, and depression. B12 deficiency may cause demyelination of nerves in the peripheral and central nervous system and has been associated with peripheral neuropathy, loss of sensation in peripheral nerves, and weakness in lower extremities in older adults. 30 Vitamin B12 deficient subjects and 30 normal subjects as control were included in study. Nerve conduction study was done using the AD instrument –Nicolet in both groups. Comparison of sensory nerve parameters in Control Group and Vitamin B12 deficient subjects, there is highly significant slowing of sensory nerves median & ulnar ( $p < 0.001$ ) and significant decrease in amplitude of Median nerve and ulnar nerve ( $p < 0.01$ ) in Vitamin B12 deficient subjects compared to controls. Early diagnosis of neuropathy due to Vitamin B12 deficiency is important, because a myriad of adverse outcomes that can be progressive and irreversible neurological abnormalities can be averted with substitutive treatment.

**Key words** : Vitamin B 12 deficiency, Nerve conduction.

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

**Introduction**

Vitamin B-12 deficiency is often considered synonymous with pernicious anemia, a rare condition manifesting as megaloblastic erythropoiesis, hyperhomocystinemia, methylmalonic acidemia, and neurological syndromes (subacute combined degeneration of cord, peripheral neuropathy, dementia and depression) [1]. The primary defect in this condition is a near-total block in the gastrointestinal absorption of vitamin B-12 which requires high-dose parenteral treatment. In developing countries such as India, inadequate dietary intake of B-12 due to socio-cultural factors leads to widely prevalent asymptomatic low B-12 status. In this scenario, lower doses of oral B-12 may be effective, safer and more affordable. In the last few decades it has become increasingly apparent that inadequate dietary intake of vitamin B-12 is a widely prevalent cause of low vitamin B-12 status in many populations [2]. This predominantly affects vegetarians who do not eat adequate amounts of animal origin foods (milk, eggs, fish, chicken and meat) and lower socioeconomic groups who cannot afford them and vegans. Small doses of oral vitamin B-12 may be adequate to improve vitamin status in these individuals [3,4]. There is sparse information on this issue. There are many different causes of B12 deficiency in older adults. More than half of older adults with B12 deficiency have food-cobalamin malabsorption, defined as impaired digestion and

absorption of protein-bound B12 [5,6,7]. Other causes of low or deficient B12 are insufficient intake either from diet or supplements, pernicious anemia, gastric surgery, gastrointestinal disease, and certain medications (e.g., proton-pump inhibitors, metformin)

Vitamin B12 deficiency is clinically recognized to be associated with neurological disorders, such as dementia, cognitive impairment, and depression [8,9]. B12 deficiency may cause demyelination of nerves in the peripheral and central nervous system and has been associated with peripheral neuropathy, loss of sensation in peripheral nerves, and weakness in lower extremities in older adults [10,11,12]. In particular, vitamin B12 deficiency is associated with large fiber (type A) neuropathy; type A nerve fibers act as both sensory and motor fiber [13]. Thus, vitamin B12 may be associated with both sensory and motor peripheral nerve function. The objective of the present study was to evaluate the association between vitamin B12 status and peripheral nerve conduction parameters in middle aged adults.

**Material and Methods**

30 middle adults (both males and females) aged between 28 to 45 years were recruited after obtaining written informed consent. None of the 30 control subjects reported any symptoms suggestive of peripheral neuropathy. Subjects who underwent surgery or history of fracture or scars at the sites of

stimulation were excluded. 30 subjects with symptoms of vitamin B12 deficiency were included as patients. Following recruitment, each subject underwent anthropometry, nerve conduction assessment, vitamin B12 estimation and complete hemogram. Vitamin B12 was determined.

**Nerve Conduction Study (NCS):** Nerve conduction study was done using the AD instrument –Nicolet. For motor & sensory nerve study, output range was 20mA, max repeat rate 1Hz, pulse width 0.05ms stimulation with current ranging between 10-20mA was applied with increasing strength until desired response was obtained. Sensory nerves tested were Median, and Ulnar nerve.

**Stimulation and Recording Sites of Sensory Nerves:**

**Sensory Nerve :** Median, **Method of stimulation:** Orthodromic, **Stimulation site :** Index finger **Recording site :** Middle of the wrist .

**Sensory Nerve :** Ulnar , **Method of stimulation:** Orthodromic , **Stimulation site :** Little finger , **Recording site :** Medial wrist

**Sensory nerve conduction study was done:**

**Parameters Assessed:**

1. Sensory nerve action potential (SNAP) Amplitude.

2. Sensory nerve conduction velocity.

**Statistical Analysis:** Statistical analysis was done using SPSS version 18 (software statistical package social science). Students unpaired t test was used to compare Nerve conduction parameters between the study and control groups . p value was calculated.

P Value <0.001 - Highly Significant.

P Value <0.01 – Significant.

P Value >0.01 - Not Significant.

## Results

**Table 1: Age and Vitamin B12 Levels**

	Patients n=30 .Mean± SD	Control n=30. Mean± SD	P value
Age in years	42.64±12.06	40.24±10.40	>0.001
Vitamin B12 levels pmol/L.	170.74±24.36	330.42±16.25	<0.001

**Table 2: Nerve conduction values in vitamin B12 deficient subjects**

Nerve conduction parameters	Patients n=30 .Mean±SD	Control n=30. Mean± SD	P value
<b>Median nerve</b>			
SNCV(m/s)	40.18±4.18	62.14±3.28	<0.001
AMP(ms)	16.42±4.84	20.48±4.62	<0.01
<b>Ulnar nerve</b>			
SNCV(m/s)	44.24±4.68	60.28±3.98	<0.001
AMP(ms)	16.24±4.26	21.68±3.69	<0.01

**AMP -amplitude, SNCV-sensory nerve conduction velocity.**

Table 1 shows low Vitamin B12 levels in patients which was 170.74±24.36 pmol/L(mean±Sd)

Table-2 shows Comparison of sensory nerve parameters in Control Group and Vitamin B12 deficient subjects , there is highly significant slowing of sensory nerves median & ulnar (p<0.001) and significant decrease in amplitude of Median nerve and ulnar nerve (p<0.01) in Vitamin B12 deficient subjects compared to controls.

## Discussion

Vitamin B12 deficiency can occur as a result of malabsorption, gastrointestinal surgery, drugs, parasitic diseases, autoimmune diseases and genetic defects [14]. Low vitamin B-12 intake may lead to decreased bioavailability and functional deficiency of cobalamin. Although early noticeable symptoms of vitamin B-12 deficiency are nonspecific (unusual fatigue, digestion problems, frequent upper respiratory infections), the best-known clinical manifestations of cobalamin malabsorption are hematologic (pernicious anemia) and neurologic symptoms [15]. Though the pathophysiological

mechanism of the neurological damage is not clear vitamin B12 deficiency leads to different pathological mechanisms in the central and peripheral nervous system [16]. Vitamin B12 deficiency should be considered in the differential diagnosis of all spinal cord, peripheral nerve, and neuropsychiatric disorders. Vitamin B12 replacement should not be withheld from patients with borderline vitamin B12 levels, since the consequences of allowing myelopathy, neuropathy dementia, and mental disorders to worsen clearly outweigh any disadvantage of therapy [17]. The best-known neurologic manifestation of vitamin B12 deficiency is subacute combined degeneration. Isolated neuropathy or myelopathy may occur independently, but often appear concurrently. Peripheral neuropathy is frequently observed in symptomatic patients with vitamin B12 deficiency [18]. Lack of vitamin B12 was suggested to effect sensory nerves primarily. Nerve conduction studies of vitamin B12 deficient patients with clinically apparent neuropathy have shown decreased amplitudes of mostly sensory nerves in a majority of cases [19]. The other studies did not observe axonal changes but observed primary demyelinating sensory neuropathy [20] whereas some studies have

shown that motor nerves could also be effected due to vitamin B12 deficiency [21]. shows In our Study Comparison of sensory nerve parameters in Control Group and Vitamin B12 deficient subjects , there is highly significant slowing of sensory nerves median & ulnar ( $p < 0.001$ ) and significant decrease in amplitude of Median nerve and ulnar nerve ( $p < 0.01$ ) in Vitamin B12 deficient subjects compared to controls.

### Conclusion

In conclusion, nerve conduction study may show pathological findings in neurologically symptomatic patients with vitamin B12 deficiency and therefore nerve conduction study is a method for detection of early peripheral neuropathy in vitamin B12 deficient patients. Early diagnosis is important, because a myriad of adverse outcomes that can be progressive and irreversible neurological abnormalities can be averted with substitutive treatment

### References

1. Stabler SP. Clinical practice. Vitamin B12 deficiency. *N Engl J Med.* 2013; 368: 149–160. <https://doi.org/10.1056/NEJMc1113996> PMID: 23301732
2. Green R, Allen LH, Bjørke-Monsen A-L, Brito A, Gueant J-L, Miller JW, et al. Vitamin B12 deficiency. *Nat Rev Dis Primers.* 2017; 3: 170–40. Chan CQH, Low LL, Lee KH. Oral Vitamin B12 Replacement for the Treatment of Pernicious Anemia. *Front Med.* 2016; 3: 38. <https://doi.org/10.3389/fmed.2016.00038> PMID: 27602354
3. Vidal-Alaball J, Butler C, Cannings-John R, Goringe A, Hood K, McCaddon A, et al. Oral vitamin B12 versus intramuscular vitamin B12 for vitamin B12 deficiency. *Cochrane Database Syst Rev.* 2005; CD004655. <https://doi.org/10.1002/14651858.CD004655.pub2> PMID: 16034940
4. Sivaprasad M, Shalini T, Balakrishna N, Sudarshan M, Lopamudra P, Suryanarayana P, et al. Status of Vitamin B12 and Folate among the Urban Adult Population in South India. *Ann Nutr Metab.* 2016; 68: 94–102. <https://doi.org/10.1159/000442677> PMID: 26667891
5. Andres E, Loukili NH, Noel E, et al. Vitamin B12 (cobalamin) deficiency in elderly patients. *CMAJ.* 2004; 171: 251–259 [PubMed: 15289425]
6. Andres E, Affenberger S, Vinzio S, et al. Food-cobalamin malabsorption in elderly patients: Clinical manifestations and treatment. *Am J Med.* 2005; 118: 1154–1159. [PubMed: 16194648]
7. Allen LH. How common is vitamin B-12 deficiency? *Am J Clin Nutr.* 2009; 89: 693S–696S. [PubMed: 19116323]
8. Bernard MA, Nakonezny PA, Kashner TM. The effect of vitamin B12 deficiency on older veterans and its relationship to health. *J Am Geriatr Soc.* 1998; 46: 1199–1206. [PubMed: 9777900]
9. Johnson MA, Hawthorne NA, Brackett WR, et al. Hyperhomocysteinemia and vitamin B-12 deficiency in elderly using Title IIIc nutrition services. *Am J Clin Nutr.* 2003; 77: 211–220. [PubMed: 12499344]
10. Klee GG. Cobalamin and folate evaluation: Measurement of methylmalonic acid and homocysteine vs vitamin B(12) and folate. *Clin Chem* 2000; 46: 1277–1283. [PubMed: 10926922]
11. Saperstein DS, Barohn RJ. Peripheral neuropathy due to cobalamin deficiency. *Curr Treat Options Neurol.* 2002; 4: 197–201. [PubMed: 11931726]
12. Mold JW, Vesely SK, Keyl BA, et al. The prevalence, predictors, and consequences of peripheral sensory neuropathy in older patients. *J Am Board Fam Pract.* 2004; 17: 309–318. [PubMed: 15355943]
13. Kumar S. Vitamin B12 deficiency presenting with an acute reversible extrapyramidal syndrome. *Neurol India.* 2004; 52: 507–509. [PubMed: 15626849]
14. Scalabrino G. Cobalamin (Vitamin B12) in subacute combined degeneration and beyond: traditional interpretation and novel theories. *Exp Neurol* 2005; 192: 463–479.
15. Elmadfa I, Singer I. Vitamin-B12 and homocysteine status among vegetarians: a global perspective. *Am J Clin Nutr* 2009; 89(5): 1693–1698
16. Puri V, Chaudhry N, Goel S, Gulati P, Nehru R, Chowdhury D. Vitamin B12 deficiency: a clinical and electrophysiological profile. *Electromyogr Clin Neurophysiol.* 2005; 45(5): 273–284.
17. Petchkrua W, Little JW, Burns SP, Stiens SA, James JJ. Vitamin B12 deficiency in spinal cord injury: a retrospective study. *J Spinal Cord Med.* 2003; 26(2): 116–121. Comment in: *J Spinal Cord Med.* 2003; 26(2): 122–123.
18. Turgut B, Turgut N, Akpınar S, Balçık K, Pamuk GE, Tekgündüz E, Demir M. Dorsal nerve conduction study in vitamin B(12) deficiency with megaloblastic anemia. *J Peripher Nerv Syst* 2006; 11(3): 247–252.
19. McCombe PA, McLeod JG. The peripheral neuropathy of vitamin B12 deficiency. *J Neurol Sci* 1984; 66(1): 117–126.
20. Steiner I, Kidron D, Soffer D, Wiraguin I, Abramsky O. Sensory peripheral neuropathy of vitamin B12 deficiency: a primary demyelinating disease. *J Neurol* 1988; 235: 163–164.
21. Sakly G, Helara O, Trabeisi A, Dogui M. Reversible peripheral neuropathy induced by vitamin B12 deficiency. *Neurophysiol Clin* 2005; 35(56): 149–153.