

**Clinical Research on Non-Venereal Genital Dermatoses in Adult Males at Bihar State****Sudhir Kumar Shreevastav**

Associate Professor, Department of Skin &amp; V D, Lord Buddha Koshi Medical College and Hospital, Saharsa, Bihar, India

Received: 13-04-2023 / Revised 11-05-2023 / Accepted 28-06-2023

Corresponding author: Dr. Sudhir Kumar Shreevastav

Conflict of interest: Nil

**Abstract:****Aim:** A clinical study on non-venereal genital dermatoses in adult males at a tertiary care center.**Methods:** The prospective analytical study was conducted in the Department of skin and VD for one year. A total of consecutive 100 male patients with genital lesions of non-venereal origin were included in this study. All male patients >18 years of age who presented with genital complaints were screened for non-venereal dermatoses. The external genitalia was examined, and findings were noted. A detailed physical examination was done to see any associated lesions elsewhere in the body. Investigations such as Gram-stain, KOH mount, venereal disease research laboratory test, HIV test and histopathological examination were done as and when required establishing the diagnosis.**Results:** Most patients belong to the age group of 20-30 years (45%), followed by the age group of 30-40 years (22%). 75 patients (75%) were from the urban area while 25 patients (25%) belong to rural background. 57 (57%) patients were married and the remaining 43 (43%) patients were unmarried. Scrotum was involved in 62% and penis in 30% while both scrotum and penis were affected in 8% cases. 28% patients were farmers while 18% patients were students. A total of 16 types of non-venereal dermatoses were noted in this study. The most common disorder was vitiligo present in 20 cases, followed by pearly penile papule, which accounted for 15 cases. The other disorder encountered included fixed drug eruption (FDE) in 13; scabies in 8, scrotal dermatitis and lichen planus in 8 cases each etc.**Conclusion:** Contrary to normal belief all the lesions on genitalia are not sexually transmitted. It is very important to distinguish between venereal and non-venereal genital dermatoses, as these non-venereal disorders are a considerable concern to patients causing mental distress and feeling of guilt.**Keywords:** Genital Dermatoses, Non-Venereal, Disorders.

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

**Introduction**

Male genitalia, being a sexual organ, harbor many sexually transmitted infections (STIs). The external genitalia are a site for various inflammatory diseases which are completely unrelated to sexual transmission, such as psoriasis and zoon's balanitis which is completely or predominantly confined to this area and yet not sexually transmitted. These all-genital dermatoses which are not transmitted sexually are called as non-venereal genital dermatoses. [1] Patients with genital dermatoses are anxious and apprehensive at presentation as they believe them to be the manifestations of sexually transmitted diseases (STDs). In contrast to this popular belief, not all genital dermatoses are sexually transmitted. The diseases which are not sexually transmitted are referred to as non-venereal dermatoses. Fitzpatrick and Gentry [2] classified these dermatoses based on pathogenesis as 1) Benign abnormalities 2) congenital abnormalities 3)

Trauma and artefacts 4) Inflammatory diseases 5) Non venereal infections and infestations 6) Benign tumours 7) Premalignant lesions 8) Malignant lesions 9) Miscellaneous lesions. Genital dermatoses pose a difficulty in diagnosis as the morphology is modified by the special environment of the genitalia like heat, friction and occlusion. Non-venereal genital dermatoses may not be restricted to the genitalia alone but can also affect other areas of the body. So, examining the extra genital sites aids in the diagnosis. Even with benign lesions some patients develop venerophobia, cancer phobia. Hence it is important to be aware of these conditions and to differentiate them from venereal disease. Explaining the true and benign nature of these lesions will remove this fear. Because of the stigma associated with genital lesions, most of these patients do not approach the medical fraternity unless the disease burden is unbearable which in case of malig-

nant lesions can endanger their lives. [3] Clinicians should have an open mind to look for these genital lesions so that patients feel confident to seek medical help. A comprehensive understanding of the various presentations, their causes and appropriate treatment options is essential to effectively manage these non-venereal dermatoses and allay the associated anxiety. The aim of our study was to determine the clinical and etiological factors, various patterns of presentation of non-venereal dermatoses in male genitalia and to assess which dermatoses have a predilection for exclusive genital involvement and which occur as a part of the generalised skin involvement.


**Material and Methods**

The prospective analytical study was conducted in the Department of Skin & V D, Lord Buddha Koshi Medical College and Hospital, Saharsa, Bihar, India for one year. A total of consecutive 100 male patients with genital lesions of non-venereal origin were included in this study. All male patients >20 years of age who presented with genital complaints were screened for non-venereal dermatoses. Informed consent was obtained. A detailed history

including demographic data, chief complaints related to skin, onset and duration of disease and associated medical, or skin disorders was elicited and recorded. History of sexual exposure was also recorded. Cases having any venereal diseases were excluded from the study. The external genitalia were examined, and findings were noted. A detailed physical examination was done to see any associated lesions elsewhere in the body. Investigations such as Gram-stain, KOH mount, venereal disease research laboratory test, HIV test and histopathological examination were done as and when required establishing the diagnosis. A proforma was prepared to record the relevant details of patient, examination, investigations and diagnosis.


**Results**

A total of 100 male patients with non-venereal dermatoses of external genitalia were included in the study. The age of the patients ranged from 20 years to 60 years, with the mean age of 31.7 years. Most patients belong to the age group of 20-30 years (45%), followed by the age group of 30-40 years (22%).




75 patients (75%) were from the urban area while 25 patients (25%) belong to rural background.


RESIDENTIAL STATUS OF PATIENT	NUMBER OF PATIENTS
URBAN	75
RURAL	25



57 (57%) patients were married and the remaining 43 (43%) patients were unmarried.




Scrotum was involved in 62% and penis in 30% while both scrotum and penis were affected in 8% cases.



**Occupation**

28% patients were farmers while 18% patients were students.



A total of 16 types of non-venereal dermatoses were noted in this study [Table 1]. The most common disorder was vitiligo present in 20 cases, followed by pearly penile papule, which accounted for 15 cases. The other disorder encountered included fixed drug eruption (FDE) in 13; scabies in 8, scrotal dermatitis and lichen planus in 8 cases each etc.

**Table 1: Genital dermatoses**

Genital dermatoses	Number (n=100)	%age
Vitiligo	20	20
Pearly penile papule	15	15
Psoriasis	3	3
Scrotal dermatitis	8	8
Squamous cell carcinoma	1	1
Lichen planus	8	8
Dermatophytosis	6	6
Granuloma annul are	1	1
Scabies	11	11
Lichen nitidus	1	1
Fixed drug eruption	13	13
Sebaceous cyst	6	6
Lichen sclerosus	3	3
Lymphangiectasia scrotum	2	2
Zoon's balanitis	1	1
Papulo-necrotic tuberculid	1	1

The common presenting features were itchy genitalia, depigmentation. Other complaints were pain, burning sensation, redness, exfoliation of the skin, raised lesions over the skin, oozing, ulceration erosions and thickening of the skin. Some patients had more than one complaint.

### Discussion

As venereal dermatoses are of primary concern to the patient and causes mental stress and guilt feeling among patients, it is, therefore, utmost important to distinguish between venereal and non-venereal dermatoses. Male patients with non-venereal dermatoses usually present to genitor-urinary experts or physicians, where the training and expertise are not oriented to adequate dermatological diagnosis and treatment. [4] Disorders of genitalia have proved confusing to various specialists involved in the diagnosis and treatment. The problem is confounded by the fact that the normal characteristics of common diseases at flexural sites are lost or modified, making the diagnosis difficult for even an experienced dermatologist.

The non-venereal dermatoses of male external genitalia include wide spectrum of disease with varied etiology. [5] There are very few comprehensive studies on the pattern of non-venereal dermatoses in males from our country. [6,7] Also, our study is first of its kind from this part of the country. Acharya *et al.* [6] had done a study of 200 patients with genital lesions of non-venereal origin. Karthikeyan *et al.* [7] had done a study on the pattern of non-venereal dermatoses of male external genitalia from South India. Khoo and Cheong [8] had done a similar study on male patients at Singapore.

The age ranged from 20 to 60 years in the present study with the mean age of 31.7 years whereas the age ranged from 9 to 70 years with a mean age 33.7 years in a study by Karthikeyan *et al.* [7]

Most of the patients belong to the age group of 20–30 years (45%) in the present study which is similar to Karthikeyan *et al.* [7]

A total of 16 different non-venereal dermatoses were observed in this study [Table 1]. Karthikeyan *et al.* [7] had 25 different non-venereal dermatoses in their study.

The most common disorder was vitiligo present in 20 cases, followed by pearly penile papule, which accounted for 15 cases. The other disorder encountered included fixed drug eruption (FDE) in 13; scabies in 8, scrotal dermatitis and lichen planus in 8 cases each etc.

The study by Acharya *et al.* [6] reported infections as commonest disorder contributing 40% cases. Genital vitiligo as most common disorder (16%) was observed in an another study, which is almost similar to our study. [7] Khoo and Cheong [8] had 14.3% pearly penile papule as most common non-venereal dermatoses, which are similar to this study (16%).

75 patients (75%) were from the urban area while 25 patients (25%) belong to rural background. 57 (57%) patients were married and the remaining 43 (43%) patients were unmarried. Scrotum was involved in 62% and penis in 30% while both scrotum and penis were affected in 8% cases.

Genital vitiligo could be an exclusive finding, or it can be associated with generalized vitiligo. Genital

vitiligo accounted for 20% cases as commonest disorder in our study and is seen in all age group from young adult to older age group. This is in contrast with the study conducted by Karthikeyan *et al.* [7] where the entire patients with vitiligo were in older age group. Ten patients in our study had associated vitiligo elsewhere while eight patients had only genital vitiligo. Duration of illness ranged from 3 months to 8 years.

Pearly penile papule is a common disorder found in up to 50% of men. [9] They were present in 15% cases in our study, which is almost similar to the study conducted by Khoo and Cheong [8] They are frequently mistaken as warts and misdiagnosed as Tyson's gland or ectopic sebaceous gland of Fordyce. [5] All the patients with pearly penile papule came to visit OPD in apprehension of some venereal disease. They were counseled about the benign nature of the disease.

Fixed drug eruptions were observed in 13% of cases in our study as third most common disorder. This is in contrast with Karthikeyan *et al.*, [7] where only 3 cases had FDE and all of them because of cotrimoxazole. In our study, various drugs were implicated such as, nonsteroidal anti-inflammatory drugs, sulphonamides, ornidazole, fluconazole, ampicillin, etc., Half of our patients with FDE had oral involvement also.

Acharya *et al.* [6] in their study recorded scabies as most common non-venereal dermatoses accounting for 30 cases (15%), while it was present in only 11% cases in our study. This may be due to lesser prevalence of scabies in this population.

Lichen planus was present in 8% cases in our study that is in contrast with Puri and Puri [10] where it was seen in only 6.6% (3) cases and Karthikeyan *et al.* [7] where it was seen in only 1 case. Four of our cases had involvement of the oral mucosa also Itching particularly around scrotum is a common presenting problem. Contributory factors include, tight clothing, friction, maceration, atopy, over-washing, use of various toiletries, topical medicaments and indigenous preparations. [11-13] Scrotal dermatitis accounted for 8% cases in our study inclusive of allergic contact dermatitis, irritant contact dermatitis. Most of the patients were from the rural background. Acharya *et al.* [6] did not report any case while Karthikeyan *et al.* [7] had 13% cases of scrotal dermatitis.

Sebaceous cysts of scrotum were observed in 6% cases in our study, while it was second most common finding (14%) by Karthikeyan *et al.* [7] They were observed in only 3.7% cases by Khoo and Cheong [8] All of our cases were asymptomatic and from younger age group.

Dermatophytic infection was present in 6% cases in our study as scaly pruritic plaques over scrotum. All of them were confirmed by KOH mount.

Lichen sclerosus (LS) is chronic inflammatory dermatoses which are associated with substantial discomfort and morbidity with an unknown etiology. [14] LS was observed in 3 cases in our study, while it was seen in only 2 cases by Karthikeyan *et al.* [7] All 3 cases had phimosis and were advised circumcision. Duration of illness ranged from 6 months to 3 years. Clinical findings in cases of LS in this study were found to be in concordance with the literature review.

Around 2% of the world population has psoriasis, but it is possible that many more could have anogenital psoriasis at some time. [15] Also, psoriasis of anogenital region can present alone. Genital appearance could be challenging to interpret, especially in uncircumscribed individuals because a mucosal site is affected rather than keratinized skin. [5] Psoriasis was encountered in 3% cases in our study. Karthikeyan *et al.* [7] reported a solitary case of psoriasis of glans penis while Acharya *et al.* [6] reported 5 cases of psoriasis over genitalia. All of our cases had classical lesions of psoriasis elsewhere.

Zoon's balanitis or plasma cell balanitis was observed in 1% cases in this study that had not been reported by Acharya *et al.*, [6] Khoo and Cheong [8] Karthikeyan *et al.* [7] It is a disorder of middle and older age in uncircumcised male, the etiology remains unknown. [15]

### Conclusion

Contrary to normal belief all the lesions on genitalia are not sexually transmitted. It is very important to distinguish between venereal and non-venereal genital dermatoses, as these non-venereal disorders are a considerable concern to patients causing mental distress and feeling of guilt. This study was quiet useful in understanding the epidemiological, clinical and etiological characteristics of various non-venereal genital dermatoses. The most common etiological diagnosis in our study was vitiligo.

### Reference

1. Andreassi L, Bilenchi R. Non-infectious inflammatory genital lesions. *Clin Dermatol* 2014; 32:307-14.
2. Fitzpatrick JA, Gentry RM. Non-venereal diseases of male external genitalia. In: Moschella SL, Hurley HJ, editors. *Dermatology*. 3rd ed. I. Philadelphia: WB Saunders Company; 1992: 1008-15.
3. Neerja Puri, Asha Puri: A study on non-venereal genital dermatoses in north India. *Our Dermatol Online*. 2013;4(1): 304-7

4. Hillman RJ, Walker MM, Harris JR, Taylor-Robinson D. Penile dermatoses: A clinical and histopathological study. *Genitourin Med* 1992; 68:166-9.
5. Bunker CB, Neill SM. The genital, perianal and umbilical regions. In: Burns T, Breathnach S, Cox N, Griffiths C, editors. *Rook's Textbook of Dermatology*. 7th ed. Oxford: Blackwell Science; 2004; 68.1-104.
6. Acharya KM, Ranpara H, Sakhia JJ, Kaur C. A study of 200 cases of genital lesions of non-venereal origin. *Indian J Dermatol Venereol Leprol* 1998; 64:68-70.
7. Karthikeyan KE, Jaishankar TJ, Thappa DM. Non-venereal dermatoses of male genital region-prevalence and pattern in a referral centre in South India. *Indian J Dermatol* 2001; 46:18-22.
8. Khoo LS, Cheong WK. Common genital dermatoses in male patients attending a public sexually transmitted disease clinic in Singapore. *Ann Acad Med Singapore* 1995;24:505-9.
9. Sonnex C, Dockerty WG. Pearly penile papules: A common cause of concern. *Int J STD AIDS* 1999; 10:726-7.
10. Puri N, Puri A. A study of non-venereal genital dermatoses in North India. *Our Dermatol Online* 2013; 4:304-7.
11. Ramam M, Khaitan BK, Singh MK, Gupta SD. Frictional sweat dermatitis. *Contact Dermatitis* 1998; 38:49.
12. Bauer A, Geier J, Elsner P. Allergic contact dermatitis in patients with anogenital complaints. *J Reprod Med* 2000; 45:649-54.
13. Hindson TC. Studies in contact dermatitis. *Trans St Johns Hosp Dermatol Soc* 1966; 52:1-9.
14. Powell JJ, Wojnarowska F. Lichen sclerosus. *Lancet* 1999; 353:1777-83.
15. Zoon JJ. Balanitis and vulvitis plasma cellularis. *Dermatologica* 1955; 111:157.