

A Hospital Based Epidemiological Study to Assess the Spectrum of Congenital Heart Diseases in a Tertiary Care Centre

Priya Verma¹, Gopal Shankar Sahni²

¹Assistant Professor, Department of Pediatrics, Shri Krishna Medical College and Hospital, Muzaffarpur, Bihar, India

²Associate Professor & HOD, Department of Pediatrics, Shri Krishna Medical College and Hospital, Muzaffarpur, Bihar, India

Received 10-12-2023 / Revised: 27-12-2023 / Accepted: 13-01-2024

Corresponding Author: Dr. Priya Verma

Conflict of interest: Nil

Abstract

Aim: The aim of the present study was to assess the spectrum of congenital heart diseases in a tertiary care centre in Bihar Region.

Methods: The present study was conducted in the Department of Pediatrics, Shri Krishna Medical College and Hospital, Muzaffarpur, Bihar, India and 200 patients were studied.

Results: We studied 200 patients in age group 0 to 5 years with clinical suspicion of CHD out of which 116 were male and 84 were female and maximum patients belonged to 1-12 months of age. The difficulty in breathing emerged as the most common presenting complaint, followed by feeding difficulty. The cases with VSD were found to be 32%, ASD were 30%, PDA were 8%, TOF were 5%, 13% cases were combination of multiple defects.

Conclusion: We concluded that VSD 32% was the commonest heart disease, followed by ASD 30%, congenital heart diseases showed male preponderance. VSD was the commonest acyanotic heart disease and TOF was the commonest cyanotic heart disease. The majority of patients were seen in the age group 1 to 12 months of age. Difficulty in breathing was the most common presenting complaint followed by feeding difficulty.

Keywords: Congenital Heart Disease, Tertiary Care Centre, Bihar.

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

Introduction

Congenital heart disease (CHD) is an abnormality in the structure or function of the cardio-circulatory system present at birth, even if diagnosed later. [1] It varies in severity, occurring from communications between cavities that spontaneously regress to major malformations that even require several procedures, by catheterization or by surgical means. It may result in mortality in the intrauterine, childhood, or adulthood period. [2] CHD is the most common (28%) major congenital anomaly and thus signifies a major global health problem. [3] The birth prevalence of CHD varies among studies worldwide and is mostly reported between 8 and 12 per 1000. [4,5] With a prevalence of 9 per 1000, approximately 1.35 million newborns are born with CHD every year globally. [5]

The prevalence of CHD in India in recent studies reports an increased prevalence of 8.5-13.6 per 1000 children. [6,7] In India, CHD contributes significantly to infant mortality (10%). [8] The common morbidities reported in children with

CHD include developmental delay and cognitive deficits (20-30%). [9,10] A significant proportion of children born with CHD may lead a normal, productive life if diagnosed early and appropriate medical/surgical intervention is instituted. Thus, early detection of CHD and timely intervention are important for a better outcome in these children. [11,12]

Congenital heart disease is the most common congenital problem effecting nearly 25% of all children with congenital malformation [13] with an incidence of almost 8/1000 live births. [14] Early diagnosis and prompt treatment has great implication on prognosis and can result in significant decrease in morbidity and mortality. The relatively high birth rate in India, an average of 150,000 children are born with congenital abnormalities with 50,000 requiring surgery in the first year of life and only one thousand cardiac surgeries are done in the early infancy. [15]

Despite vast improvements in medical field, congenital heart disease is still one of the leading

cause of death and can present in different age groups from birth to adolescence. [16] The presentation of children with CHD can be very variable; they may be asymptomatic and discovered only accidentally when a murmur is noted during check-up for an unrelated illness or routine neonatal checkup, [17] or may present with symptoms like cyanosis, clubbing of fingers nails to full blown congestive heart failure. [16,17]

The aim of the present study was to assess the spectrum of congenital heart diseases in a tertiary care centre in Bihar Region.

Materials and Methods

The present study was conducted in the Department of Pediatrics, Shri Krishna Medical College and Hospital, Muzaffarpur, Bihar, India for one year and 200 patients were studied.

Inclusion Criteria: All children <5 years with clinically suspected congenital heart disease were included.

Exclusion Criteria: Patients with previously diagnosed heart diseases were excluded.

Procedure: In a period of 12 months, any patient <5 years with strong clinical suspicion of congenital heart disease attending inpatient and outpatient department of Shri Krishna Medical College and Hospital, Muzaffarpur, Bihar, India (including children's emergency and SNCU) was selected for study.

CHD was suspected based on the clinical features and an abnormal pulse oximetry value (<90%). A detailed history of patients was taken from their reliable informant with special emphasis on complaints suggestive of any cardiac illness. Developmental milestones and anthropometry was also assessed, general and systemic examination of patients was done and patients were also screened for presence of any other anomaly. All this was done using a pre-structured proforma. These patients were screened using NADAS criteria and were further classified into NADAS positive and negative. The confirmation of presence congenital heart disease was done using echocardiography. All those patients who were not found to have congenital heart disease on echocardiography were excluded.

Results

Table 1: Age and sex distribution of congenital heart diseases

Age	Male	Female	Total	
	No.	No.	No.	%
0 to 1 month	40	36	76	38
1 to 12 months	60	32	92	46
1 to 5 years	16	16	32	16
Total	116	84	200	100

We studied 200 patients in age group 0 to 5 years with clinical suspicion of CHD out of which 116 were male and 84 were female and maximum patients belonged to 1-12 months of age.

Table 2: Clinical profile of patients presenting with congenital heart diseases

Symptoms	N (%)
Bluish discoloration of body	50 (25)
Recurrent respiratory infections	48 (24)
Palpitations/chest pulsations	22 (11)
Swelling of face and limbs	4 (2)
Forehead sweating	48 (24)
Impaired consciousness/unresponsiveness	14 (7)
Suck rest suck cycle/feeding difficulty	58 (29)
Difficulty in breathing	192 (96)
Any other symptom	24 (12)

The difficulty in breathing emerged as the most common presenting complaint, followed by feeding difficulty.

Table 3: Spectrum of congenital heart diseases in age group 0 to 5 years

S. No.	Echo status	Male	Female	Total	
				No.	%
1.	VSD	44	20	64	32
2.	ASD	28	32	60	30
3.	PDA	4	12	16	8
4.	TOF	8	2	10	5
5.	Complex (others)	18	8	26	13
6.	TAPVC	4	0	4	2

7.	AV canal defect	2	2	4	2
8.	Single ventricle	2	0	2	1
9.	HOCM	2	0	2	1
10.	TGA	4	0	4	2
11.	Pentalogy of fallot	0	2	2	1
12.	Tricuspid atresia	0	4	4	2
13.	Hypoplastic left heart syndrome	0	2	2	1

The cases with VSD were found to be 32%, ASD were 30%, PDA were 8%, TOF were 5%, 13% cases were combination of multiple defects.

Discussion

Congenital heart defects are the most common type of congenital anomaly, which constitutes an important group of pediatric illness and major cause of childhood mortality and morbidity. [18] Congenital malformations of the heart and circulation are not fixed anatomic defects that appear at birth but instead are anomalies in flux that originate in the early embryo, evolve during gestation, survive the dramatic circulatory alterations at birth, and change considerably during extra uterine life. [19]

We studied 200 patients in age group 0 to 5 years with clinical suspicion of CHD out of which 116 were male and 84 were female and maximum patients belonged to 1-12 months of age. The difficulty in breathing emerged as the most common presenting complaint, followed by feeding difficulty. The cases with VSD were found to be 32%, ASD were 30%, PDA were 8%, TOF were 5%, 13% cases were combination of multiple defects. This correlated with study done by Doshi et al [20] (2022) which included children between the age group of 0-18 years.

A study done by Kapoor et al [21] gives prevalence of 26.4/1000 patients. VSD was the commonest lesion (21.3%), followed by ASD (18.9%) and PDA (14.6%). Tetralogy of Fallot was the commonest cyanotic heart disease (4.6%), which correlates with our study. The prevalence in our study was also similar to the study done by Bibi et al [22] (2018) which showed a male preponderance with 57 (64%) male patients as compared to 32 (36%) female patients. Ventricular septal defect (VSD) was the commonest cardiac lesion being present in 34 (38.2%) patients. The prevalence also correlates with the study done by Bhardwaj et al (2014) and Samanek et al (1989) with both having VSD as the commonest congenital heart disease followed by ASD. [23,24]

The study done by Khalil et al [25] gives the incidence of CHD per 1000 live births. They studied 10964 live births and observed the incidence of 3.9/1000 live births. Patent ductus arteriosus (41.9%) and ventricular septal defects (VSD) (34.9%), were the commonest lesions with

the incidence of 1.6 and 1.4/1000 live births, respectively.

Conclusion

We concluded that VSD 32% was the commonest heart disease, followed by ASD 30%, congenital heart diseases showed male preponderance. VSD was the commonest acyanotic heart disease and TOF was the commonest cyanotic heart disease. The majority of patients were seen in the age group 1 to 12 months of age. Difficulty in breathing was the most common presenting complaint followed by feeding difficulty.

References

1. Bonow RO, Mann DL, Zipes DP, Libby P. Braunwald's heart disease e-book: A textbook of cardiovascular medicine. Elsevier Health Sciences; 2011 Feb 25.
2. Go AS, Mozaffarian D, Roger VL, Benjamin EJ, Berry JD, Borden WB, Bravata DM, Dai S, Ford ES, Fox CS, Franco S, Fullerton HJ, Gillespie C, Hailpern SM, Heit JA, Howard VJ, Huffman MD, Kissela BM, Kittner SJ, Lackland DT, Lichtman JH, Lisabeth LD, Magid D, Marcus GM, Marelli A, Matchar DB, McGuire DK, Mohler ER, Moy CS, Mussolino ME, Nichol G, Paynter NP, Schreiner PJ, Sorlie PD, Stein J, Turan TN, Virani SS, Wong ND, Wood D, Turner MB; American Heart Association Statistics Committee and Stroke Statistics Subcommittee. Heart disease and stroke statistics--2013 update: a report from the American Heart Association. *Circulation*. 2013 Jan 1;127 (1):e6-e245.
3. Dolk H, Loane M, Garne E; European Surveillance of Congenital Anomalies (EUROCAT) Working Group. Congenital heart defects in Europe: prevalence and perinatal mortality, 2000 to 2005. *Circulation*. 2011 Mar 1;123 (8): 841-9.
4. Hoffman JIe. The global burden of congenital heart disease. *Cardiovasc J Afr*. 2013 May; 24 (4):141-5.
5. van der Linde D, Konings EE, Slager MA, Witsenburg M, Helbing WA, Takkenberg JJ, Roos-Hesselink JW. Birth prevalence of congenital heart disease worldwide: a systematic review and meta-analysis. *J Am Coll Cardiol*. 2011 Nov 15;58(21):2241-7.
6. Bhat NK, Dhar M, Kumar R, Patel A, Rawat A, Kalra BP. Prevalence and pattern of con-

- genital heart disease in Uttarakhand, India. *Indian J Pediatr.* 2013 Apr;80(4):281-5.
7. Smitha R, Karat SC, Narayanappa D, Krishnamurthy B, Prasanth SN, Ramachandra NB. Prevalence of congenital heart diseases in Mysore. *Indian Journal of Human Genetics.* 2006; 12(1):11-6.
 8. Saxena A. Congenital heart disease in India: a status report. *Indian J Pediatr.* 2005 Jul;72(7): 595-8.
 9. Limperopoulos C, Majnemer A, Shevell MI, Rosenblatt B, Rohlicek C, Tchervenkov C. Neurodevelopmental status of newborns and infants with congenital heart defects before and after open heart surgery. *J Pediatr.* 2000 Nov;137(5):638-45.
 10. Shillingford AJ, Glanzman MM, Ittenbach RF, Clancy RR, Gaynor JW, Wernovsky G. Inattention, hyperactivity, and school performance in a population of school-age children with complex congenital heart disease. *Pediatrics.* 2008 Apr;121(4):e759-67.
 11. Brown KL, Ridout DA, Hoskote A, Verhulst L, Ricci M, Bull C. Delayed diagnosis of congenital heart disease worsens preoperative condition and outcome of surgery in neonates. *Heart.* 2006 Sep;92(9):1298-302.
 12. Peterson C, Dawson A, Grosse SD, Riehle-Colarusso T, Olney RS, Tanner JP, Kirby RS, Correia JA, Watkins SM, Cassell CH. Hospitalizations, costs, and mortality among infants with critical congenital heart disease: how important is timely detection? *Birth Defects Res A Clin Mol Teratol.* 2013 Oct;97(10):664-72.
 13. Aburawi EH. The burden of congenital heart disease in Libya. *Libyan J Med.* 2006 Sep 8;1(2):120-2.
 14. Camm AJ, Lüscher TF, Serruys PW, editors. *The ESC textbook of cardiovascular medicine.* Oxford University Press, USA; 2009 Aug 27.
 15. Saxena A. Treatment of rheumatic carditis. *Indian J Pediatr.* 2002 Jun;69(6):513-6.
 16. Wren C, Richmond S, Donaldson L. Presentation of congenital heart disease in infancy: implications for routine examination. *Arch Dis Child Fetal Neonatal Ed.* 1999 Jan;80(1):F49-53.
 17. Kuehl KS, Loffredo CA, Ferencz C. Failure to diagnose congenital heart disease in infancy. *Pediatrics.* 1999 Apr;103(4 Pt 1):743-7.
 18. Bhardwaj R, Rai SK, Yadav AK, Lakhota S, Agrawal D, Kumar A, et al. Epidemiology congenital heart disease in India. *Congenit Heart Dis.* 2015;10(5):437-46.
 19. Sommer RJ, Hijazi ZM, Rhodes JF. Pathophysiology of congenital heart disease in the adult: part I: shunt lesions. *Circulat.* 2008;117(8):1.
 20. Doshi M, Jain H, Saini A. Clinical and anthropological profile of children presenting with features of congenital heart disease at tertiary care centre. *International Journal of Contemporary Pediatrics.* 2022 Jun;9(6):614.
 21. Kapoor R, Gupta S. Prevalence of congenital heart disease, Kanpur, India. *Indian Pediatr.* 2008;45(4):309-11.
 22. Bibi S, Gilani SY, Bibi S. spectrum of congenital heart disease in full term neonates. *J Ayub Med Coll Abbottabad.* 2018;30(1):67-70.
 23. Bhardwaj R, Rai SK, Yadav AK. Epidemiology of congenital heart disease in India. *Congenit Heart Dis.* 2015;10(5):437-46.
 24. Samánek M, Slavík Z, Zborilová B, Hrobovná V, Vorísková M, Skovráněk J. Prevalence, treatment, and outcome of heart disease in live-born children: a prospective analysis of 91,823 live-born children. *Pediatr Cardiol.* 1989;10(4):205-11.
 25. Khalil A, Aggarwal R, Thirupuram S, Arora R. Incidence of congenital heart disease among hospital live births in India. *Indian Pediatr.* 1994;31(5):519-27.