

## Clinical-Epidemiological Assessment of Patients with Acute Kidney Injury Following Acute Gastroenteritis

Gangesh Kumar Gunjan<sup>1</sup>, Umesh Chandra Jha<sup>2</sup>

<sup>1</sup>Senior Resident, Department of General Medicine, Darbhanga Medical College and Hospital, Darbhanga, Bihar, India

<sup>2</sup>Associate Professor and HOD, Department of General Medicine, Darbhanga Medical College and Hospital, Darbhanga, Bihar, India

Received: 05-11-2023 / Revised: 17-12-2023 / Accepted: 28-01-2024

Corresponding Author: Dr. Gangesh Kumar Gunjan

Conflict of interest: Nil

### Abstract

**Aim:** The aim of the present study was to assess the clinical profile of patients with acute kidney injury following acute gastroenteritis.

**Methods:** The Present study was conducted in Department of General Medicine, Darbhanga Medical College and Hospital, Darbhanga, Bihar, India from for 12 months. After applying inclusion and exclusion criteria, total 100 patients of AGE with AKI were considered for this study.

**Results:** Most common age group in this study was age group of 61–70 years (30%), followed by age group of 51–60 years (23%). Mean age of study patients was  $54.6 \pm 12.8$  years. Male patients (65%) were more than female patients (35%). According to clinical presentation most common symptom was loose stools (100%), followed by fever (78%) and vomiting (70%). Other complaints were shortness of breath (20%) and altered sensorium (16%). AKI was staged for severity according to the KDIGO criteria. At the time of diagnosis most patients were in stage 1 (58%), while 31% and 11% were in stage 2 and 3 respectively.

**Conclusion:** Acute kidney injury in patients with acute gastroenteritis had good prognosis if detected earlier. Early recognition of AKI is essential to ensure prompt and appropriate management, and to avoid progression to deadlier stages of the disease.

**Keywords:** Acute Gastroenteritis, Acute Kidney Injury, KDIGO criteria.

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

### Introduction

Acute gastroenteritis (AGE) is characterized by rapid onset of diarrhea, which may be accompanied by vomiting, nausea, or fever, that usually lasts less than 14 days. [1] In the United States (US) alone, approximately 179 million episodes of AGE occur each year, resulting in approximately 600 000 hospitalizations and 5000 deaths. [2] Clinically, gastroenteritis can manifest as mild, moderate, or severe disease with the main symptoms of abdominal pain, diarrhea, nausea, and vomiting. A severe form of AGE includes bloody stools, fever, and sepsis. [3–5] AGE is usually diagnosed clinically based on the clinical history, physical examination, and evaluation of risk factors. The diagnosis can be supported based on the results of a complete blood count (CBC), basic metabolic panel, stool analysis and culture, and serum electrolyte levels. [6,7]

Most AGE cases are caused by viruses. Bacterial AGE accounts for approximately 20% of cases. Viral gastroenteritis usually affects children aged

<2 years, whereas bacterial gastroenteritis occurs more often in those who are older than 2 years. However, diarrhea that lasts for at least 14 days is most frequently caused by parasitic infections. [8,9] Rotavirus is the most common cause of severe AGE in children and is responsible for an estimated 527 000 deaths in children younger than 5 years of age. [10] Although AGE usually remits spontaneously and only requires minimal medical intervention, severe complications, such as dehydration, may result in prolonged vomiting and diarrhea.

Dehydration occurs when the body loses a lot of water, electrolytes as well as other fluids; thus, an increase in serum osmolality of greater than 294 mOsm/kg is considered dehydration. [11] Dehydration is classified into three stages: mild, moderate, and severe. The assessment of the severity of dehydration in children is as follows: mild, 3–5% of dehydration; moderate, 6–9% of dehydration; and severe, more than 10% of dehydration. The percentage of dehydration is

determined through physical findings and vital signs. [12] In adults, mild dehydration is defined as a water loss equal to 1% of the body weight, which may result in headache, weakness, dizziness, and lethargy. Moderate dehydration can result in a dry mouth, low urinary output, tachycardia, and loss of skin elasticity. The body cannot regulate its temperature and blood pressure in case of severe dehydration. Patients with mild-to-moderate dehydration should receive oral rehydration therapy as first-line treatment. [13,14]

The aim of the present study was to assess the clinical profile of patients with acute kidney injury following acute gastroenteritis.

### Materials and Methods

The Present study was conducted in Department of General Medicine, Darbhanga Medical College and Hospital, Darbhanga, Bihar, India for 12 months. After applying inclusion and exclusion criteria, total 100 patients of AGE with AKI were considered for this study.

### Inclusion Criteria

- Patients above 18 years, admitted with acute gastroenteritis.
- Patients with acute kidney injury (AKI), diagnosed by KDIGO criteria.

### Exclusion Criteria

- Age less than 18 years.
- Patients with chronic renal insufficiency.
- Patients who are initially considered as AKI but subsequently found to be suffering from long-standing renal disease.

An informed consent was taken from patients for participation. Demographic and clinical details,

thorough examination findings were noted. Renal function tests with electrolytes are done daily and recorded. Other laboratory parameters such as CBC, ESR, Urine examination, Stool Examination, HIV, blood glucose, liver function test are also done. ABG, ECG, abdominal ultrasound examination were done whenever required. AKI was diagnosed according to Acute Kidney Injury Working Group of KDIGO (Kidney Disease: Improving Global Outcomes)<sup>2</sup> which diagnoses AKI by an absolute increase in Serum Creatinine, at least 0.3 mg/dL within 48 hours or by a 50% increase in Serum Creatinine from baseline within 7 days, or a urine volume of less than 0.5 mL/kg/h for at least 6 hours. Primary treatment was done with adequate fluid replacement and antibiotics (gram negative spectrum for ongoing gastroenteritis). Dialysis was done in patients with hyperkalemia, pulmonary edema and severe metabolic acidosis, who did not respond to medical treatment and prophylactically in patients whose creatinine is more than 4 mg/dl. Data recorded include patients' characteristics, comorbid medical conditions, dialysis requirement, total duration of hospital stay, and complications that occurred in their hospital stay were recorded. and final outcome. The clinical and laboratory parameters were analyzed to assess the role of each of these factors as the possible outcome i.e. recovery or death. All patients were followed up till discharge or death. Statistical analysis was done using descriptive statistics. Results were expressed as mean and standard deviation for continuous data and frequency as number and percentage.

### Results

**Table 1: Age distribution**

Age groups in years	Male	Female	Total
19-30	2	2	4 (4%)
31-40	4	5	9 (9%)
41-50	12	8	20 (20%)
51-60	15	8	23 (23%)
61-70	22	8	30 (30%)
71-80	8	3	11 (13%)
>80	2	1	3 (3%)
Total	65	35	100

Most common age group in this study was age group of 61-70 years (30%), followed by age group of 51-60 years (23%). Mean age of study patients was  $54.6 \pm 12.8$  years. Male patients (65%) were more than female patients (35%).

**Table 2: Symptom wise distribution**

Symptoms	N	%
Loose stools	100	100
Fever	78	78
Vomiting	70	70
Shortness of breath	20	20
Altered sensorium	16	16

According to clinical presentation most common symptom was loose stools (100%), followed by fever (78%) and vomiting (70%). Other complaints were shortness of breath (20%) and altered sensorium (16%).

**Table 3: AKI staging (KDIGO)**

Stage	Serum creatinine	Urine output	No. of patients (%)
1	1.5–1.9 times baseline OR	<0.5 ml/kg/h for 6–12 hours	58 (58%)
	≥0.3 mg/dl (≥26.5 μmol/l) increase		
2	2.0–2.9 times baseline	<0.5 ml/kg/h for ≥12 hours	31 (31%)
3	3.0 times baseline OR	<0.3 ml/kg/h for ≥24 hours	11 (11%)
	Increase in serum creatinine to ≥4.0 mg/dl (≥353.6 μmol/l) OR	OR	
	Initiation of renal replacement therapy OR	Anuria for ≥12 hours	
	In patients <18 years, decrease in eGFR to <35 ml/min per 1.73 m <sup>2</sup>		

AKI was staged for severity according to the KDIGO criteria. At the time of diagnosis most patients were in stage 1 (58%), while 31% and 11% were in stage 2 and 3 respectively.

### Discussion

Acute kidney injury (AKI) previously known as Acute Renal Failure, is a clinical syndrome characterized by an abrupt decline in glomerular filtration rate sufficient to decrease the elimination of nitrogenous waste products (urea and creatinine) and other uremic toxins. [15] The causes of AKI have traditionally been divided into three broad categories as prerenal (due to renal hypoperfusion), intrinsic (due to renal parenchymal pathology), postrenal (due to urinary tract obstruction). [16]

Most common age group in this study was age group of 61–70 years (30%), followed by age group of 51–60 years (23%). Mean age of study patients was 54.6 ± 12.8 years. Male patients (65%) were more than female patients (35%). Early in the course of AKI, optimization of the hemodynamic status and correction of any volume deficit will have a salutary effect on kidney function. Diuretics do not have any significant effect on progression or outcome of AKI. [17] In that study, mean age of presentation 46.5 ± 11.35 years, in males was 47.3 ± 11.35 and that of females was 45.72 ± 10.23 years. 63.33 % were Males and 36.7% were females. Similar findings were noted in present study. Acute kidney injury (AKI) is not an individual disease entity, but rather, is a syndrome that is characterized by a group of symptoms that occur together. AKI is a consequence of several underlying conditions; 20% of hospitalized patients have AKI, and 10% of these patients require renal replacement therapy. [18,19] According to the latest Kidney Disease Improving Global Outcomes (KDIGO) clinical guideline for the classification and management of AKI, an increase in serum creatinine (SCr) by ≥0.3 mg/dL within 48 hours and to ≥1.5 times the baseline level over 7 days or decreased urinary output <0.5 mL/kg/h for 6–12 hours indicates AKI. [20]

According to clinical presentation most common symptom was loose stools (100%), followed by fever (78%) and vomiting (70%). Other complaints were shortness of breath (20%) and altered sensorium (16%). AKI was staged for severity according to the KDIGO criteria. At the time of diagnosis most patients were in stage 1 (58%), while 31% and 11% were in stage 2 and 3 respectively. In the study by Satri V et al [21] sepsis was found to be the most common cause of AKI followed by acute gastroenteritis, while Eswarappa M et al [22] noted acute gastroenteritis as most common cause of AKI in their study. On stool culture and sensitivity, we found 88.5% patients with sterile stool cultures. 10.3% patients had E. coli on culture while 1 patient has Salmonella growth on stool culture.

Inbanathan J et al [23] noted that 70 % patients were treated conservatively and 30 % required hemodialysis. 52% were discharged within 1 week, while 37% of patients were discharged between 1-3 weeks. 11% of patients were treated for more than 3 weeks. 4% mortality was noted. The prognosis of patients with AKI is directly related to cause of renal failure and, to great extent, to the duration of renal failure before therapeutic intervention. Prognostic factors are older age, multiorgan failure (i.e., the more the organ that fails, the worse is prognosis), circulatory failure, vasopressor support, and need for renal replacement therapy (RRT). Factors influencing patient survival in acute kidney injury are severity of injury and underlying disease and other factors such as age, severity of coexistent illnesses and associated complications like Intravascular overload, hyperkalemia and other metabolic complications and systemic life-threatening complications like cardiac arrhythmia, myocardial infarction, pulmonary embolism, gastrointestinal ulcers, seizures, coma, hemolysis, bleeding tendencies and severe infections. [24]

In the appropriate context, early detection requires a high degree of suspicion that AKI is occurring. Diagnosis requires a combination of a clinical history, a thorough physical examination, an

accurate assessment of kidney function, appropriate imaging, and when indicated, a kidney biopsy.

### Conclusion

Acute kidney injury in patients with acute gastroenteritis had good prognosis if detected earlier. Early recognition of AKI is essential to ensure prompt and appropriate management, and to avoid progression to deadlier stages of the disease.

### References

1. Hartman S, Brown E, Loomis E, Russell HA. Gastroenteritis in children. *Am Fam Physician*. 2019;99(3):159–65.
2. Hall AJ, Rosenthal M, Gregoricus N, Greene SA, Ferguson J, Henao OL, et al... Incidence of acute gastroenteritis and role of Norovirus, Georgia, USA, 2004-2005. *Emerg Infect Dis*. 2011;17(8):1381–8.
3. Shane AL, Mody RK, Crump JA, Tarr PI, Steiner TS, Kotloff K, et al... 2017 Infectious Diseases Society of America clinical practice guidelines for the diagnosis and management of infectious diarrhea. *Clin Infect Dis*. 2017; 65(12): e45–e80.
4. Zollner-Schwetz I, Krause R. Therapy of acute gastroenteritis: role of antibiotics. *Clin Microbiol Infect*. 2015;21(8):744–9.
5. Bányai K, Estes MK, Martella V, Parashar UD. Viral gastroenteritis. *Lancet*. 2018;392 (10142):175–86.
6. DuPont HL. Acute infectious diarrhea in immunocompetent adults. *N Engl J Med*. 2014;370(16):1532–40.
7. Connor BA. Travelers' health - travelers' diarrhea; Updated 2015.
8. European Society for Paediatric Gastroenterology, Hepatology, and Nutrition/European Society for Paediatric Infectious Diseases. European Society for Paediatric Gastroenterology, Hepatology, and Nutrition/European Society for Paediatric Infectious Diseases Evidence-based guidelines for the management of acute gastroenteritis in children in Europe. *J Pediatr Gastroenterol Nutr*. 2008;46(Suppl 2) (Suppl 2): S81–S122.
9. Managing acute gastroenteritis Among children: oral rehydration, maintenance, and nutritional therapy. 2004;114(2):507-.
10. Telmesani AM. Oral Rehydration Salts, zinc supplement and rota virus vaccine in the management of childhood acute diarrhea. *J Family Community Med*. 2010;17(2):79–82.
11. Aylor K, Jones EB. Adult dehydration. In: StatPearls [Internet]. Updated 2022. Treasure Island (Florida): StatPearls Publishing; 2022.
12. Chen J, Wan CM, Gong ST, Fang F, Sun M, Qian Y, et al... Chinese clinical practice guidelines for acute infectious diarrhea in children. *World J Pediatr*. 2018;14(5):429–36.
13. American Medical Directors Association. Dehydration and fluid maintenance in the long-term care setting. Columbia, Maryland: Author; 2009.
14. Miller HJ. Dehydration in the older adult. *J Gerontol Nurs*. 2015;41(9):8–13.
15. Susantitaphong P, Cruz DN, Cerda J, Abulfaraj M, Alqahtani F, Koulouridis I, Jaber BL. World incidence of AKI: a meta-analysis. *Clinical journal of the American Society of Nephrology*. 2013 Sep 1;8(9):1482-93.
16. Kellum JA, Lameire N, Aspelin P, Barsoum RS, Burdmann EA, Goldstein SL, Herzog CA, Joannidis M, Kribben A, Levey AS, MacLeod AM. Kidney disease: improving global outcomes (KDIGO) acute kidney injury work group. KDIGO clinical practice guideline for acute kidney injury. *Kidney international supplements*. 2012 Mar;2(1):1-38.
17. Ho KM, Sheridan DJ. Meta-analysis of frusemide to prevent or treat acute renal failure. *Bmj*. 2006 Aug 24;333(7565):420.
18. Levey AS, James MT. Acute kidney injury. *Ann Intern Med*. 2017;167(9), ITC66–ITC80.
19. Bellomo R, Kellum JA, Ronco C. Acute kidney injury. *Lancet*. 2012;380(9843):756–66.
20. Ricci Z, Romagnoli S. Acute kidney injury: diagnosis and classification in adults and children. *Contrib Nephrol*. 2018; 193:1–12.
21. Satri V, Kumar VS, Satyanarayana V, Ramakrishna BS, Rao PS, Madhusudan M. Epidemiology and outcome of acute kidney injury in patients presenting to emergency department—Our experience. *Journal of Clinical and Scientific Research*. 2020 Apr 1;9(2):77-81.
22. Eswarappa M, Gireesh MS, Ravi V, Kumar D, Dev G. Spectrum of acute kidney injury in critically ill patients: A single center study from South India. *Indian journal of nephrology*. 2014 Sep 1;24(5):280-5.
23. Inbanathan J, Lavanya BU. Clinical Profile of Renal Involvement in Acute Gastroenteritis Patients. *Int J Sci Stud* 2016;4(8):48-52.
24. Sean M. Bagshaw .Short- and long-term survival after acute kidney injury *Nephrol. Dial. Transplant*. 2008; 23:23-29.