

**A Study to Evaluate the Circulatory Levels of Hs-CRP in Patients of Non-Alcoholic Fatty Liver Disease: A Comparative Study**Gangesh Kumar Gunjan<sup>1</sup>, Umesh Chandra Jha<sup>2</sup><sup>1</sup>Senior Resident, Department of General Medicine, Darbhanga Medical College and Hospital, Darbhanga, Bihar, India<sup>2</sup>Associate Professor and HOD, Department of General Medicine, Darbhanga Medical College and Hospital, Darbhanga, Bihar, India

Received: 04-01-2024 / Revised: 10-02-2024 / Accepted: 24-02-2024

Corresponding Author: Dr. Gangesh Kumar Gunjan

Conflict of interest: Nil

**Abstract****Aim:** The aim of the present study was to evaluate the circulatory levels of Hs-CRP in patients of non-alcoholic fatty liver disease and compare them with clinically healthy controls.**Material & Methods:** The study included 100 NAFLD patients and 100 clinically healthy controls. This study was carried out at Darbhanga Medical College and Hospital, Darbhanga, Bihar, India for one year. The study information was given to all the study participants and their written consent was collected.**Results:** The lipid profile variables like TG and LDL were high patients than controls. The present study observed a significant increase anthropometric parameters and lipid parameters in patients than controls. The inflammatory marker C-reactive protein levels increased in NASH group than NAFLD.**Conclusion:** The present study concluded that Hs-CRP can be considered as an important marker of inflammatory status in NAFLD and NASH.**Keywords:** Inflammatory Markers, NAFLD, NASH, Hs-CRP.

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

**Introduction**

Non-alcoholic fatty liver disease (NAFLD) encompasses a spectrum of clinic-pathologic entities that have in common the presence of fat accumulation in the liver in the absence of significant alcohol consumption. [1,2] It is characterized by triglyceride accumulation in hepatocytes, which occurs without alcohol abuse. It is one of the most commonly encountered chronic liver diseases. [3] The clinical spectrum of NAFLD ranges from asymptomatic steatosis to steatohepatitis, fibrosis and cirrhosis. Most NAFLD patients have asymptomatic simple steatosis without adverse sequelae. [4] Inflammation is present in the early phase of NAFLD and is an essential driver in the initiation and progression of liver damage. [5]

NAFLD is strongly associated with the major components of the metabolic syndrome. Increasing recognition of the importance of NAFLD and its strong relationship with the metabolic syndrome has stimulated an interest in the possible role of NAFLD in the development of cardiovascular disease (CVD). Mortality is even higher amongst individuals with non-alcoholic steatohepatitis

(NASH), and this mortality was primarily cardiovascular, rather than liver related. [6]

Liver is the center for the production of classical biomarkers of inflammation and endothelial dysfunction. Their secretion is partly dependent on factors that are up-regulated in the presence of insulin resistance and the metabolic syndrome. A series of proinflammatory proteins and cytokines are implicated in hepatic inflammation have been studied to test their usefulness as non-invasive soluble biomarkers for NAFLD/NASH diagnosis and prognosis. [7]

IL-6 and TNF- $\alpha$  are the major stimuli responsible for increased hepatic production of C-reactive protein (CRP), fibrinogen and other acute-phase proteins. [8,9] HsCRP is a potent indicator of subclinical inflammation and plays a crucial role in various inflammatory cascades and may lead to cardiovascular diseases. [10] It is synthesized mainly in liver, however it is also produced at many other sources. CRP and HsCRP are the two names given to the same protein. CRP is measured in a broader range as it is a non-specific marker of inflammation and increases in many inflammatory conditions. [10,11] Correlations of C-reactive

protein levels with anthropometric profile, percentage of body fat and lipids in healthy adolescents and young adults has been observed in young Indian population. [11] NAFLD is a classical disease due to abnormal deposition of fat in liver and may develop some inflammatory condition which may lead to NASH. [12] HsCRP is one of the inflammatory molecules identified and shown to be associated with development of NASH. HsCRP has been identified as a culprit in vulnerable atherosclerotic plaques and is associated with ACS. [13]

Hence the aim of this study was to evaluate the circulatory levels of HsCRP in patients of non-alcoholic fatty liver.

### Material & Methods

The study included 100 NAFLD patients and 100 clinically healthy controls. This study was carried out at Darbhanga Medical College and Hospital, Darbhanga, Bihar, India for one year. The study information was given to all the study participants and their written consent was collected.

### Inclusion Criteria:

Clinically proven NAFLD patients (NAFLD was detected by ultrasound), Age and sex matched healthy individuals control group.

### Exclusion Criteria:

Patient with significant alcohol intake (>20 g/day) type 2 diabetes mellitus (T2DM), cardiovascular disease (CVD), presence of other liver diseases (alcoholic liver disease, viral hepatitis, autoimmune hepatitis, primary biliary cirrhosis, biliary obstruction, drug-induced liver damage etc.), severe end organ damage, human immunodeficiency virus infection, pregnancy and lactation, were excluded from the study

### Blood Sampling and Methodology:

Fasting venous blood samples were collected for biochemical investigations. Routine biochemical parameters like fasting blood sugar, lipid profile, liver function tests and Serum HsCRP levels were estimated in clinical laboratory of D.Y Patil hospital and research centre, Nerul, Navi Mumbai. Cardiac profile is also evaluated by ECG, ECHO findings as well angiographic results.

### STATISTICAL ANALYSIS

SPSS software (version 17) was used for Statistical analysis of the data.

### Results

**Table 1: Demographical Characteristics of study subjects**

Variables	Controls (n=100)	Patients (n=100)	P value
BMI	23.37 ( $\pm$ 3.87)	25.35 ( $\pm$ 5.65)	0.07
WC	92.8 ( $\pm$ 10.5)	100.22 ( $\pm$ 12.16)	0.07
WHR	0.95 ( $\pm$ 0.55)	1.07 ( $\pm$ 0.43)	0.01
TG (mg%)	110.68 ( $\pm$ 43.21)	158.22 ( $\pm$ 80.11)	0.06
TC (mg%)	172.24 ( $\pm$ 30.12)	212.38 ( $\pm$ 47.63)	0.02
LDL (mg%)	99.30 ( $\pm$ 32.80)	120.16 ( $\pm$ 45.32)	0.07
ALT(IU/L)	75.5 $\pm$ 10.5	17.3 $\pm$ 3.4	0.01

The lipid profile variables like TG and LDL were high patients than controls.

**Table 2: Serum HsCRP**

Variables	NAFLD	NASH	P value
Serum HsCRP	2.12 ( $\pm$ 0.18)	3.43 ( $\pm$ 0.57)	0.07

The present study observed a significant increase anthropometric parameters and lipid parameters in patients than controls. The inflammatory marker C-reactive protein levels increased in NASH group than NAFLD.

### Discussion

Fatty liver is a clinical condition characterized by and commonly understood as the accumulation of lipids within hepatocytes. Traditionally, for practical purposes a, hepatic fat content exceeding 5% of the liver weight has been considered as fatty liver. [14] In the past, fatty liver was well-thought-out as a benign and reversible pathology and represented a nonspecific response of the liver to metabolic stress of various origin.<sup>2</sup>

However, it is increasingly seen as a part of the spectrum that can be as benign as fatty liver to a fatal condition of hepatocellular carcinoma (HCC). Alcohol consumption has been the most common factor responsible for fatty liver, however, it may also be seen in non-alcoholics and the clinical entity is known as non-alcoholic fatty liver disease (NAFLD). It is commonly associated with various components of metabolic syndrome including obesity, diabetes mellitus, and dyslipidemia. [15] Various reports also suggest that NAFLD may be associated with low-grade inflammation in liver. [16]

The lipid profile variables like TG and LDL were high patients than controls. The present study observed a significant increase anthropometric

parameters and lipid parameters in patients than controls. The inflammatory marker C-reactive protein levels increased in NASH group than NAFLD. High serum hs-CRP reflects its synthesis is in response to a pathological process<sup>9</sup>. Proinflammatory mediators like IL-6 and TNF- $\alpha$  plays an important role in production of HsCRP in liver. In vivo release of interleukin-6 (IL-6), linked closely to hs-CRP pathway. Hence IL-6 levels considered as an important indicator for subclinical inflammation which is broadly specific. Various studies have shown the association of IL-6 with path physiological conditions, like obesity, type 2 diabetes and CVD. [17,18] While the pathophysiology of NAFLD is not completely understood, accumulation of triglycerides in hepatocytes is due to various factors like- presence of oxidative stress, lipid peroxidation, pro-inflammatory cytokines (e.g. TNF- $\alpha$ , IL-6). [19]

Some studies also observed the adiponectin m-RNAs in liver biopsies of these patients. Further they have noted the appearance of the m-RNAs of adiponectin receptors also. [20] Animal studies showed that increased systemic levels of TNF- $\alpha$  is a result of high amount of fatty acids present in the liver which mediate hepatic production of TNF- $\alpha$ . [21] Hepatocyte damage facilitates activation of liver-specific macrophages ('Kupffer Cells') which secretes more TNF- $\alpha$  and IL-6 into the blood. [22] A recent follow-up study was done by Lee et al [23] to address whether hs-CRP levels within the normal range can predict the development of NAFLD in healthy male subjects. In the 7-years follow-up, it was found that the risk for NAFLD increased as the hs-CRP level increased ( $P < 0.001$ ). As the hs-CRP level increased within the healthy cohort, the risk of developing NAFLD increased.

### Conclusion

The present study concluded that Hs-CRP can be considered as an important marker of inflammatory status in NAFLD and NASH.

### References

- Farrell G.C., George J., Hall P., de la M., McCullough A.J. Blackwell Publishing; 2005. Fatty Liver Disease: NASH and Related Disorders.
- Farrell G.C., Larter C.Z. Nonalcoholic fatty liver disease: from steatosis to cirrhosis. *Hepato Balt Md*. 2006;43(2 Suppl 1): S99-S112.
- Angulo P. Nonalcoholic fatty liver disease. *N Engl J Med*. 2002; 346: 1221-1231.
- Musso G, Gambino R, Cassader M, Pagano G. Meta-analysis: natural history of non-alcoholic fatty liver disease (NAFLD) and diagnostic accuracy of non-invasive tests for liver disease severity. *Ann Med*. 2011; 43: 617-649.
- Tilg H, Moschen AR. Evolution of inflammation in non-alcoholic fatty liver disease: the multiple parallel hits hypothesis. *Hepatology*. 2010; 52:1836-46.
- Ekstedt M, Franzén LE, Mathiesen UL, et al. Long-term follow-up of patients with NAFLD and elevated liver enzymes. *Hepato Balt Md*. 2006; 44:865e873.
- Jamialahmadi T, Bo S, Abbasifard M, Sathyapalan T, Jangjoo A, Moallem SA, Almahmeed W, Ashari S, Johnston TP, Sahebkar A. Association of C-reactive protein with histological, elastographic, and sonographic indices of non-alcoholic fatty liver disease in individuals with severe obesity. *Journal of Health, Population and Nutrition*. 2023 Dec;42(1):1-9.
- Van Gaal LF, Mertens IL, De Block CE. Mechanisms linking obesity with cardiovascular disease. *Nature*. 2006; 444:875e880.
- Shoelson SE. Inflammation and insulin resistance. *J Clin Invest*. 2006; 116:1793e1801.
- Blake GJ, Ridker PM. Inflammatory biomarkers and cardiovascular risk prediction. *J Intern Med*. 2002;252: 283-94.
- Vikram NK, Misra A, Dwivedi M, Sharma R, Pandey RM, Luthra K, Chatterjee A, Dhingra V, Jaikhanani BL, Talwar KK, Guleria R. Correlations of C-reactive protein levels with anthropometric profile, percentage of body fat and lipids in healthy adolescents and young adults in urban North India. *Atherosclerosis*. 2003 Jun 1;168(2):305-13.
- Tarantino G, Conca P, Pasanisi F, et al. could inflammatory markers help diagnose nonalcoholic steatohepatitis. *Eur J Gastroenterol Hepatol*. 2009; 21:504-11.
- Deepali Vidhate, James Thomas, Becky Thomas, Simran Dhadijala, Arpit Murarka. Association of high sensitivity C-reactive protein (hs-CRP) with unstable angina: An alarm for vulnerable plaques in CAD patients. *International Journal of Clinical Biochemistry and Research*, July-September, 2018;5(3):458-463.
- Sherlock S, Dooley J, Iber F. Diseases of the liver and biliary system. Oxford: Blackwell science; 2002 Jan.
- El-Serag HB, Tran T, Everhart JE. Diabetes increases the risk of chronic liver disease and hepatocellular carcinoma. *Gastroenterology*. 2004 Feb 1;126(2):460-8.
- Rodríguez-Hernández H, Simental-Mendía LE, Rodríguez-Ramírez G, Reyes-Romero MA. Obesity and inflammation: epidemiology, risk factors, and markers of inflammation. *International journal of endocrinology*. 2013 Oct; 2013.

17. Vidhate D, Thomas J, Gupte A. Association of circulating levels of Adiponectin and IL-6 with incidence and severity of the coronary artery disease in patients undergoing Coronary Artery Bypass Graft (CABG). *International Journal*. 2021 Mar;4(2):215.
18. Vidhate D, Thomas J, Thomas B, Dhadijala S, Murarka A. Association of high sensitivity C-reactive protein (hs-CRP) with unstable angina: An alarm for vulnerable plaques in CAD patients.
19. Duvnjak M, Lerotić I, Baršić N, Tomašić V, Jukić LV, Velagić V. Pathogenesis and management issues for non-alcoholic fatty liver disease. *World journal of gastroenterology: WJG*. 2007 Sep 9;13(34):4539.
20. Kaser S, Moschen A, Cayon A, Kaser A, Crespo J, Pons-Romero F, Ebenbichler CF, Patsch JR, Tilg H. Adiponectin and its receptors in non-alcoholic steatohepatitis. *Gut*. 2005 Jan;54(1):117.
21. Feldstein AE, Werneburg NW, Canbay A, Guicciardi ME, Bronk SF, Rydzewski R, Burgart LJ, Gores GJ. Free fatty acids promote hepatic lipotoxicity by stimulating TNF- $\alpha$  expression via a lysosomal pathway. *Hepatology*. 2004 Jul;40(1):185-94.
22. Wieckowska A, Papouchado BG, Li Z, Lopez R, Zein NN, Feldstein AE. Increased hepatic and circulating interleukin-6 levels in human nonalcoholic steatohepatitis. *Official journal of the American College of Gastroenterology ACG*. 2008 Jun 1;103(6):1372-9.
23. Lee J, Yoon K, Ryu S, Chang Y, Kim HR. High-normal levels of hs-CRP predict the development of non-alcoholic fatty liver in healthy men. *PLoS One*. 2017 Feb 24;12(2):e0172666.